

# Geometric Lang-Vojta's conjecture in $\mathbb{P}^2$

Amos Turchet

Università degli studi di Udine

On Lang and Vojta's conjectures

Centre International de Rencontres Mathématiques

March 6th 2014



# Table of contents

- 1 Hyperbolicity
- 2 Geometric Lang-Vojta's conjecture
- 3 Logarithmic Geometry



# Basic Definition

## Definition

A complex manifold  $M$  is said to be (Brody)-**hyperbolic** if every holomorphic map  $f : \mathbb{C} \rightarrow M$  is constant.



# Basic Definition

## Definition

A complex manifold  $M$  is said to be (Brody)-**hyperbolic** if every holomorphic map  $f : \mathbb{C} \rightarrow M$  is constant.

Brody Theorem implies that for compact manifolds *Brody-hyperbolicity* is equivalent to being hyperbolic in the sense of *Kobayashi*.



# Basic Definition

## Definition

A complex manifold  $M$  is said to be (Brody)-**hyperbolic** if every holomorphic map  $f : \mathbb{C} \rightarrow M$  is constant.

Brody Theorem implies that for compact manifolds *Brody-hyperbolicity* is equivalent to being hyperbolic in the sense of *Kobayashi*.

## Fundamental problem

Describe geometric attributes of  $X$  that implies hyperbolicity.



# Hyperbolicity for RS

In dimension 1, i.e. for Riemann surfaces, the following easy fact holds:

## Theorem

*If  $X$  is a Riemann surface of genus at least 2, every holomorphic map  $\mathbb{C} \rightarrow X$  is constant. Therefore  $X$  is hyperbolic.*



# Hyperbolicity for RS

In dimension 1, i.e. for Riemann surfaces, the following easy fact holds:

## Theorem

*If  $X$  is a Riemann surface of genus at least 2, every holomorphic map  $\mathbb{C} \rightarrow X$  is constant. Therefore  $X$  is hyperbolic.*

The theorem follows essentially from Liouville's Theorem.



# Hyperbolicity for RS

In dimension 1, i.e. for Riemann surfaces, the following easy fact holds:

## Theorem

*If  $X$  is a Riemann surface of genus at least 2, every holomorphic map  $\mathbb{C} \rightarrow X$  is constant. Therefore  $X$  is hyperbolic.*

The theorem follows essentially from Liouville's Theorem. In particular the geometric condition *genus*  $\geq 2$  implies hyperbolicity.





# Hyperbolicity for affine curves

There is an analogous results for affine curves, i.e. complements of a finite set of points in a RS.

## Theorem

*If  $X$  is a Riemann surface of genus  $g(X)$ , and  $S$  a finite set of points of  $X$ . Then*

$$X \setminus S \text{ is hyperbolic} \iff 2g(X) - 2 + \#S > 0$$



# Hyperbolicity for affine curves

There is an analogous results for affine curves, i.e. complements of a finite set of points in a RS.

## Theorem

*If  $X$  is a Riemann surface of genus  $g(X)$ , and  $S$  a finite set of points of  $X$ . Then*

$$X \setminus S \text{ is hyperbolic} \iff 2g(X) - 2 + \#S > 0$$

Follows essentially from Picard's Little Theorem.



# Hyperbolicity for affine curves

There is an analogous results for affine curves, i.e. complements of a finite set of points in a RS.

## Theorem

*If  $X$  is a Riemann surface of genus  $g(X)$ , and  $S$  a finite set of points of  $X$ . Then*

$$X \setminus S \text{ is hyperbolic} \iff 2g(X) - 2 + \#S > 0$$

Follows essentially from Picard's Little Theorem. Notice how the geometric condition  $2g(X) - 2 + \#S > 0$  now encodes information about the points at infinity of  $X$ .



# Hyperbolicity for affine curves

There is an analogous results for affine curves, i.e. complements of a finite set of points in a RS.

## Theorem

*If  $X$  is a Riemann surface of genus  $g(X)$ , and  $S$  a finite set of points of  $X$ . Then*

$$X \setminus S \text{ is hyperbolic} \iff 2g(X) - 2 + \#S > 0$$

Follows essentially from Picard's Little Theorem. Notice how the geometric condition  $2g(X) - 2 + \#S > 0$  now encodes information about the points at infinity of  $X$ .

Q: What happens in higher dimensions?



# Hyperbolicity in higher dimensions

Complete geometric description of hyperbolic manifolds in dimension greater than one is still out of reach.



# Hyperbolicity in higher dimensions

Complete geometric description of hyperbolic manifolds in dimension greater than one is still out of reach.

Kobayashi proposed a famous conjecture for manifold embedded in projective space

## Conjecture (Kobayashi)

- (a) *a very general hypersurface  $D \subset \mathbb{P}^{n+1}$  with  $n \geq 2$  of degree  $\deg D \geq 2n + 1$  is hyperbolic;*
- (b)  *$\mathbb{P}^n \setminus D$  with  $n \geq 2$  is hyperbolic for a very general hypersurface  $D \subset \mathbb{P}^n$  of degree  $\deg D \geq 2n + 1$ .*



# Hyperbolicity in higher dimensions

Complete geometric description of hyperbolic manifolds in dimension greater than one is still out of reach.

Kobayashi proposed a famous conjecture for manifold embedded in projective space

## Conjecture (Kobayashi)

- (a) *a very general hypersurface  $D \subset \mathbb{P}^{n+1}$  with  $n \geq 2$  of degree  $\deg D \geq 2n + 1$  is hyperbolic;*
- (b)  *$\mathbb{P}^n \setminus D$  with  $n \geq 2$  is hyperbolic for a very general hypersurface  $D \subset \mathbb{P}^n$  of degree  $\deg D \geq 2n + 1$ .*

Results in many important cases: works by Y.T. Siu, Demailly, El Goul, McQuillan, Diverio, Merker, Rousseau, Pacienza...



# Hyperbolicity in higher dimensions

Problems:

- Achieve the conjecture's bound is really hard;
- Full description of hyperbolic manifold not embedded in  $\mathbb{P}^n$  seems very difficult to discuss;





# Hyperbolicity in higher dimensions

Problems:

- Achieve the conjecture's bound is really hard;
- Full description of hyperbolic manifold not embedded in  $\mathbb{P}^n$  seems very difficult to discuss;

**Strategy:** Lower the expectation.



# Hyperbolicity for Algebraic Varieties

Instead of complex manifolds we focus on algebraic projective varieties.



# Hyperbolicity for Algebraic Varieties

Instead of complex manifolds we focus on algebraic projective varieties.

## Theorem (Demailly)

*Let  $X$  be a compact complex hyperbolic **algebraic** variety. Then there exists an  $\epsilon > 0$  such that every compact irreducible curve  $\mathcal{C} \subset X$  satisfies:*

$$(\star) \quad -\chi(\tilde{\mathcal{C}}) = 2g(\tilde{\mathcal{C}}) - 2 \geq \epsilon \deg \tilde{\mathcal{C}}$$

*where  $\tilde{\mathcal{C}}$  is the normalization of the curve  $\mathcal{C}$ , and the degree is calculated respect to an ample divisor in  $X$ .*



# Hyperbolicity for Algebraic Varieties

Instead of complex manifolds we focus on algebraic projective varieties.

## Theorem (Demailly)

Let  $X$  be a compact complex hyperbolic *algebraic* variety. Then there exists an  $\epsilon > 0$  such that every compact irreducible curve  $\mathcal{C} \subset X$  satisfies:

$$(\star) \quad -\chi(\tilde{\mathcal{C}}) = 2g(\tilde{\mathcal{C}}) - 2 \geq \epsilon \deg \tilde{\mathcal{C}}$$

where  $\tilde{\mathcal{C}}$  is the normalization of the curve  $\mathcal{C}$ , and the degree is calculated respect to an ample divisor in  $X$ .

Property  $(\star)$  is defined in a pure algebraic way.



# Algebraic Hyperbolicity

## Definition (Demailly)

A projective algebraic variety  $X$  is said to be *algebraically hyperbolic* if  $(\star)$  holds.



# Algebraic Hyperbolicity

## Definition (Demailly)

A projective algebraic variety  $X$  is said to be *algebraically hyperbolic* if  $(\star)$  holds.

Hyperbolicity implies algebraic hyperbolicity; hence alg. hyperbolicity can be used to test whether a projective variety can be hyperbolic.



# Algebraic Hyperbolicity

## Definition (Demailly)

A projective algebraic variety  $X$  is said to be *algebraically hyperbolic* if  $(\star)$  holds.

Hyperbolicity implies algebraic hyperbolicity; hence alg. hyperbolicity can be used to test whether a projective variety can be hyperbolic.

Algebraic Hyperbolicity is directly related to Vojta conjectures.



# Algebraic Hyperbolicity for affine varieties

## Definition

Given a projective algebraic variety  $X$  and a normal crossing divisor  $D$ ,  $X \setminus D$  is said to be *algebraically hyperbolic* if there exists an  $\epsilon > 0$  such that every compact irreducible curve  $\mathcal{C} \subset X$  satisfies:

$$-\chi(\tilde{\mathcal{C}}) = 2g(\tilde{\mathcal{C}}) - 2 + \#S \geq \epsilon \deg \tilde{\mathcal{C}}$$

where  $S$  is the set  $\nu^{-1}(D)$  and  $\nu : \tilde{\mathcal{C}} \rightarrow \mathcal{C}$  is the normalization.





# Algebraic Hyperbolicity for affine varieties

## Definition

Given a projective algebraic variety  $X$  and a normal crossing divisor  $D$ ,  $X \setminus D$  is said to be *algebraically hyperbolic* if there exists an  $\epsilon > 0$  such that every compact irreducible curve  $\tilde{\mathcal{C}} \subset X$  satisfies:

$$-\chi(\tilde{\mathcal{C}}) = 2g(\tilde{\mathcal{C}}) - 2 + \#S \geq \epsilon \deg \tilde{\mathcal{C}}$$

where  $S$  is the set  $\nu^{-1}(D)$  and  $\nu : \tilde{\mathcal{C}} \rightarrow \mathcal{C}$  is the normalization.

We call  $X \setminus D$  **weakly algebraically hyperbolic** if the previous bound reads

$$\max\{1, 2g(\tilde{\mathcal{C}}) - 2 + \#S\} \geq \epsilon \deg \tilde{\mathcal{C}}$$



# The conjecture

## Conjecture (Geometric Lang-Vojta)

Let  $\tilde{X}$  be a smooth projective algebraic surface with canonical divisor  $K_{\tilde{X}}$  and let  $D$  be a reduced normal crossing divisor on  $\tilde{X}$ . Let  $X = \tilde{X} \setminus D$  be the complement of the support of  $D$ . *If  $X$  is of log-general type, i.e. if  $D + K_{\tilde{X}}$  is big, then  $X$  is weakly algebraically hyperbolic.*



# The conjecture

## Conjecture (Geometric Lang-Vojta)

Let  $\tilde{X}$  be a smooth projective algebraic surface with canonical divisor  $K_{\tilde{X}}$  and let  $D$  be a reduced normal crossing divisor on  $\tilde{X}$ . Let  $X = \tilde{X} \setminus D$  be the complement of the support of  $D$ . *If  $X$  is of log-general type, i.e. if  $D + K_{\tilde{X}}$  is big, then  $X$  is weakly algebraically hyperbolic.*

Previous conjecture can be seen as a geometric version of the famous Lang-Vojta's conjecture predicting degeneracy of integral points on log-general type surfaces.



# The conjecture

## Conjecture (Geometric Lang-Vojta)


Let  $\tilde{X}$  be a smooth projective algebraic surface with canonical divisor  $K_{\tilde{X}}$  and let  $D$  be a reduced normal crossing divisor on  $\tilde{X}$ . Let  $X = \tilde{X} \setminus D$  be the complement of the support of  $D$ . *If  $X$  is of log-general type, i.e. if  $D + K_{\tilde{X}}$  is big, then  $X$  is weakly algebraically hyperbolic.*

Previous conjecture can be seen as a geometric version of the famous Lang-Vojta's conjecture predicting degeneracy of integral points on log-general type surfaces.

The “weakly” cannot be suppressed: there are examples of log general type surface which contains  $\mathbb{G}_m$



# Log general type surface, not AH



# The case of $\mathbb{P}^2$

When  $X = \mathbb{P}^2$  previous conjecture takes the form:

## Conjecture

*Let  $D$  be a reduced plane curve with normal crossing and let  $X = \mathbb{P}^2 \setminus D$ . If  $\deg D \geq 4$ , then  $X$  is weakly algebraically hyperbolic.*



# The case of $\mathbb{P}^2$

When  $X = \mathbb{P}^2$  previous conjecture takes the form:

## Conjecture

*Let  $D$  be a reduced plane curve with normal crossing and let  $X = \mathbb{P}^2 \setminus D$ . If  $\deg D \geq 4$ , then  $X$  is weakly algebraically hyperbolic.*

As in example before: in the case  $\deg D = 4$  the complement  $X$  is not (algebraic) hyperbolic!



# Known results

Lang-Vojta Conjecture for  $\mathbb{P}^2$  is known for complements of very general curves of degree at least 5 (Xi Chen, Pacienza and Rousseau).





# Known results

Lang-Vojta Conjecture for  $\mathbb{P}^2$  is known for complements of very general curves of degree at least 5 (Xi Chen, Pacienza and Rousseau).

When the degree of  $D$  is 4 the conjecture is known for **generic**  $D$  having at least three components:

- The four line case follows from an extension of Mason's ABC theorem (Brownawell and Masser);
- The three component case can be reduced to a  $S$ -unit  $gcd$  problem (Corvaja and Zannier).



# Known results

Lang-Vojta Conjecture for  $\mathbb{P}^2$  is known for complements of very general curves of degree at least 5 (Xi Chen, Pacienza and Rousseau).

When the degree of  $D$  is 4 the conjecture is known for **generic**  $D$  having at least three components:

- The four line case follows from an extension of Mason's ABC theorem (Brownawell and Masser);
- The three component case can be reduced to a  $S$ -unit  $gcd$  problem (Corvaja and Zannier).

All these results have structural obstruction for extensions to the remaining cases, namely  $D$  with less than three irreducible components and  $\deg D = 4$ .



# Goal and ideas

**Goal:** prove Geometric LV for the complement in  $\mathbb{P}^2$  of a (very) generic quartic (even irreducible!).



# Goal and ideas

**Goal:** prove Geometric LV for the complement in  $\mathbb{P}^2$  of a (very) generic quartic (even irreducible!).

**Idea:** consider the three component case as a deformation of the irreducible case.



# Goal and ideas

**Goal:** prove Geometric LV for the complement in  $\mathbb{P}^2$  of a (very) generic quartic (even irreducible!).

**Idea:** consider the three component case as a deformation of the irreducible case.

A similar idea is used in Xi Chen's proof of LV Conjecture: he degenerate the boundary  $D$  to a union of hyperplanes and then applying the known results for  $\mathbb{P}^2 \setminus \bigcup_{i=1,5} H_i$ .

His argument is involved and requires the degree of  $D$  to be at least 5 to work.



# Goal and ideas

**Goal:** prove Geometric LV for the complement in  $\mathbb{P}^2$  of a (very) generic quartic (even irreducible!).

Idea: consider the three component case as a deformation of the irreducible case.




A similar idea is used in Xi Chen's proof of LV Conjecture: he degenerate the boundary  $D$  to a union of hyperplanes and then applying the known results for  $\mathbb{P}^2 \setminus \bigcup_{i=1,5} H_i$ .

His argument is involved and requires the degree of  $D$  to be at least 5 to work.

In our case there are problems that need to be addressed.



# An example



# Problems

- No control on the variations of the number of points of intersection.





# Problems

- No control on the variations of the number of points of intersection.
- Curves can become reducible.



# Problems

- No control on the variations of the number of points of intersection.
- Curves can become reducible.
- A component of the curve could coincide with an irreducible component of the divisor.



# Problems

- No control on the variations of the number of points of intersection.
- Curves can become reducible.
- A component of the curve could coincide with an irreducible component of the divisor.

**Solution:** use logarithmic Geometry (take care of multiplicities of intersection).



# Main result

Theorem (T. - WAH for complements of quartics)

$\mathbb{P}^2 \setminus D$  is weakly algebraic hyperbolic for every simple normal crossing divisor  $D$  of degree 4 which flattly and log smoothly deforms to a conic and two lines.



# Main result

## Theorem (T. - WAH for complements of quartics)

$\mathbb{P}^2 \setminus D$  is weakly algebraic hyperbolic for every simple normal crossing divisor  $D$  of degree 4 which flattly and log smoothly deforms to a conic and two lines.

### Restrictions:

- Need to consider only **simple** normal crossing divisors (possible extensions are announced).



# Main result

## Theorem (T. - WAH for complements of quartics)

$\mathbb{P}^2 \setminus D$  is weakly algebraic hyperbolic for every simple normal crossing divisor  $D$  of degree 4 which flattly and log smoothly deforms to a conic and two lines.

### Restrictions:

- Need to consider only **simple** normal crossing divisors (possible extensions are announced).
- Deformations need to be **logarithmically smooth** which is stronger than flat.



# Main result

## Theorem (T. - WAH for complements of quartics)

$\mathbb{P}^2 \setminus D$  is weakly algebraic hyperbolic for every simple normal crossing divisor  $D$  of degree 4 which flattly and log smoothly deforms to a conic and two lines.

### Restrictions:

- Need to consider only **simple** normal crossing divisors (possible extensions are announced).
- Deformations need to be **logarithmically smooth** which is stronger than flat.
- Argument requires to work in logarithmic category.



# Sketch of the argument

We start from Corvaja and Zannier's result (CZ in the sequel) for the complement of a conic and two lines that read as follows

**Theorem (Corvaja and Zannier, 2008)**

*Let  $X = \mathbb{P}^2 \setminus D$ , where  $D$  is a quartic consisting of the union of a smooth conic and two lines in general position. Let  $\tilde{C}$  be a smooth complete algebraic curve and  $S \subset \tilde{C}$  a finite set of points. Then for every morphism  $f : \tilde{C} \rightarrow \mathbb{P}^2$  such that  $f^{-1}(D) \subset S$  the following holds:*

$$\deg f(\tilde{C}) \leq 2^{15} \cdot 35 \cdot \max\{1, \chi_S(\tilde{C})\}$$





# Sketch of the argument

We start from Corvaja and Zannier's result (CZ in the sequel) for the complement of a conic and two lines that read as follows

## Theorem (Corvaja and Zannier, 2008)

*Let  $X = \mathbb{P}^2 \setminus D$ , where  $D$  is a quartic consisting of the union of a smooth conic and two lines in general position. Let  $\tilde{C}$  be a smooth complete algebraic curve and  $S \subset \tilde{C}$  a finite set of points. Then for every morphism  $f : \tilde{C} \rightarrow \mathbb{P}^2$  such that  $f^{-1}(D) \subset S$  the following holds:*

$$\deg f(\tilde{C}) \leq 2^{15} \cdot 35 \cdot \max\{1, \chi_S(\tilde{C})\}$$

The main idea is to reformulate this results in terms of vanishing of certain moduli spaces of log stable maps  $\mathcal{K}_\Gamma(\mathbb{P}^2, D)$ .



# Sketch of the argument

We start from Corvaja and Zannier's result (CZ in the sequel) for the complement of a conic and two lines that read as follows

**Theorem (Corvaja and Zannier, 2008)**

*Let  $X = \mathbb{P}^2 \setminus D$ , where  $D$  is a quartic consisting of the union of a smooth conic and two lines in general position. Let  $\tilde{C}$  be a smooth complete algebraic curve and  $S \subset \tilde{C}$  a finite set of points. Then for every morphism  $f : \tilde{C} \rightarrow \mathbb{P}^2$  such that  $f^{-1}(D) \subset S$  the following holds:*

$$\deg f(\tilde{C}) \leq 2^{15} \cdot 35 \cdot \max\{1, \chi_S(\tilde{C})\}$$

The main idea is to reformulate this results in terms of vanishing of certain moduli spaces of log stable maps  $\mathcal{K}_\Gamma(\mathbb{P}^2, D)$ .

Then use the properness of the stack  $\mathcal{K}_\Gamma(\mathbb{P}^2, D)$  to extend the result for very general  $D$  deforming flatly to a conic and two lines.



# Why Logarithmic Geometry

- If in the deformation argument multiplicities of intersection are fixed,  $S$  can be controlled.



# Why Logarithmic Geometry

- If in the deformation argument multiplicities of intersection are fixed,  $S$  can be controlled.
- Log Geometry gives a way to define point of intersection even for irreducible components coinciding with a component of the divisor.



# Why Logarithmic Geometry

- If in the deformation argument multiplicities of intersection are fixed,  $S$  can be controlled.
- Log Geometry gives a way to define point of intersection even for irreducible components coinciding with a component of the divisor.
- Each curve  $\mathcal{C}$  has a “natural” log structure coming from the set  $S$ .



# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:



# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:

$$\mathcal{M}(V) = \{f \in \mathcal{O}_X(V) : f|_{V \setminus D} \in \mathcal{O}_X(V \setminus D)^*\}$$



# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:

$$\mathcal{M}(V) = \{f \in \mathcal{O}_X(V) : f|_{V \setminus D} \in \mathcal{O}_X(V \setminus D)^*\}$$

- $\mathcal{M}$  has the structure of sheaf of **monoids** on  $X$  (sum of invertible functions needs not to be invertible);





# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:

$$\mathcal{M}(V) = \{f \in \mathcal{O}_X(V) : f|_{V \setminus D} \in \mathcal{O}_X(V \setminus D)^*\}$$

- $\mathcal{M}$  has the structure of sheaf of **monoids** on  $X$  (sum of invertible functions needs not to be invertible);
- There exists a well-defined map  $\alpha : \mathcal{M} \rightarrow \mathcal{O}_X$  which is the identity when restricted to  $\mathcal{O}_X^*$ ;



# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:

$$\mathcal{M}(V) = \{f \in \mathcal{O}_X(V) : f|_{V \setminus D} \in \mathcal{O}_X(V \setminus D)^*\}$$

- $\mathcal{M}$  has the structure of sheaf of **monoids** on  $X$  (sum of invertible functions needs not to be invertible);
- There exists a well-defined map  $\alpha : \mathcal{M} \rightarrow \mathcal{O}_X$  which is the identity when restricted to  $\mathcal{O}_X^*$ ;
- Derivatives of sections of  $\mathcal{M}$  generates  $H^0(X, \Omega_X^1(D))$  where  $\Omega_X^1(D)$  is the sheaf of differential forms with logarithmic poles along  $D$  (name *logarithmic geometry*).



# Fundamental example

$X$  a smooth variety,  $D$  a normal crossing divisor on  $X$ . Consider the following sheaf:

$$\mathcal{M}(V) = \{f \in \mathcal{O}_X(V) : f|_{V \setminus D} \in \mathcal{O}_X(V \setminus D)^*\}$$

- $\mathcal{M}$  has the structure of sheaf of **monoids** on  $X$  (sum of invertible functions needs not to be invertible);
- There exists a well-defined map  $\alpha : \mathcal{M} \rightarrow \mathcal{O}_X$  which is the identity when restricted to  $\mathcal{O}_X^*$ ;
- Derivatives of sections of  $\mathcal{M}$  generates  $H^0(X, \Omega_X^1(D))$  where  $\Omega_X^1(D)$  is the sheaf of differential forms with logarithmic poles along  $D$  (name *logarithmic geometry*).
- The normal crossing condition can be rephrased in local terms if one considers the étale topology instead of the Zariski topology.



# Logarithmic Scheme

## Definition

A *logarithmic scheme* is a couple  $(X, \mathcal{M})$  where  $X$  is a scheme and  $\mathcal{M}$  is a sheaf of monoid on the étale site of  $X$ , called a **log structure** together with a morphisms of sheaves of monoids  $\alpha : \mathcal{M} \rightarrow \mathcal{O}_X$  such that  $\alpha^{-1}\mathcal{O}_X^* \rightarrow \mathcal{O}_X^*$  is an isomorphism.

A *morphism of log schemes*  $(X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$  is a couple  $(f, f^b)$  where  $f : \underline{X} \rightarrow \underline{Y}$  is a morphisms of schemes and  $f^b : f^*\mathcal{M}_Y \rightarrow \mathcal{M}_X$  is a morphism of log structures on  $X$ .



# Logarithmic Scheme

## Definition

A *logarithmic scheme* is a couple  $(X, \mathcal{M})$  where  $X$  is a scheme and  $\mathcal{M}$  is a sheaf of monoid on the étale site of  $X$ , called a **log structure** together with a morphisms of sheaves of monoids  $\alpha : \mathcal{M} \rightarrow \mathcal{O}_X$  such that  $\alpha^{-1}\mathcal{O}_X^* \rightarrow \mathcal{O}_X^*$  is an isomorphism.

A *morphism of log schemes*  $(X, \mathcal{M}_X) \rightarrow (Y, \mathcal{M}_Y)$  is a couple  $(f, f^b)$  where  $f : \underline{X} \rightarrow \underline{Y}$  is a morphisms of schemes and  $f^b : f^*\mathcal{M}_Y \rightarrow \mathcal{M}_X$  is a morphism of log structures on  $X$ .

Here  $f^*\mathcal{M}_Y$  is the logarithmic structure associated to the map  $f^{-1}(\mathcal{M}_Y) \rightarrow f^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$ , called the inverse image.



# Log curves are prestable

- Logarithmic geometry appears naturally when considering stable curves.



# Log curves are prestable

- Logarithmic geometry appears naturally when considering stable curves.
- Every log curve is naturally a pointed nodal curve.



# Log curves are prestable

- Logarithmic geometry appears naturally when considering stable curves.
- Every log curve is naturally a pointed nodal curve.
- Every pointed and at most nodal curve carries a “canonical” log structure





# Back to $\mathbb{P}^2, D$

Goal: extend CZ result for  $D$  a conic and two lines to log-stable curves.



# Back to $\mathbb{P}^2, D$

Goal: extend CZ result for  $D$  a conic and two lines to log-stable curves.

First we need to address the problem of curves  $\mathcal{C}$  with more than one irreducible components. How to properly extend  $S = \mathcal{C} \cap D$ ?



# Back to $\mathbb{P}^2$ , $D$

Goal: extend CZ result for  $D$  a conic and two lines to log-stable curves.

First we need to address the problem of curves  $\mathcal{C}$  with more than one irreducible components. How to properly extend  $S = \mathcal{C} \cap D$ ?

We need to take into account the nodes, i.e.  $S$  will be the set of distinguished points (marked points and nodes).



# Back to $\mathbb{P}^2$ , $D$


Goal: extend CZ result for  $D$  a conic and two lines to log-stable curves.


First we need to address the problem of curves  $\mathcal{C}$  with more than one irreducible components. How to properly extend  $S = \mathcal{C} \cap D$ ?


We need to take into account the nodes, i.e.  $S$  will be the set of distinguished points (marked points and nodes).


Moreover we would like to have a well defined notion of  $S$  in **each** irreducible component. Therefore nodes should count as two points (this can be made precise using log charts).













# log stability

Consider  $\varphi : \tilde{\mathcal{C}} \rightarrow \mathbb{P}^2$ : log-structures are given respectively by  $S$  and  $D$ . Log stability in this case is equivalent to usual stability plus the following conditions:



# log stability

Consider  $\varphi : \tilde{\mathcal{C}} \rightarrow \mathbb{P}^2$ : log-structures are given respectively by  $S$  and  $D$ . Log stability in this case is equivalent to usual stability plus the following conditions:

- 1 For every irreducible component  $\mathcal{C}_i$  of  $\tilde{\mathcal{C}}$  such that  $\mathcal{C}_i$  maps to a degree one irreducible component of  $D$ ,  $S_{\mathcal{C}_i}$  contains at least **three** points.



# log stability

Consider  $\varphi : \tilde{\mathcal{C}} \rightarrow \mathbb{P}^2$ : log-structures are given respectively by  $S$  and  $D$ . Log stability in this case is equivalent to usual stability plus the following conditions:

- 1 For every irreducible component  $\mathcal{C}_i$  of  $\tilde{\mathcal{C}}$  such that  $\mathcal{C}_i$  maps to a degree one irreducible component of  $D$ ,  $S_{\mathcal{C}_i}$  contains at least **three** points.
- 2 For every irreducible component  $\mathcal{C}_j$  of  $\mathcal{C}$  such that  $\mathcal{C}_j$  maps to the degree two irreducible component of  $D$ ,  $S_{\mathcal{C}_j}$  contains at least **four** points.



# Extension to log stable maps

With previous definition the following extension of CZ holds

## Proposition

Given  $\tilde{C}, S, D$  as above, let  $\varphi : \tilde{C} \rightarrow \mathbb{P}^2$  be a non-constant *stable* log-morphism such that  $\varphi^{-1}(D) \subset S$ . Then the degree of the image  $\varphi(\tilde{C})$  verifies:

$$\deg(\varphi(\tilde{C})) \leq A \cdot \max\{1, \chi(\tilde{C} \setminus S)\}$$



# Extension to log stable maps

With previous definition the following extension of CZ holds

## Proposition

Given  $\tilde{C}, S, D$  as above, let  $\varphi : \tilde{C} \rightarrow \mathbb{P}^2$  be a non-constant *stable* log-morphism such that  $\varphi^{-1}(D) \subset S$ . Then the degree of the image  $\varphi(\tilde{C})$  verifies:

$$\deg(\varphi(\tilde{C})) \leq A \cdot \max\{1, \chi(\tilde{C} \setminus S)\}$$

Here  $A$  is the same constant appearing in CZ Theorem.



# Moduli interpretation

Pivotal observation: previous Proposition can be rephrased in the following way:



# Moduli interpretation

Pivotal observation: previous Proposition can be rephrased in the following way:

Given a log-stable curve  $\mathcal{C}$  as before, the degree of the image of every log-stable map to  $(\mathbb{P}^2, \mathcal{M}_D)$  corresponds to the degree of the Chow class of the image curve.



# Moduli interpretation

Pivotal observation: previous Proposition can be rephrased in the following way:

Given a log-stable curve  $\mathcal{C}$  as before, the degree of the image of every log-stable map to  $(\mathbb{P}^2, \mathcal{M}_D)$  corresponds to the degree of the Chow class of the image curve.

In particular for a map  $f : (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow (\mathbb{P}^2, \mathcal{M}_D)$  if  $f_*([\mathcal{C}]) = \beta$  for  $\beta \in A^1(\mathbb{P}^2)$  then  $\deg f(\mathcal{C}) = \deg \beta$ .





# Moduli interpretation

Pivotal observation: previous Proposition can be rephrased in the following way:

Given a log-stable curve  $\mathcal{C}$  as before, the degree of the image of every log-stable map to  $(\mathbb{P}^2, \mathcal{M}_D)$  corresponds to the degree of the Chow class of the image curve.

In particular for a map  $f : (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow (\mathbb{P}^2, \mathcal{M}_D)$  if  $f_*([\mathcal{C}]) = \beta$  for  $\beta \in A^1(\mathbb{P}^2)$  then  $\deg f(\mathcal{C}) = \deg \beta$ .

Hence once  $\beta \in A^1(\mathbb{P}^2)$   $g = g(\beta)$  and  $n = \#S$  are fixed a log-stable maps  $f : (\mathcal{C}, \mathcal{M}_{\mathcal{C}}) \rightarrow (\mathbb{P}^2, \mathcal{M}_D)$  from a genus  $g$ ,  $n$ -marked curve exists only if  $\deg \beta$  verifies the inequality of Proposition.



# The moduli space of log stable maps

Previous remark can be made precise using the moduli space  $\mathcal{K}_\Gamma(\mathbb{P}^2, \mathcal{M}_D)$  for a discrete data  $\Gamma$ .



# The moduli space of log stable maps

Previous remark can be made precise using the moduli space  $\mathcal{K}_\Gamma(\mathbb{P}^2, \mathcal{M}_D)$  for a discrete data  $\Gamma$ .

## Definition (Discrete data)

Let  $\Gamma = (\beta, g, n, \vec{c})$  be a fourple consisting of the following data:

- $\beta \in H^2(X, \mathbb{Z})$  is a curve class;
- $n, g$  are two non-negative integers (*marked points* and *genus*);
- $\vec{c}$  is a  $n$ -vector of non-negative integers (*multiplicities*) that verify:

$$\sum_{i=1}^n c_i = c_1(D) \cap \beta$$



# The moduli space of log stable maps

Previous remark can be made precise using the moduli space  $\mathcal{K}_\Gamma(\mathbb{P}^2, \mathcal{M}_D)$  for a discrete data  $\Gamma$ .

## Definition (Discrete data)

Let  $\Gamma = (\beta, g, n, \vec{c})$  be a fourple consisting of the following data:

- $\beta \in H^2(X, \mathbb{Z})$  is a curve class;
- $n, g$  are two non-negative integers (*marked points* and *genus*);
- $\vec{c}$  is a  $n$ -vector of non-negative integers (*multiplicities*) that verify:

$$\sum_{i=1}^n c_i = c_1(D) \cap \beta$$

We denote by  $\mathcal{K}_\Gamma(\mathbb{P}^2, \mathcal{M}_D)$  the moduli space of log-stable map to  $(\mathbb{P}^2, \mathcal{M}_D)$  of type  $\Gamma$ .



CZ equivalent to  $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta) = \emptyset$

### Proposition

*Given a curve  $\mathcal{B} \subset X$  such that  $S = \mathcal{B} \cap D$  and  $\deg(\mathcal{B}) > A\chi_s(\mathcal{B})$ , let  $\beta$  denote the corresponding element of  $A^1(\mathbb{P}^2)$ . Then the moduli space  $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$  is empty for  $g = g(\mathcal{B})$ ,  $n = \#S$  and every vector of multiplicities.*

### Proposition

*Suppose that for every plane curve  $\mathcal{B}$  with  $\deg \mathcal{B} > A\chi_{S_B} \mathcal{B}$ , where  $S_B = \mathcal{B} \cap D$ ,  $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$  are empty if  $g = g(\mathcal{B})$  and  $n = \#S_B$ . Then every log-stable map  $f$  from a genus  $g$  curve  $\mathcal{C}$  to  $X$  with log-structure  $f^*\mathcal{M}_D$  (and  $S = S_{\mathcal{C}} = f^{-1}(D)$ ) verifies*

$$\deg f(\mathcal{C}) \leq A\chi_S(\mathcal{C})$$



# Using properness of $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$

We have reduced the weak algebraic hyperbolicity problem to the emptiness of a stack. Now we use the following Theorem:



# Using properness of $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$

We have reduced the weak algebraic hyperbolicity problem to the emptiness of a stack. Now we use the following Theorem:

Theorem (Chen, Abramovich-Chen, Gross-Siebert)

$\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$  is a proper DM Stack.



## Using properness of $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$

We have reduced the weak algebraic hyperbolicity problem to the emptiness of a stack. Now we use the following Theorem:

Theorem (Chen, Abramovich-Chen, Gross-Siebert)

$\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$  is a proper DM Stack.

The Theorem implies that the condition  $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta) = \emptyset$  extends to a very generic  $D$  of degree 4 over a flat family which is logarithmically smooth.





# Using properness of $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$

We have reduced the weak algebraic hyperbolicity problem to the emptiness of a stack. Now we use the following Theorem:

Theorem (Chen, Abramovich-Chen, Gross-Siebert)

$\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta)$  is a proper DM Stack.

The Theorem implies that the condition  $\mathcal{K}_\Gamma((\mathbb{P}^2, \mathcal{M}_D), \beta) = \emptyset$  extends to a very generic  $D$  of degree 4 over a flat family which is logarithmically smooth.

With previous propositions this gives the conclusion.



# Final comments

- Theorem uses in an essential way that  $\mathcal{M}_D$  comes for a **simple** normal crossing divisor on  $D$  (i.e.  $\mathbb{P}^2, \mathcal{M}_D$  is a Deligne-Faltings pair).



# Final comments

- Theorem uses in an essential way that  $\mathcal{M}_D$  comes for a **simple** normal crossing divisor on  $D$  (i.e.  $\mathbb{P}^2, \mathcal{M}_D$  is a Deligne-Faltings pair).
- Bound obtained works only for minimal **stable** maps.



# Final comments

- Theorem uses in an essential way that  $\mathcal{M}_D$  comes for a **simple** normal crossing divisor on  $D$  (i.e.  $\mathbb{P}^2, \mathcal{M}_D$  is a Deligne-Faltings pair).
- Bound obtained works only for minimal **stable** maps.
- Log smoothness hypothesis can possibly be removed (work in progress).



# Final comments

- Theorem uses in an essential way that  $\mathcal{M}_D$  comes for a **simple** normal crossing divisor on  $D$  (i.e.  $\mathbb{P}^2, \mathcal{M}_D$  is a Deligne-Faltings pair).
- Bound obtained works only for minimal **stable** maps.
- Log smoothness hypothesis can possibly be removed (work in progress).
- Possibilities to extend to varieties other than  $\mathbb{P}^2$ .



Thank you for your attention

