

Promise Constraints and Cheese

Antoine Mottet
TU Hamburg

Dagstuhl 20.05.2022

$A, B \dots$ relational structures with $A \rightarrow B$

$\text{PCSP}(A, B)$: decide if $\underbrace{X \rightarrow A}_{\text{YES}}$ or $\underbrace{X \not\rightarrow B}_{\text{NO}}$

$A, B \dots$ relational structures with $A \rightarrow B$

$\text{PCSP}(A, B)$: decide if $\underbrace{X \rightarrow A}_{\text{YES}}$ or $\underbrace{X \not\rightarrow B}_{\text{NO}}$

(A, B) sandwiches C if $A \rightarrow C \rightarrow B$ ($\Rightarrow \text{PCSP}(A, B) \leq \text{CSP}(C)$,
search version ...)

$A, B \dots$ relational structures with $A \rightarrow B$

$\text{PCSP}(A, B)$: decide if $\underbrace{X \rightarrow A}_{\text{YES}}$ or $\underbrace{X \not\rightarrow B}_{\text{NO}}$

(A, B) sandwiches C if $A \rightarrow C \rightarrow B$ ($\Rightarrow \text{PCSP}(A, B) \leq \text{CSP}(C)$,
search version ...)

Observation:

IF $\text{PCSP}(A, B)$ is in P, there is C s.t.

$\text{CSP}(C)$ is in P and $\text{PCSP}(A, B) \leq \text{CSP}(C)$.

Proof: M solving $\text{PCSP}(A, B)$ $L(M) = \text{CSP}(C)$ (Bodirsky-Grohe)
 \square

$A, B \dots$ relational structures with $A \rightarrow B$

$\text{PCSP}(A, B)$: decide if $\underbrace{X \rightarrow A}_{\text{YES}}$ or $\underbrace{X \not\rightarrow B}_{\text{NO}}$

(A, B) sandwiches C if $A \rightarrow C \rightarrow B$ ($\Rightarrow \text{PCSP}(A, B) \leq \text{CSP}(C)$,
search version ...)

Observation:

IF $\text{PCSP}(A, B)$ is in P, there is C s.t.

$\text{CSP}(C)$ is in P and $\text{PCSP}(A, B) \leq \text{CSP}(C)$.

Proof: M solving $\text{PCSP}(A, B)$ $L(M) = \text{CSP}(C)$ (Bodirsky-Grohe)
 \square

Not quite a sandwich...

Algorithm M is "monotone" if

$(M \text{ accepts } Y \text{ and } X \rightarrow Y) \Rightarrow M \text{ accepts } X$

Algorithm M is "monotone" if

$(M \text{ accepts } Y \text{ and } X \rightarrow Y) \Rightarrow M \text{ accepts } X$

Observation 2:

(A, B) sandwiches a tractable C iff $\text{PCSP}(A, B)$ solvable by
a polynomial-time "monotone" algorithm.

Algorithm M is "monotone" if

$$(M \text{ accepts } Y \text{ and } X \rightarrow Y) \Rightarrow M \text{ accepts } X$$

Observation 2:

(A, B) sandwiches a tractable C iff $\text{PCSP}(A, B)$ solvable by a polynomial-time "monotone" algorithm.

Proof: $C = \text{disjoint union of structures accepted by algorithm}$.

Corollary: \emptyset

Algorithm M is "monotone" if

$$(M \text{ accepts } Y \text{ and } X \rightarrow Y) \Rightarrow M \text{ accepts } X$$

Observation 2:

(A, B) sandwiches a tractable C iff $\text{PCSP}(A, B)$ solvable by a polynomial-time "monotone" algorithm.

Proof: $C = \text{disjoint union of structures accepted by algorithm}$.

Corollary: \emptyset

Theme: Find nicer cheese under assumption that $\text{PCSP}(A, B)$ is solved by specific algorithm.

- search $\text{PCSP}(A, B)$
- $\text{Pol}(C) \rightarrow \text{BI}(A, B)$


① AC⁰ / FIRST ORDER LOGIC AND TIME TRAVEL

PCSP(A, B) is solved by Φ if:

- $X \rightarrow A \Rightarrow X + \overline{\Phi}$
- $X \not\rightarrow B \Rightarrow X + \overline{\Phi} \quad / \quad X + \overline{\Phi} \Rightarrow X \rightarrow B$.

Question: (Oprsal '20)

Can one characterize (A, B) s.t.
 $\text{PCSP}(A, B)$ is in FO?




① AC⁰ / FIRST ORDER LOGIC AND TIME TRAVEL

$\text{PCSP}(A, B)$ is solved by Φ if:

- $X \rightarrow A \Rightarrow X + \overline{\Phi}$
- $X \not\rightarrow B \Rightarrow X + \overline{\Phi} \quad / \quad X + \overline{\Phi} \Rightarrow X \rightarrow B$

Question: (Oprsal '20)

Can one characterize (A, B) s.t.
 $\text{PCSP}(A, B)$ is in FO?



Theorem: (Atserias)

A finite. $\text{CSP}(A)$ is in FO iff $\exists \mathcal{F}$ finite
 s.t. $X \rightarrow A \Leftrightarrow \nexists F \in \mathcal{F}: F \rightarrow X$

Finite duality

Question: (Oprsal '21)

$\text{PCSP}(A, B)$ in FO iff $\exists \mathcal{F}$ finite s.t.
 $X \rightarrow A \Rightarrow (\exists F \in \mathcal{F}: F \rightarrow X) \quad \text{and} \quad (\nexists F \in \mathcal{F}: F \rightarrow X) \Rightarrow X \rightarrow B$?


① AC⁰ / FIRST ORDER LOGIC AND TIME TRAVEL

PCSP(A, B) is solved by Φ if:

- $X \rightarrow A \Rightarrow X + \overline{\Phi}$
- $X \not\rightarrow B \Rightarrow X + \overline{\Phi} / X + \overline{\Phi} \Rightarrow X \rightarrow B$

Theorem: (Atserias)

A finite. $CSP(A)$ is in FO iff $\exists \mathcal{F}$ finite
s.t. $X \rightarrow A \Leftrightarrow \nexists F \in \mathcal{F}: F \rightarrow X$



Question: Opršal '22

CASSETTE SOME ALGORITHMS.

- local consistency
- Gelench - Adams
- CLP {Ciardo, Zivny}
- finite domain
- definability in various logics.

Theorem: (Rossman '08)

Yes.

Theorem: (Rossman '08)

Yes.

For finite CSPs:

- finite duality \Rightarrow tree duality
- characterization by "1-tolerant" polymorphisms
- $\text{CSP}(A) \in \text{FO?}$ in NP

Larose, Loten, Tardif

Theorem: (Rossman '08)

Yes.

For finite CSPs:

- finite duality \Rightarrow tree duality
- characterization by "1-tolerant" polymorphisms
- $\text{CSP}(A) \in \text{FO} ?$ in NP

Larose, Loten, Tardif

Theorem: (M.)

For finite A, B , TFAE:

- $\text{PCSP}(A, B) \in \text{FO}$
- (A, B) has finite duality $\xrightarrow{5} \text{Rossman}$
- (A, B) has finite tree duality
- $\exists C$ finite s.t. $A \rightarrow C \rightarrow B$ and $\text{CSP}(C) \in \text{FO}$.

Theorem: (Rossman '08)

Yes.

For finite CSPs:

- finite duality \Rightarrow tree duality
- characterization by "1-tolerant" polymorphisms
- $\text{CSP}(A) \in \text{FO}?$ in NP

Larose, Loten, Tardif

Theorem: (M.)

For finite A, B , TFAE:

- $\text{PCSP}(A, B) \in \text{FO}$
- (A, B) has finite duality $\xrightarrow{5} \text{Rossman}$
- (A, B) has finite tree duality
- $\exists C$ finite s.t. $A \rightarrow C \rightarrow B$ and $\text{CSP}(C) \in \text{FO}$.

But: • no characterization by 1-tolerant polymorphisms (necessary not sufficient)
• $\text{PCSP}(A, B) \in \text{FO}?$ decidable?

Proof: \mathcal{F} finite duality for (A, B) : $X \rightarrow A \Rightarrow \nexists F \in \mathcal{F} : F \rightarrow X$
 $X \not\rightarrow B \Rightarrow \exists F \in \mathcal{F} : F \rightarrow X$

Wlog every $F \in \mathcal{F}$ is connected.


$\exists U$ ω -categorical s.t. $X \rightarrow U$ iff $\nexists F \in \mathcal{F} : F \rightarrow X$.
 (Cherlin, Shelah, Shi)

$$A \rightarrow A \Rightarrow \nexists F \in \mathcal{F} : F \rightarrow A \Rightarrow A \rightarrow U$$

$$U \rightarrow U \Rightarrow \nexists F \in \mathcal{F} : F \rightarrow U \Rightarrow U \rightarrow B.$$

$\exists U^+$ ω -categorical Ramsey expansion of U
 (Hubička, Nešetřil)

$$A \rightarrow U \xrightarrow{\pi} U /_{\text{Aut}(U^+)} \xrightarrow{h} B$$



Proof (cont'd):

$$A \rightarrow U \xrightarrow{\pi} U /_{\text{Aut}(U^+)} \xrightarrow{h} B$$

and U has finite duality.

U has 1-tolerant polymorphism: $f: U^k \rightarrow U$ s.t. $f(R^0, \dots, R^e, U^n, R^0, \dots, R^e) \subseteq R^0$
 ✓ relation R^0 of U

- [BPT] $\Rightarrow U$ has 1-tolerant polymorphism that is canonical wrt U^+
- $\Rightarrow U /_{\text{Aut}(U^+)} \text{ has 1-tolerant polymorphism}$
- $\Rightarrow \underline{U /_{\text{Aut}(U^+)}}$ has finite duality. □

Proof (cont'd):

$$A \xrightarrow{\pi} U \xrightarrow{\text{Aut}(U^+)} B$$

and U has finite duality.

U has 1-tolerant polymorphism: $f: U^k \rightarrow U$ s.t. $f(R^0, \dots, R^e, U^n, R^0, \dots, R^u) \subseteq R^U$
 ✓ relation R^U of U

- [BPT] $\Rightarrow U$ has 1-tolerant polymorphism that is canonical wrt U^+
- $\Rightarrow U/\text{Aut}(U^+)$ has 1-tolerant polymorphism
- $\Rightarrow \underline{U/\text{Aut}(U^+)}$ has finite duality. □

(IK_3, IK_g) has binary 1-tolerant polymorphism but $\text{PCSP}(IK_3, IK_g)$ not in FD.
 Open: bound on $|C|$?


② PCSPs WITH BOUNDED WIDTH

(k, l) -consistency solves $\text{PCSP}(A, B)$ if

X (k, l) -consistent wrt $A \Rightarrow X \rightarrow B$.

}

\exists family of sets $P_K \subseteq A^K$ ($K \subseteq X$, $|K| \leq k$)
 s.t. $P_K \subseteq$ partial homomorphisms $X[K] \rightarrow A$
 and $\forall K \subseteq L \rightarrow P_K \subseteq P_L$




CLASSIFY SOME ALGORITHMS.

- local consistency ?
- Gelenbe - Adams
- CLAP [Ciardo, Zivny]
- finite duality ✓
- definability in various logics.

Fix A

X (k, l) -consistent




0
1
A

Fix A

X (k, l) -consistent

(X, P) decoration by a (k, l) -system




0
1
A

$\mathcal{K}^+ := \left\{ X + \begin{matrix} \text{non-trivial} \\ (k, l) \text{-system} \end{matrix} \mid X (k, l) \text{-consistent wrt } A \right\}$

Fix A

X (k, l) -consistent
 (X, P) decoration by a (k, l) -system




0
 1
 A

$\mathcal{K}^+ := \left\{ X + \begin{array}{c} \text{non-trivial} \\ (k, l)-\text{system} \end{array} \mid X \text{ } (k, l)\text{-consistent wrt. } A \right\}$

Lemma:

\mathcal{K}^+ has the (strong) amalgamation property:

$(Y, P), (Z, Q)$ (k, l) -consistent and
 $P|_{Y \cap Z} = Q|_{Y \cap Z}$
 $\Rightarrow Y \cup Z$ (k, l) -consistent.



$\rightsquigarrow \exists (U, P)$ universal limit object, ω -categorical (and more)

$$\mathcal{K}^+ := \left\{ X \text{ non-trivial } (k,l)\text{-system} \quad \middle| \quad X \text{ (k,l)-consistent w.r.t. } A \right\}$$

$$X \rightarrow U \Leftrightarrow X \text{ (k,l)-consistent w.r.t. } A$$

- \$A\$ \$(k,l)\$-consistent w.r.t. \$A \Rightarrow A \rightarrow U\$

$$\mathcal{K}^+ := \left\{ X \text{ non-trivial } (k,l)\text{-system} \quad \middle| \quad X \text{ (k,l)-consistent w.r.t. } A \right\}$$

$$X \rightarrow U \Leftrightarrow X \text{ (k,l)-consistent w.r.t. } A$$

- \$A(k,l)\$-consistent w.r.t. \$A \Rightarrow A \rightarrow U\$
- \$U \rightarrow U \Rightarrow U \text{ \$(k,l)\$-consistent w.r.t. } A\$

$$\mathcal{K}^+ := \left\{ X \text{ non-trivial } (k,l)\text{-system} \quad \middle| \quad X(h,l) \text{-consistent wrt. } A \right\}$$

$$X \rightarrow U \Leftrightarrow X(h,l) \text{-consistent wrt. } A$$

- $A(h,l)$ -consistent wrt. $A \Rightarrow A \rightarrow U$
- $U \rightarrow U \Rightarrow U(h,l)$ -consistent wrt. A
- U has width (k,l) :

$$\begin{aligned} & X(h,l) \text{-consistent wrt. } U \\ \Rightarrow & X(h,l) \text{-consistent wrt. } A \\ \Rightarrow & X \rightarrow U. \end{aligned}$$

Theorem: (M.)

$A, 1 \leq k \leq l$. There exists \mathbb{U} ω -categorical^{††} s.t.
 $A \rightarrow \mathbb{U}$, \mathbb{U} has width (k, l) , and for all B , TFAE:

- (A, B) has width (k, l)
- $\mathbb{U} \rightarrow B$.

\mathbb{U} is universal with these properties.

Polymorphisms?

Theorem: (M.)

$A, 1 \leq k \leq l$. There exists \mathbb{U} ω -categorical^{††} s.t.
 $A \rightarrow \mathbb{U}$, \mathbb{U} has width (k, l) , and for all B , TFAE:

- (A, B) has width (k, l)
- $\mathbb{U} \rightarrow B$.

\mathbb{U} is universal with these properties.

Polymorphisms?

Theorem) (Atserias, Dalmau)

If (A, B) has bounded width then (A, B) has WNU(n) $\forall n \geq 3$.

Indicator structure:

$$\Sigma = w(yxx) \approx w(xyx) \approx w(xxy)$$

$$A = \{0, 1\}$$

$$\Sigma(A) = \frac{A^3}{\sim_{\Sigma}}$$

$$001 \sim_{\Sigma} 010 \sim_{\Sigma} 100$$

Observation: $\Sigma(A) \rightarrow A$ iff $\text{Pol}(A)$ satisfies Σ

$X \rightarrow U \Leftrightarrow X \text{ } (h, l)\text{-consistent wrt. } A$

Proposition:

For every set Σ of height 1 identities, TFAE:

- $\text{Pol}(U)$ satisfies Σ
- $\text{Pol}(A, U)$ satisfies Σ
- $\Sigma(A)$ (h, l) -consistent wrt. A

$\forall n \text{ WNU}(n) \times$ (not even $\forall n$ large)

$\forall n \text{ WNU}(n) \times$ (not even $\forall n$ large)

$\forall n \text{ dWNU}(n)$ ✓
(Barto terms)

$\text{WNU}(n) \Rightarrow \text{dWNU}(n)$

$n=3:$

$$f_1(yxx) \approx f_3(xyx) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_3(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$

$\forall n$ large phNU(n) ✓

(not height 1)

$$\alpha_1 f(yx\cdots x) \approx \alpha_2 f(xy\cdots x) \approx \cdots$$

$\forall n \text{ WNU}(n) \times$ (not even $\forall n$ large)

$\forall n \text{ dWNU}(n)$ ✓
(Barto terms)

$\text{WNU}(n) \Rightarrow \text{dWNU}(n)$

$n=3:$

$$f_1(yxx) \approx f_3(xyx) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_3(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$

...

$\forall n$ large $\text{pWNU}(n)$ ✓
(not height 1)

$$\alpha_1 f(yx\cdots x) \approx \alpha_2 f(xy\cdots x) \approx \cdots$$

Corollary:

A ω -categorical with bounded width,
 $\exists \mathbb{U}$ ω -categorical⁺⁺ s.t. $A \supseteq \mathbb{U}$ and \mathbb{U} has

dWNU polymorphisms of all arities.

Existence of dWNU(3) :

$$f_1(yxx) \approx f_3(yxy) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_4(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$


:

$$A = (\{0,1\}, \underbrace{x+y+z=0}_{x+y+z=1})$$

$\Sigma(A)$ consistent?

$$\text{Domain} = [f_1, 100], [f_1, 010], \dots \\ [f_3, "], [f_3, 010]$$

Let Π be a "tree" with $\Pi \rightarrow \Sigma(A)$.



Existence of dWNU(3) :


$$f_1(yxx) \approx f_3(yxy) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_4(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$

:




$$A = (\{0,1\}, \underbrace{x+y+z=0}_{x+y+z=1})$$

$\Sigma(A)$ consistent?

$$\text{Domain} = [f_1, 100], [f_1, 010], \dots \\ [f_3, 010]$$

Let Π be a "tree" with $\Pi \rightarrow \Sigma(A)$.



Existence of dWNU(3) :

$$f_1(yxx) \approx f_3(yxy) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_4(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$



:

$$A = (\{0,1\}, \underbrace{x+y+z=0}_{}, \underbrace{x+y+z=1}_{})$$

$\Sigma(A)$ consistent?

$$\text{Domain} = [f_1, 100], [f_1, 010], \dots \\ [f_3, "], [f_3, 010]$$

Let Π be a "tree" with $\Pi \rightarrow \Sigma(A)$.



Existence of dWNU(3) :


$$f_1(yxx) \approx f_3(yyx) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_4(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$

:



$$A = \left(\{0,1\}, \underbrace{x+y+z=0}_{x+y+z=1} \right)$$


$\Sigma(A)$ consistent?

$$\text{Domain} = [f_1, 100], [f_1, 010], \dots$$

\vdots

$[f_3, 010]$

Let Π be a "tree" with $\Pi \rightarrow \Sigma(A)$.



Existence of dWNU(3) :

$$f_1(yxx) \approx f_3(yxy) \approx f_4(xyx)$$

$$f_1(xyx) \approx f_2(xyx) \approx f_4(xxy)$$

$$f_1(xxy) \approx f_2(xxy) \approx f_3(xxy)$$

$$f_3(yxx) \approx f_2(yxx) \approx f_4(yxx)$$



:

$$A = (\{0,1\}, \underbrace{x+y+z=0}_{x+y+z=1})$$

$\Sigma(A)$ consistent?

$$\text{Domain} = [f_1, 100], [f_1, 010], \dots \\ [f_3, 010]$$

Let Π be a "tree" with $\Pi \rightarrow \Sigma(A)$.



So $\Pi \rightarrow A$

Open:

can \mathbb{U} be taken finite? decidable property?

Proof strategy:

① construct Ramsey expansion (exists for width 1)

② smooth approximations to make pWNUs (1-)canonical

Open: can \mathbb{U} be taken finite? decidable property?

Proof strategy:

① construct Ramsey expansion (exists for width 1)

② smooth approximations to make pWNUs (1-)canonical

If ① fails then counterexample to conjecture in Ramsey theory (Bodirsky - Pinsker - Trankov)

If ① holds then search PCSP(A, B) tractable if bounded width.
+ probable collapse of bounded width hierarchy.

Open: can \mathbb{U} be taken finite? decidable property?

Proof strategy:

① construct Ramsey expansion

② smooth approximations to make pWNUs (1-)canonical

If ① fails then counterexample to conjecture in Ramsey theory (Bodirsky - Pinsker - Trankov)

If ① holds then search PCSP(A, B) tractable if bounded width.
+ probable collapse of bounded width hierarchy.

CLASSIFY SOME ALGORITHMS.

- local consistency ?
- Glenshi - Adams
- CLAT [Ciardo, Zivny]
- finite duality ✓
- definability in various logics.

General argument:

Suppose one has a pair \mathcal{A} ... algorithm s.t.
 M ... minion

\mathcal{A} accepts X iff $X \rightarrow F_M(\mathcal{A})$

General argument:

Suppose one has a pair \mathcal{A} ... algorithm s.t.
 \mathcal{M} ... minion

\mathcal{A} accepts X iff $X \rightarrow F_{\mathcal{M}}(A)$

$K = \{(X, h) \mid h: X \rightarrow F_{\mathcal{M}}(A)\}$ $U = K$ -universal object
(Fraïssé-limit)

General argument:

Suppose one has a pair \mathcal{A} ... algorithm s.t.
 \mathcal{M} ... minion

\mathcal{A} accepts X iff $X \rightarrow F_{\mathcal{M}}(A)$

$K = \{(X, h) \mid h: X \rightarrow F_{\mathcal{M}}(A)\}$ $U = K$ -universal object
(Fraïssé-limit)

Then:

- \mathcal{A} accepts X iff $X \rightarrow U$
- \mathcal{A} solves PCSP(A, B) iff $U \rightarrow B$
- $U \rightarrow F_{\mathcal{M}}(A)$ if \mathcal{M} locally finite.
- $F_{\mathcal{M}}(A) \rightarrow U$

General argument:

Suppose one has a pair \mathcal{A} ... algorithm s.t.
 \mathcal{M} ... minion

\mathcal{A} accepts X iff $X \xrightarrow{\text{ok}} F_{\mathcal{M}}(A^{\otimes k})$

$K = \{(X, h) \mid h: X \xrightarrow{\text{ok}} F_{\mathcal{M}}(A^{\otimes k})\}$ $U = K$ -universal object
(Fraïssé-limit)

Then:

- \mathcal{A} accepts X iff $X \rightarrow U$
- \mathcal{A} solves PCSP(A, B) iff $U \rightarrow B$
- $U \rightarrow F_{\mathcal{M}}(A)$ if \mathcal{M} locally finite. \times
- $F_{\mathcal{M}}(A) \rightarrow U$ \times