

MMSNP: Proof of the Universal-Algebraic Dichotomy Conjecture

Manuel Bodirsky, Florent Madelaine, **Antoine Mottet**

September 25, 2018

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity:

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

-
- ▶ Input: undirected graph G
 - ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: in P.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg($$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x})) \quad)$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?
(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

MMSNP and FPP are computationally equivalent.

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,
- ▶ Solve question posed by Lutz and Wolter (ICDT'15),

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,
- ▶ Solve question posed by Lutz and Wolter (ICDT'15),
- ▶ Confirmation of conjecture by Bodirsky-M (LICS'16).

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

Definition

$\mathfrak{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathfrak{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathfrak{A} ,
- ▶ **Question:** is there a homomorphism $\mathfrak{A} \rightarrow \mathfrak{B}$?

Definition

$\mathfrak{B} = (B; E)$ a **graph**.


$\text{CSP}(\mathfrak{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathfrak{A} ,
- ▶ **Question:** is there a homomorphism $\mathfrak{A} \rightarrow \mathfrak{B}$?


In general, the forbidden patterns problem (FPP) for \mathcal{F} is **not a CSP**, but a **finite union** of CSPs.


Proposition


Every FPP reduces in polynomial-time to a finite number of FPP of connected structures.











Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F}).$$

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F}).$$

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

FPP of finite sets of finite **connected** structures

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$.

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

FPP of finite sets of finite **connected** structures



CSPs in the BP class


Obstructions \mathcal{F} :



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :



Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

Conjecture (Bodirsky-Pinsker, 2011)

*Let \mathfrak{B} be a reduct of a finitely bounded homogeneous structure.
If there is no uniformly continuous $h1$ homomorphism
 $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$, then $\text{CSP}(\mathfrak{B})$ is in P .*

Conjecture (Bodirsky-Pinsker, 2011)

*Let \mathfrak{B} be a reduct of a finitely bounded homogeneous structure.
If there is no uniformly continuous $h1$ homomorphism
 $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$, then $\text{CSP}(\mathfrak{B})$ is in P .*

Theorem (Bodirsky-Madelaine-M, 2018)

*Let \mathfrak{B} be ω -categorical such that $\text{CSP}(\mathfrak{B})$ is in MMSNP.
If there is no uniformly continuous $h1$ homomorphism
 $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$, then $\text{CSP}(\mathfrak{B})$ is in P .*

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)


\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Rephrased: do $\text{CSP}(\mathfrak{B}, \bullet, \bullet)$ and $\text{CSP}(\mathfrak{B})$ have same complexity?



A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Proposition (Bodirsky, LMCS 2007)

*For ω -categorical model-complete cores, it is possible to add **constants** without changing the complexity.*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Proposition (Bodirsky, LMCS 2007)

*For ω -categorical model-complete cores, it is possible to add **constants** without changing the complexity.*

Good news: we can choose the MMSNP structure \mathfrak{B} so that (\mathfrak{B}, \neq) is an ω -categorical model-complete core.

Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :



Suppose the vertex x is precoloured in the input:



Proposition

The input precoloured graph is colourable iff the graph obtained by adding the gadgets is colourable.

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

Definition

$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

Definition

$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)

Definition


$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)




Every canonical function determines a **behaviour** on colours.



Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.
Only **finitely many** behaviours of a given arity.

Every canonical function determines a **behaviour** on colours.




We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

Theorem (Bodirsky-M, LICS'16)

Let \mathfrak{B} be in the BP class. If $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers operation modulo $\overline{\text{Aut}(\mathfrak{A})}$ that is canonical with respect to $\text{Aut}(\mathfrak{A})$, then $\text{CSP}(\mathfrak{B})$ is in P.

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

Theorem (Bodirsky-M, LICS'16)

Let \mathfrak{B} be in the BP class. If $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers operation modulo $\overline{\text{Aut}(\mathfrak{A})}$ that is canonical with respect to $\text{Aut}(\mathfrak{A})$, then $\text{CSP}(\mathfrak{B})$ is in P.

\Rightarrow let \mathcal{C} be the clone of canonical polymorphisms of the MMSNP structure \mathfrak{B} . If there is no h1 homomorphism $\mathcal{C} \rightarrow \mathcal{P}$, then $\text{CSP}(\mathfrak{B})$ is in P.

1. We suppose that there is a h1 homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{P}$.

1. We suppose that there is a h1 homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{P}$.
2. \mathcal{C} is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of \mathcal{C} that is isomorphic to \mathcal{P} .

1. We suppose that there is a h1 homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{P}$.
2. \mathcal{C} is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of \mathcal{C} that is isomorphic to \mathcal{P} .
3. Using this subfactor and Cthulhu partitions[†], we show that the polymorphisms of \mathfrak{B} **canonize in a unique way**,

(†): Coined collectively by the institute of algebra in Dresden.

1. We suppose that there is a h1 homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{P}$.
2. \mathcal{C} is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of \mathcal{C} that is isomorphic to \mathcal{P} .
3. Using this subfactor and Cthulhu partitions[†], we show that the polymorphisms of \mathfrak{B} **canonize in a unique way**[‡],

(†): Coined collectively by the institute of algebra in Dresden.

(‡): Loosely speaking.

1. We suppose that there is a h1 homomorphism $\xi: \mathcal{C} \rightarrow \mathcal{P}$.
2. \mathcal{C} is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of \mathcal{C} that is isomorphic to \mathcal{P} .
3. Using this subfactor and Cthulhu partitions[†], we show that the polymorphisms of \mathfrak{B} **canonize in a unique way**[‡],
4. Define an h1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ by canonizing and composing with ξ .

(†): Coined collectively by the institute of algebra in Dresden.

(‡): Loosely speaking.

σ : set of colour symbols. A **trivial subfactor** of \mathcal{C} is a partition $S \uplus T \subseteq \sigma$ such that the equivalence relation with blocks S and T is a congruence of \mathcal{C} with the property that the clone induced by \mathcal{C} on $\{S, T\}$ is isomorphic to \mathcal{P} .

Proposition

S, T trivial subfactor of \mathcal{C} . $\exists (B, E)$ undirected graph s.t.:

- ▶ (B, E) contains an edge from S to T but does not contain *pseudo-loops*;
- ▶ the connected components of N are included in S , included in T , or bipartite with the bipartition induced by S and T ;
- ▶ E is preserved by $\text{Pol}(\mathfrak{B})$.

We say that $\{S, T\}$ is a Cthulhu partition of (B, E) .

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathfrak{B}, <)$ is ω -categorical and Ramsey.

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathfrak{B}, <)$ is ω -categorical and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathfrak{B}, <)$ is ω -categorical and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Given $f \in \text{Pol}(\mathfrak{B})$, and $\xi: \mathcal{C} \rightarrow \mathcal{P}$ given by subfactor $\{S, T\}$, define $\phi(f) := \xi(g)$ where g is an arbitrary function in $\mathcal{C} \cap \overline{\text{Aut}(\mathfrak{B}, <)f\text{Aut}(\mathfrak{B}, <)}$.

Proposition

$\phi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ is a well-defined uniformly continuous height 1 homomorphism.

Summing up:

Theorem

Let \mathfrak{B} be a MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous height 1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

Let \mathfrak{B} be a MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous height 1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$,*
- 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

Let \mathfrak{B} be a MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous height 1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$,*
- 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*
- 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

Let \mathfrak{B} be a MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous height 1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$,*
 - 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*
 - 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,*
 - 4. $\text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a colour-canonical pseudo-Siggers,*
- and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.*

Summing up:

Theorem

Let \mathfrak{B} be a MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous height 1 homomorphism $\text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$,*
- 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*
- 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,*
- 4. $\text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a colour-canonical pseudo-Siggers,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Items 3. and 4. can be checked effectively.

- ▶ When is $\text{CSP}(\mathfrak{B})$ in Datalog? Is it decidable?
Answered for **monochromatic** obstructions.
- ▶ MMSNP_2 : instead of colouring vertices, we colour edges. It is **more expressive** than MMSNP , but it is open whether it has a complexity dichotomy (Lutz et al.).
Example: is it possible to colour the edges of an input graph and avoid:

