

Infinite-domain CSPs: part 1

Why? What?

Why?

- Linear programming, diophantine equations, semi-definite programs, ...

Why?

- ~~Linear programming, diophantine equations, semi-definite programs, ...~~

Why?

- ~~Linear programming, diophantine equations, semi-definite programs, ...~~
- Why CSPs with **finite** domains?
 k -coloring, Boolean satisfiability, linear equations over \mathbb{Z}_p , ...

Why?

- ~~Linear programming, diophantine equations, semi-definite programs, ...~~
- Why CSPs with **finite** domains?
 k -coloring, Boolean satisfiability, linear equations over \mathbb{Z}_p , ...

Finite-domain CSPs are not studied because of individual problems. Why then?


1. Hope for a general understanding of a **class** of problems
Want to understand **why** problems are easy/hard
2. Connection between tractability and closure properties
3. Induction
4. Natural class due to connection with Monotone Monadic SNP without inequalities (MMSNP)

Why?

- Linear programming, diophantine equations, semi-definite programs, ...
- Why CSPs with **finite** domains?
 k -coloring, Boolean satisfiability, linear equations over \mathbb{Z}_p , ...

Finite-domain CSPs are not studied because of individual problems. Why then?

1. Hope for a general understanding of a **class** of problems
Want to understand **why** problems are easy/hard
2. Connection between tractability and closure properties
3. Induction
4. Natural class due to connection with Monotone Monadic SNP without inequalities (MMSNP)



Template-free CSPs

$\text{CSP}(\mathbb{A})$

Given \mathbb{X} , does there exist a homomorphism $\mathbb{X} \rightarrow \mathbb{A}$?

Template-free CSPs

$\text{CSP}(\mathbb{A})$

Given \mathbb{X} , does there exist a homomorphism $\mathbb{X} \rightarrow \mathbb{A}$? $\leftrightarrow \mathcal{C} = \{\mathbb{X} \mid \exists h: \mathbb{X} \rightarrow \mathbb{A}\}$

$\text{CSP}(\mathbb{A})$

Given \mathbb{X} , does there exist a homomorphism $\mathbb{X} \rightarrow \mathbb{A}$? $\leftrightarrow \mathcal{C} = \{\mathbb{X} \mid \exists h: \mathbb{X} \rightarrow \mathbb{A}\}$

Let \mathcal{C} be a class of "instances".

When is this the set of yes-instances of a non-uniform CSP?

$\text{CSP}(\mathbb{A})$

Given \mathbb{X} , does there exist a homomorphism $\mathbb{X} \rightarrow \mathbb{A}$? $\leftrightarrow \mathcal{C} = \{\mathbb{X} \mid \exists h: \mathbb{X} \rightarrow \mathbb{A}\}$

Let \mathcal{C} be a class of "instances".

When is this the set of yes-instances of a non-uniform CSP?

Observation.

$\exists \mathbb{A} : \mathcal{C} = \text{CSP}(\mathbb{A})$ iff \mathcal{C} is closed under disjoint unions and relaxations (inverse homomorphisms)

$\text{CSP}(\mathbb{A})$

Given \mathbb{X} , does there exist a homomorphism $\mathbb{X} \rightarrow \mathbb{A}$? $\leftrightarrow \mathcal{C} = \{\mathbb{X} \mid \exists h: \mathbb{X} \rightarrow \mathbb{A}\}$

Let \mathcal{C} be a class of "instances".

When is this the set of yes-instances of a non-uniform CSP?




Observation.

$\exists \mathbb{A} : \mathcal{C} = \text{CSP}(\mathbb{A})$ iff \mathcal{C} is closed under disjoint unions and relaxations (inverse homomorphisms)

- Take $\mathbb{A} = \bigsqcup_{\mathbb{X} \in \mathcal{C}} \mathbb{X}$
- Can be a useful point of view, but most likely not




Template-free CSPs: 3-coloring

Goal: understand finite 3-colorable graphs



Template-free CSPs: 3-coloring




Goal: understand finite 3-colorable graphs



Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ is 3-colorable}\}$

Template-free CSPs: 3-coloring




Goal: understand finite 3-colorable graphs



Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ is 3-colorable}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Goal: understand finite 3-colorable graphs






Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ is 3-colorable}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Step 3: $\text{CSP}(G^*)$ is the problem of deciding “ $G \in \mathcal{C}$?”

Goal: understand finite 3-colorable graphs



Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ is 3-colorable}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$




Step 3: $\text{CSP}(G^*)$ is the problem of deciding " $G \in \mathcal{C}$?"



Step 4: $G^* \rightleftharpoons K_3$ so $\text{CSP}(G^*) = \text{CSP}(K_3)$




Template-free CSPs: Acyclicity

Goal: understand finite acyclic directed graphs



Template-free CSPs: Acyclicity


Goal: understand finite acyclic directed graphs



Step 1: $\mathcal{C} = \{G \text{ finite digraph } | G \text{ acyclic}\}$

Template-free CSPs: Acyclicity

Goal: understand finite acyclic directed graphs




Step 1: $\mathcal{C} = \{G \text{ finite digraph } | G \text{ acyclic}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Template-free CSPs: Acyclicity

Goal: understand finite acyclic directed graphs




Step 1: $\mathcal{C} = \{G \text{ finite digraph } | G \text{ acyclic}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Step 3: $\text{CSP}(G^*)$ is the problem of deciding “ $G \in \mathcal{C}$?”

Template-free CSPs: Acyclicity

Goal: understand finite acyclic directed graphs




Step 1: $\mathcal{C} = \{G \text{ finite digraph } | G \text{ acyclic}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Step 3: $\text{CSP}(G^*)$ is the problem of deciding " $G \in \mathcal{C}$?"


 Step 4: $\text{CSP}(G^*) = \text{CSP}(\mathbb{N}; <)$. Acyclic digraphs are those with a topological ordering.

Template-free CSPs: Monochromatic triangle



Goal: understand \exists 2-edge colorings without monochromatic triangle

Template-free CSPs: Monochromatic triangle



Goal: understand \exists 2-edge colorings without monochromatic triangle

Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ 2-edge-colorable without monochromatic triangle}\}$


Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Step 3: $\text{CSP}(G^*)$ is the problem of deciding “ $G \in \mathcal{C}$?”



Step 4: $G^* \rightleftharpoons ?$

Template-free CSPs: Monochromatic triangle



Goal: understand \exists 2-edge colorings without monochromatic triangle

Step 1: $\mathcal{C} = \{G \text{ finite graph } | G \text{ 2-edge-colorable without monochromatic triangle}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$



Step 3: $\text{CSP}(G^*)$ is the problem of deciding “ $G \in \mathcal{C}$?”



Step 4: $G^* \rightleftharpoons ?$

Observation. $\forall n \geq 1 \exists G$ triangle-free that cannot be partitioned into n independent sets.

Template-free CSPs: cyclic orientability





Goal: understand graphs that can be oriented in a way that every triangle becomes a directed cycle

Step 1: $\mathcal{C} = \{G \text{ finite graph} \mid G \text{ cyclically orientable}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

Step 3: $\text{CSP}(G^*)$ is the problem of deciding “ $G \in \mathcal{C}$?”

Step 4: $G^* \rightleftharpoons ?$



Goal: understand graphs that can be oriented in a way that every triangle becomes a directed cycle

Step 1: $\mathcal{C} = \{G \text{ finite graph} \mid G \text{ cyclically orientable}\}$

Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$


Step 3: $\text{CSP}(G^*)$ is the problem of deciding " $G \in \mathcal{C}$?"



Step 4: $G^* \rightleftharpoons ?$


Observation. $\forall n \geq 1 \exists G$ triangle-free that cannot be partitioned into n independent sets.

Template-free CSPs: cyclic orientability




Can one not see this as a label cover instance?


Template-free CSPs: cyclic orientability




Can one not see this as a label cover instance?




Template-free CSPs: cyclic orientability




Can one not see this as a label cover instance?





Can one not see this as a label cover instance?



This potato is implied, added automatically:

- 1 constraint per triangle
- This type of constraint is not accessible as a “real” ternary constraint

Similarities between these problems

All problems so far:

- label cover over a finite set
- some **structural** constraints are "automatically" added
- structural constraints are not part of the "normal" constraints
- scope of structural constraints is of **bounded** size
- Except for 3-coloring, not $\text{CSP}(\mathbb{A})$ for a finite \mathbb{A} , yet CSPs of bad structures (disjoint unions)

Similarities between these problems

All problems so far:





- label cover over a finite set
- some **structural** constraints are "automatically" added
- structural constraints are not part of the "normal" constraints
- scope of structural constraints is of **bounded** size
- Except for 3-coloring, not $\text{CSP}(\mathbb{A})$ for a finite \mathbb{A} , yet CSPs of bad structures (disjoint unions)

What templates to study these problems?

How to study the complexity of the CSPs using these templates?

Orbit-finiteness,
reducts of finitely bounded homogeneous structures





Template-free CSPs: cyclic orientability



Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$


This step is wasteful in terms of vertices and edges

Template-free CSPs: cyclic orientability







Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

This step is wasteful in terms of vertices and edges




Template-free CSPs: cyclic orientability







Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$

This step is wasteful in terms of vertices and edges




Template-free CSPs: cyclic orientability




Step 2: Take $G^* = \bigsqcup_{G \in \mathcal{C}} G$


This step is wasteful in terms of vertices and edges



\oplus



$=$



Template-free CSPs: cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$ (where $(V; E) * (V; A) = (V; E, A)$)


structure with two edge relations



Template-free CSPs: cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$ (where $(V; E) * (V; A) = (V; E, A)$)


\mathcal{V} has the (free) amalgamation property:



structure with two edge relations

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$ (where $(V; E) * (V; A) = (V; E, A)$)

\mathcal{V} has the (free) amalgamation property:



Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:


- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

structure with two edge relations

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$ (where $(V; E) * (V; A) = (V; E, A)$)

\mathcal{V} has the (free) amalgamation property:



Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$. This is called **homogeneity**

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

Template for cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$

Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

Template for cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$


Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

Here:

- $G^* = (V; E, A)$ is an infinite graph with cyclic orientation



$$G^* = (V; E, A)$$

Template for cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$


Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

Here:

- $G^* = (V; E, A)$ is an infinite graph with cyclic orientation
- $G = (V; E)$ is a **reduct** of G^*



Template for cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$


Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .

Here:

- $G^* = (V; E, A)$ is an infinite graph with cyclic orientation
- $G = (V; E)$ is a **reduct** of G^*
- The finite subgraphs of $(V; E)$ are exactly the finite graphs that admit *some* cyclic orientation



Template for cyclic orientability

Define $\mathcal{V} = \{G * \vec{G} \mid \vec{G} \text{ cyclic orientation of } G\}$


Theorem (Fraïssé). If \mathcal{V} is closed under substructures and has the amalgamation property, there exists G^* such that:

- The finite substructures of G^* are exactly those in \mathcal{V}
- For any isomorphism $f: G \rightarrow G'$ between finite substructures of G^* , there exists $\alpha \in \text{Aut}(G^*)$ s.t. $\alpha|_G = f$.

G^* is unique up to isomorphism: it is the **Fraïssé limit** of \mathcal{V} .


Here:

- $G^* = (V; E, A)$ is an infinite graph with cyclic orientation
- $G = (V; E)$ is a **reduct** of G^*
- The finite subgraphs of $(V; E)$ are exactly the finite graphs that admit *some* cyclic orientation
- $\text{CSP}(V; E) = \mathcal{C} = \text{CSP}(\bigsqcup_{G \in \mathcal{C}} G)$



Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

Homogeneity

$$\overset{\nu}{\bullet} \simeq \overset{\nu'}{\bullet} \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(\nu) = \nu'$$




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

Homogeneity

$$\bullet \stackrel{v}{\sim} \bullet \stackrel{v'}{\sim} \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(v) = v'$$

$$e \stackrel{e}{\sim} e' \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(e) = e'$$




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

Homogeneity

$$\bullet \stackrel{v}{\sim} \bullet \stackrel{v'}{\sim} \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(v) = v'$$

$$e \stackrel{e}{\sim} e' \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(e) = e'$$



Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

Homogeneity


$$\bullet \stackrel{\nu}{\sim} \bullet \stackrel{\nu'}{\sim} \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(\nu) = \nu'$$

$$\begin{array}{c} e \\ \downarrow \\ \bullet \end{array} \simeq \begin{array}{c} e' \\ \downarrow \\ \bullet \end{array} \Rightarrow \exists \alpha \in \text{Aut}(G^*) : \alpha(e) = e'$$

Finitely many directed graphs on n vertices
 $\rightsquigarrow \text{Aut}(G^*) \curvearrowright V^n$ has finitely many orbits

$\text{Aut}(G^*)$ is **oligomorphic**



G^* is **ω -categorical**



Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?


$CSP(V; E)$ as label cover




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

$CSP(V; E)$ as label cover





Elements of E




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

$CSP(V; E)$ as label cover




Infinitely many vertices/edges to choose from




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

$CSP(V; E)$ as label cover




Infinitely many vertices/edges to choose from
Up to automorphisms, only two solutions




Template for cyclic orientability

Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

$CSP(V; E)$ as label cover




Infinitely many vertices/edges to choose from
Up to automorphisms, only two solutions



Template for cyclic orientability


Why is G^* “better” than $\bigsqcup_{G \in \mathcal{C}} G$?

$CSP(V; E)$ as label cover



Infinitely many vertices/edges to choose from

Up to automorphisms, only two solutions



Conjecture. Let \mathcal{C} be a **CSP**

Then \mathcal{C} is in **P** or **NP-complete**.

Template-free CSPs: The conjecture


Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

Then \mathcal{C} is in **P** or **NP-complete**.

Template-free CSPs: The conjecture

Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class \mathcal{K}** (the **certificates**) and a **verifier $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$** such that:

Then \mathcal{C} is in **P** or **NP-complete**.




Template-free CSPs: The conjecture

Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)


Then \mathcal{C} is in **P** or **NP-complete**.



Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$ (bigger certificates are fine)

Then \mathcal{C} is in P or NP-complete.




(\subseteq means non-induced substructure)

Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$ (bigger certificates are fine)
- $\forall \mathbb{B} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{A} \subseteq \mathbb{B}$, $\mathbb{A} * \mathbb{K}|_{\mathbb{A}} \in \mathcal{V}$ (smaller instances are fine)

Then \mathcal{C} is in P or NP-complete.




(\subseteq means non-induced substructure)

Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $A \in \mathcal{C}$, there exists $K \in \mathcal{K}$ such that $A * K \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall A * K \in \mathcal{V}$ and $K \hookrightarrow L \in \mathcal{K}$, $\exists A \subseteq B \in \mathcal{K}$ such that $B * L \in \mathcal{V}$ (bigger certificates are fine)
- $\forall B * K \in \mathcal{V}$ and $A \subseteq B$, $A * K|_A \in \mathcal{V}$ (smaller instances are fine)
- $\forall K \in \mathcal{K}$, $\{A \in \mathcal{C} \mid A * K \in \mathcal{V}\}$ has a maximal element

Then \mathcal{C} is in P or NP-complete.



(\subseteq means non-induced substructure)


Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$ (bigger certificates are fine)
- $\forall \mathbb{B} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{A} \subseteq \mathbb{B}$, $\mathbb{A} * \mathbb{K}|_{\mathbb{A}} \in \mathcal{V}$ (smaller instances are fine)
- $\forall \mathbb{K} \in \mathcal{K}$, $\{\mathbb{A} \in \mathcal{C} \mid \mathbb{A} * \mathbb{K} \in \mathcal{V}\}$ has a maximal element

Then \mathcal{C} is in P or NP-complete.

Under those conditions:

- $\mathcal{C} = \text{CSP}(\mathbb{A})$, where \mathbb{A} is a **reduct of a finitely bounded homogeneous structure** \mathbb{A}^* .
- $\text{Aut}(\mathbb{A})$ is oligomorphic



(\subseteq means non-induced substructure)


Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$ (bigger certificates are fine)
- $\forall \mathbb{B} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{A} \subseteq \mathbb{B}$, $\mathbb{A} * \mathbb{K}|_{\mathbb{A}} \in \mathcal{V}$ (smaller instances are fine)
- $\forall \mathbb{K} \in \mathcal{K}$, $\{\mathbb{A} \in \mathcal{C} \mid \mathbb{A} * \mathbb{K} \in \mathcal{V}\}$ has a maximal element

Then \mathcal{C} is in P or NP-complete.

Examples:

- order CSPs, phylogenetic CSPs
- **GMSNP**: problems of coloring edges of a graph while avoiding forbidden configurations



(\subseteq means non-induced substructure)


Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$ (yes-instances have a certificate)
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$ (bigger certificates are fine)
- $\forall \mathbb{B} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{A} \subseteq \mathbb{B}$, $\mathbb{A} * \mathbb{K}|_{\mathbb{A}} \in \mathcal{V}$ (smaller instances are fine)
- $\forall \mathbb{K} \in \mathcal{K}$, $\{\mathbb{A} \in \mathcal{C} \mid \mathbb{A} * \mathbb{K} \in \mathcal{V}\}$ has a maximal element

Then \mathcal{C} is in P or NP-complete.

Non-examples:

- \mathcal{C} = satisfiable systems of affine equations over \mathbb{Z}
- ... feasible linear programs
- ... feasible SDPs



(\subseteq means non-induced substructure)

The conjecture is proved...

The conjecture is proved...

... when \mathcal{K} is the class of:

- maps to a finite set

(Bulatov, Zhuk)

The conjecture is proved...

... when \mathcal{K} is the class of:

- maps to a finite set (Bulatov, Zhuk)
- linear orders (Bodirsky, Kára)
- phylogenetic trees (Bodirsky, Jonsson, Pham)
- graphs (Bodirsky, Pinsker)
- K_n -free graphs for some $n \geq 3$ (Bodirsky, Martin, Pinsker, Pongrácz)
- equivalence relations (with c classes of size s) (Kompatscher, Pham)
- partial orders (Bodirsky, M.)
- unary structures (M., Pinsker)
- tournaments (M., Nagy, Pinsker)
- (K_n^r -free) r -uniform hypergraphs

The conjecture is proved...

... when \mathcal{K} is the class of:

- maps to a finite set (Bulatov, Zhuk)
- linear orders (Bodirsky, Kára)
- phylogenetic trees (Bodirsky, Jonsson, Pham)
- graphs (Bodirsky, Pinsker)
- K_n -free graphs for some $n \geq 3$ (Bodirsky, Martin, Pinsker, Pongrácz)
- equivalence relations (with c classes of size s) (Kompatscher, Pham)
- partial orders (Bodirsky, M.)
- unary structures (M., Pinsker)
- tournaments (M., Nagy, Pinsker)
- (K_n^r -free) r -uniform hypergraphs

... and in several other cases where also \mathcal{V} is restricted:

- MMSNP (Feder, Vardi) (Bodirsky, Madelaine, M.)
- Orientability with forbidden tournaments (Bodirsky, Guzmán-Pro) (Feller, Pinsker) (Bitter, M.)

Algebra & "Topology"

Polymorphisms

Definition. A **polymorphism** of \mathbb{A} is a function $f: \mathbb{A}^n \rightarrow \mathbb{A}$ such that for every relation $R^{\mathbb{A}}$ we have

$$\mathbf{a}^1, \dots, \mathbf{a}^n \in R^{\mathbb{A}} \implies f(\mathbf{a}^1, \dots, \mathbf{a}^n) \in R^{\mathbb{A}}$$
$$\qquad\qquad\qquad ||$$
$$(f(a_1^1, \dots, a_1^n), \dots, f(a_r^1, \dots, a_r^n))$$

Polymorphisms

Definition. A **polymorphism** of \mathbb{A} is a function $f: \mathbb{A}^n \rightarrow \mathbb{A}$ such that for every relation $R^\mathbb{A}$ we have

$$\mathbf{a}^1, \dots, \mathbf{a}^n \in R^\mathbb{A} \implies f(\mathbf{a}^1, \dots, \mathbf{a}^n) \in R^\mathbb{A}$$

Even if \mathbb{A} is infinite:

- If $h_1, \dots, h_n: \mathbb{X} \rightarrow \mathbb{A}$ are solutions, $f \circ (h_1, \dots, h_n)$ is a solution
- $\text{Pol}(\mathbb{A})$ is a clone (contains projections, closed under composition)
- If $R \subseteq A^r$ is **pp-definable**, then R is preserved by $\text{Pol}(\mathbb{A})$


Polymorphisms

Definition. A **polymorphism** of \mathbb{A} is a function $f: \mathbb{A}^n \rightarrow \mathbb{A}$ such that for every relation $R^\mathbb{A}$ we have

$$\mathbf{a}^1, \dots, \mathbf{a}^n \in R^\mathbb{A} \implies f(\mathbf{a}^1, \dots, \mathbf{a}^n) \in R^\mathbb{A}$$

Even if \mathbb{A} is infinite:

- If $h_1, \dots, h_n: \mathbb{X} \rightarrow \mathbb{A}$ are solutions, $f \circ (h_1, \dots, h_n)$ is a solution
- $\text{Pol}(\mathbb{A})$ is a clone (contains projections, closed under composition)
- If $R \subseteq A^r$ is **pp-definable**, then R is preserved by $\text{Pol}(\mathbb{A})$



projection of the solution set of a CSP instance

Lemma. Let \mathbb{X} be a finite structure, $x_1, \dots, x_n \in X$.

There exists a pp-formula $\varphi_{\mathbb{X}}(x_1, \dots, x_n)$ such that for all \mathbb{A} :

$$\mathbb{A} \models \varphi_{\mathbb{X}}(a_1, \dots, a_n) \iff \exists h: \mathbb{X} \rightarrow \mathbb{A} \forall i : h(x_i) = a_i$$

Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$. Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable

$$\begin{array}{ccccccccc} & & \mathbf{a}^1 & & \mathbf{a}^2 & & & & \mathbf{a}^n \\ & & || & & || & & & & || \\ \mathbf{b}^1 = & \left(\begin{array}{cccc} a_1^1 & a_1^2 & \cdots & a_1^n \end{array} \right) & & & & & & & \end{array}$$

$$\begin{array}{ccc} \vdots & \vdots & \vdots \end{array}$$


$$\mathbf{b}^r = \left(\begin{array}{cccc} a_r^1 & a_r^2 & \cdots & a_r^n \end{array} \right)$$

Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$
- Consider a chain $B \subseteq B_2 \subseteq \dots$ with $\bigcup B_i = A^n$
Then $R_1 \supseteq R_2 \supseteq \dots$




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$
- Consider a chain $B \subseteq B_2 \subseteq \dots$ with $\bigcup B_i = A^n$
Then $R_1 \supseteq R_2 \supseteq \dots$
- Every finite $\bigcap R_i \neq \emptyset$, so $\bigcap_{i \geq 1} R_i \neq \emptyset$.
This is the **standard compactness argument**.




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$
- Consider a chain $B \subseteq B_2 \subseteq \dots$ with $\bigcup B_i = A^n$
Then $R_1 \supseteq R_2 \supseteq \dots$
- Every finite $\bigcap R_i \neq \emptyset$, so $\bigcap_{i \geq 1} R_i \neq \emptyset$.
This is the **standard compactness argument**.
- For some finite B , $\varphi_{\mathbb{B}}$ defines $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{A}^n \rightarrow \mathbb{A}\}$




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$
- Consider a chain $B \subseteq B_2 \subseteq \dots$ with $\bigcup B_i = A^n$
Then $R_1 \supseteq R_2 \supseteq \dots$
- Every finite $\bigcap R_i \neq \emptyset$, so $\bigcap_{i \geq 1} R_i \neq \emptyset$.
This is the **standard compactness argument**.
- For some finite B , $\varphi_{\mathbb{B}}$ defines $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{A}^n \rightarrow \mathbb{A}\}$
- If $\mathbf{a}^1, \dots, \mathbf{a}^n$ generate R under $\text{Pol}(\mathbb{A})$, this is exactly R




Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

- For $B \subseteq A^n$ finite and $\mathbf{a}^1, \dots, \mathbf{a}^n \in A^r$,
 $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$ is pp-definable
- $\varphi_{\mathbb{B}}$ defines $R_1 = \{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{B} \rightarrow \mathbb{A}\}$
- Subset of A^r closed under $\text{Aut}(\mathbb{A})$
- Consider a chain $B \subseteq B_2 \subseteq \dots$ with $\bigcup B_i = A^n$
Then $R_1 \supseteq R_2 \supseteq \dots$
- Every finite $\bigcap R_i \neq \emptyset$, so $\bigcap_{i \geq 1} R_i \neq \emptyset$.
This is the **standard compactness argument**.
- For some finite B , $\varphi_{\mathbb{B}}$ defines $\{f(\mathbf{a}^1, \dots, \mathbf{a}^n) \mid f: \mathbb{A}^n \rightarrow \mathbb{A}\}$
- If $\mathbf{a}^1, \dots, \mathbf{a}^n$ generate R under $\text{Pol}(\mathbb{A})$, this is exactly R
- We can always find finitely many generators, by ω -categoricity



Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$. Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

Corollary. If \mathbb{A} is ω -categorical, the complexity of $\text{CSP}(\mathbb{A})$ only depends on $\text{Pol}(\mathbb{A})$.

$$\text{Pol}(\mathbb{A}) \subseteq \text{Pol}(\mathbb{B}) \implies \text{CSP}(\mathbb{B}) \leq_P \text{CSP}(\mathbb{A})$$

Polymorphisms and pp-definability

Theorem (Bodirsky, Nešetřil). Let \mathbb{A} be ω -categorical and $R \subseteq A^r$.

Then R is pp-definable in \mathbb{A} iff R is preserved by $\text{Pol}(\mathbb{A})$.

Corollary. If \mathbb{A} is ω -categorical, the complexity of $\text{CSP}(\mathbb{A})$ only depends on $\text{Pol}(\mathbb{A})$.

$$\text{Pol}(\mathbb{A}) \subseteq \text{Pol}(\mathbb{B}) \implies \text{CSP}(\mathbb{B}) \leq_P \text{CSP}(\mathbb{A})$$

This order is way too fine, no hope of understanding it all (already continuum many clones on $\{0, 1, 2\}$)

Definition (Minion/clone homomorphism). Arity-preserving map $\xi: \text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ such that:

- minor conditions are preserved
- all equational conditions are preserved


(minion homomorphism)

(clone homomorphism)

Definition (Minion/clone homomorphism). Arity-preserving map $\xi: \text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ such that:

- minor conditions are preserved
- all equational conditions are preserved

(minion homomorphism)
(clone homomorphism)




Clones on $\{0, 1\}$ up to \subseteq

Definition (Minion/clone homomorphism). Arity-preserving map $\xi: \text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ such that:

- minor conditions are preserved
- all equational conditions are preserved

(minion homomorphism)
(clone homomorphism)



Definition. $f_n \rightarrow f$ if for all **finite** $S \subseteq A$, $\exists N \forall n \geq N : f_n|_S = f|_S$

This turns $\text{Pol}(\mathbb{A})$ into a **topological clone** with nice properties:

- It is closed: if $f_n \rightarrow f$ and $\{f_n : n \geq 1\} \subseteq \text{Pol}(\mathbb{A})$, then $f \in \text{Pol}(\mathbb{A})$
- If \mathbb{A} is ω -categorical, $\text{Pol}(\mathbb{A})^{(n)}/\text{Aut}(\mathbb{A})$ is **compact**

(Bodirsky, Pinsker)

Definition. $f_n \rightarrow f$ if for all **finite** $S \subseteq A$, $\exists N \forall n \geq N : f_n|_S = f|_S$

This turns $\text{Pol}(\mathbb{A})$ into a **topological clone** with nice properties:

- It is closed: if $f_n \rightarrow f$ and $\{f_n : n \geq 1\} \subseteq \text{Pol}(\mathbb{A})$, then $f \in \text{Pol}(\mathbb{A})$
- If \mathbb{A} is ω -categorical, $\text{Pol}(\mathbb{A})^{(n)}/\text{Aut}(\mathbb{A})$ is **compact**

(Bodirsky, Pinsker)

Theorem (Barto, Opršal, Pinsker). Let \mathbb{A} be ω -categorical, and \mathbb{B} be finite.

The following are equivalent:

- There exists a **uniformly continuous** minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$
- \mathbb{B} is pp-constructible in \mathbb{A} , and $\text{CSP}(\mathbb{B}) \leq_P \text{CSP}(\mathbb{A})$.

Definition. $f_n \rightarrow f$ if for all **finite** $S \subseteq A$, $\exists N \forall n \geq N : f_n|_S = f|_S$

This turns $\text{Pol}(\mathbb{A})$ into a **topological clone** with nice properties:

- It is closed: if $f_n \rightarrow f$ and $\{f_n : n \geq 1\} \subseteq \text{Pol}(\mathbb{A})$, then $f \in \text{Pol}(\mathbb{A})$
- If \mathbb{A} is ω -categorical, $\text{Pol}(\mathbb{A})^{(n)}/\text{Aut}(\mathbb{A})$ is **compact**

(Bodirsky, Pinsker)

Theorem (Barto, Opršal, Pinsker). Let \mathbb{A} be ω -categorical, and \mathbb{B} be finite.

The following are equivalent:

- There exists a **uniformly continuous** minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$
- \mathbb{B} is pp-constructible in \mathbb{A} , and $\text{CSP}(\mathbb{B}) \leq_P \text{CSP}(\mathbb{A})$.

$$\begin{array}{ccc} \text{Pol}(\mathbb{A}) & \xrightarrow{\xi} & \text{Pol}(\mathbb{B}) \\ \text{restr}_F \searrow & \nearrow & \\ & \text{Pol}(\mathbb{F}, \mathbb{A}) & \end{array}$$

The conjecture (again)

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure.

Suppose

- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in \mathcal{P} .


The conjecture (again)

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure.

Suppose

- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in P.




The conjecture (again)

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure.

Suppose

- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in P.




The conjecture (again)

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure.

Suppose

- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in P.




The conjecture (again)

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure.

Suppose

- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in P.



Answer 1: Topology is irrelevant (for clone homomorphisms to $\text{Pol}(\mathbb{K}_3)$)

Theorem (Barto, Pinsker). Let \mathbb{B} be an ω -categorical structure that is a **core**.

The following are equivalent:

- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no **uniformly continuous** clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- $\text{Pol}(\mathbb{B})$ has a 6-ary operation s satisfying

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y) \quad (\text{pseudo-Siggers modulo } \text{Aut}(\mathbb{B}))$$


Answer 1: Topology is irrelevant (for clone homomorphisms to $\text{Pol}(\mathbb{K}_3)$)

Theorem (Barto, Pinsker). Let \mathbb{B} be an ω -categorical structure that is a **core**.

The following are equivalent:

- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no **uniformly continuous** clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- $\text{Pol}(\mathbb{B})$ has a 6-ary operation s satisfying

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y) \quad (\text{pseudo-Siggers modulo } \text{Aut}(\mathbb{B}))$$



Loop lemma


Answer 1: Topology is irrelevant (for clone homomorphisms to $\text{Pol}(\mathbb{K}_3)$)

Theorem (Barto, Pinsker). Let \mathbb{B} be an ω -categorical structure that is a **core**.

The following are equivalent:

- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no **uniformly continuous** clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- $\text{Pol}(\mathbb{B})$ has a 6-ary operation s satisfying

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y) \quad (\text{pseudo-Siggers modulo } \text{Aut}(\mathbb{B}))$$



Answer 1: Topology is irrelevant (for clone homomorphisms to $\text{Pol}(\mathbb{K}_3)$)




Theorem (Barto, Pinsker). Let \mathbb{B} be an ω -categorical structure that is a **core**.

The following are equivalent:

- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no **uniformly continuous** clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- $\text{Pol}(\mathbb{B})$ has a 6-ary operation s satisfying

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y)$$

(pseudo-Siggers modulo $\text{Aut}(\mathbb{B})$)



Loop lemma

Free algebra

Answer 1: Topology is irrelevant (for clone homomorphisms to $\text{Pol}(\mathbb{K}_3)$)

Theorem (Barto, Pinsker). Let \mathbb{B} be an ω -categorical structure that is a **core**.

The following are equivalent:

- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no **uniformly continuous** clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- For every finite tuple c , $\text{Pol}(\mathbb{B}, c)$ has no clone homomorphism to $\text{Pol}(\mathbb{K}_3)$
- $\text{Pol}(\mathbb{B})$ has a 6-ary operation s satisfying

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y) \quad (\text{pseudo-Siggers modulo } \text{Aut}(\mathbb{B}))$$

Several improvements recently:

- Loop lemma without the constants
(also interesting for finite structures!) (Barto, Bodor, Kozik, M., Pinsker)
- Loop lemma for directed graphs
 \rightsquigarrow **4-ary** pseudo-Siggers (Brunar, Kozik, Nagy, Pinsker)


Answer 2: Topology is relevant (for minion homomorphisms to $\text{Pol}(K_3)$)

Theorem (Gillibert, Jonušas, Kompatscher, M., Pinsker).

There exists an ω -categorical \mathbb{A} with finite signature such that:

- $\text{Pol}(\mathbb{A})$ does not satisfy any minor condition
 - ~ \exists minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$
- $\text{Pol}(\mathbb{B}, \mathbb{A})$ satisfies a non-trivial minor condition for every finite $\mathbb{B} \subseteq \mathbb{A}$
 - ~ \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$

Note: \mathbb{A} not subject to the dichotomy conjecture



Conjecture. No uniformly continuous minion/clone homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$ implies $\text{CSP}(\mathbb{A})$ in P .

Conjecture. No uniformly continuous minion/clone homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$ implies $\text{CSP}(\mathbb{A})$ in P.

Theorem. For a finite \mathbb{A} , the following are equivalent:

- $\text{CSP}(\mathbb{A})$ is in P
- \nexists minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$
- $\text{Pol}(\mathbb{A})$ has a 6-ary operation (Siggers)

$$s(x, y, x, z, y, z) \approx s(y, x, z, x, z, y)$$

- $\text{Pol}(\mathbb{A})$ has for some $k \geq 3$ a weak near unanimity operation (Maróti, McKenzie)

$$w(x, \dots, x, x, y) \approx w(x, \dots, x, y, x) \approx \dots \approx w(y, x, \dots, x, x)$$

- $\text{Pol}(\mathbb{A})$ has for all large prime $n \geq 3$ a cyclic operation (Barto, Kozik)

$$c(x_1, \dots, x_n) \approx c(x_2, \dots, x_n, x_1)$$

Theorem (Bodirsky, M., Olšák, Opršal, Pinsker, Willard).

\forall minor condition Σ , \exists reduct \mathbb{A} of a finitely bounded homogeneous structure such that:

- $\text{Pol}(\mathbb{A})$ satisfies some non-trivial minor condition
- $\text{Pol}(\mathbb{A})$ does not satisfy Σ
- $\text{CSP}(\mathbb{A})$ is in P (even in AC^0)

Easy CSPs without given identities
(within the conjecture)

Identities and complexity

Theorem (Bodirsky, M., Olšák, Opršal, Pinsker, Willard).

\forall minor condition Σ , \exists reduct \mathbb{A} of a finitely bounded homogeneous structure such that:

- $\text{Pol}(\mathbb{A})$ satisfies some non-trivial minor condition
- $\text{Pol}(\mathbb{A})$ does not satisfy Σ
- $\text{CSP}(\mathbb{A})$ is in P (even in AC^0)

Easy CSPs without given identities
(within the conjecture)

Theorem (Gillibert, Kompatscher, Jonušas, M., Pinsker).

\forall equational condition Σ true for all CSPs in AC^0 ,
 $\exists \mathbb{A}$ ω -categorical s.t. $\text{Pol}(\mathbb{A}) \models \Sigma$ and $\text{CSP}(\mathbb{A})$ has arbitrary Turing degree.

Hard CSPs with given identities
(outside the conjecture)

Identities and complexity

Theorem (Bodirsky, M., Olšák, Opršal, Pinsker, Willard).

\forall minor condition Σ , \exists reduct \mathbb{A} of a finitely bounded homogeneous structure such that:

- $\text{Pol}(\mathbb{A})$ satisfies some non-trivial minor condition
- $\text{Pol}(\mathbb{A})$ does not satisfy Σ
- $\text{CSP}(\mathbb{A})$ is in P (even in AC^0)

Easy CSPs without given identities
(within the conjecture)

Theorem (Gillibert, Kompatscher, Jonušas, M., Pinsker).

\forall equational condition Σ true for all CSPs in AC^0 ,
 $\exists \mathbb{A}$ ω -categorical s.t. $\text{Pol}(\mathbb{A}) \models \Sigma$ and $\text{CSP}(\mathbb{A})$ has arbitrary Turing degree.

Hard CSPs with given identities
(outside the conjecture)

Other bad news:

- Bounded width cannot be characterized by identities
(within the conjecture)
- Non-trivial $\text{Pol}(\mathbb{A})$ w/o pseudo cyclic polymorphisms
(within the conjecture)
- Non-trivial $\text{Pol}(\mathbb{A})$ w/o pseudo weak near unanimity operations
(outside the conjecture)

(Bodirsky, Rydval)

(Bodirsky, Kára)

(Barto, Bodor, Kozik, M., Pinsker)


Summary: why?

Conjecture. Let \mathcal{C} be a **CSP** such that there exists a **finitely bounded Fraïssé class** \mathcal{K} (the **certificates**) and a **verifier** $\mathcal{V} \subseteq \mathcal{C} * \mathcal{K}$ such that:

- For all $\mathbb{A} \in \mathcal{C}$, there exists $\mathbb{K} \in \mathcal{K}$ such that $\mathbb{A} * \mathbb{K} \in \mathcal{V}$
- $\forall \mathbb{A} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{K} \hookrightarrow \mathbb{L} \in \mathcal{K}$, $\exists \mathbb{A} \subseteq \mathbb{B} \in \mathcal{K}$ such that $\mathbb{B} * \mathbb{L} \in \mathcal{V}$
- $\forall \mathbb{B} * \mathbb{K} \in \mathcal{V}$ and $\mathbb{A} \subseteq \mathbb{B}$, $\mathbb{A} * \mathbb{K}|_{\mathbb{A}} \in \mathcal{V}$
- $\forall \mathbb{K} \in \mathcal{K}$, $\{\mathbb{A} \in \mathcal{C} \mid \mathbb{A} * \mathbb{K} \in \mathcal{V}\}$ has a maximal element


(yes-instances have a certificate)
(bigger certificates are fine)
(smaller instances are fine)

Then \mathcal{C} is in P or NP-complete.



Summary: why?

1. Hope for a general understanding of a class of problems
2. Connection between tractability and closure properties
3. Induction + Compactness
4. Naturally defined class



Outline of Part 2

Smooth Approximations

How to prove tractability/hardness of such CSPs

Smooth Approximations 1
2022

M., Pinsker

Smooth Approximations 2
2021

M., Nagy, Pinsker, Wrona

Smooth Approximations 3
2024

M., Nagy, Pinsker

∞ CSPs and Finite PCSPs

Descriptive complexity: bounded width, first-order solvability ...

New algorithms for order CSPs and hypergraph CSPs

Conjecture (Bodirsky, Pinsker). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure with certificates \mathcal{K} . Suppose


- \nexists uniformly continuous minion homomorphism $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(K_3)$, or (equivalently)
- if \mathbb{B} is the core of \mathbb{A} , $\nexists c, \xi : \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ uniformly continuous clone homomorphism.

Then $\text{CSP}(\mathbb{A})$ is in P.

The conjecture is proved when \mathcal{K} is the class of:



- maps to a finite set (Bulatov, Zhuk)
- linear orders (Bodirsky, Kára)
- phylogenetic trees (Bodirsky, Jonsson, Pham)
- graphs (Bodirsky, Pinsker)
- K_n -free graphs for some $n \geq 3$ (Bodirsky, Martin, Pinsker, Pongrácz)
- equivalence relations (with c classes of size s)
- partial orders (Kompatscher, Pham)
- unary structures (Bodirsky, M.)
- tournaments (M., Pinsker)
- (K_n^r -free) r -uniform hypergraphs (M., Nagy, Pinsker)

The “reduction to the finite”: cyclic orientability





The “reduction to the finite”: cyclic orientability

G orientable iff the LC instance has a solution



The “reduction to the finite”: cyclic orientability



G orientable iff the LC instance has a solution



- Domain: structures in \mathcal{K} of bounded size
- Unary relations: original constraints
- Binary relations: matching substructures

The “reduction to the finite”: cyclic orientability

G orientable iff the LC instance has a solution



- Domain: structures in \mathcal{K} of bounded size
- Unary relations: original constraints
- Binary relations: matching substructures

Observation. For every reduct \mathbb{A} of a finitely bounded homogeneous structure with certificates \mathcal{K} , $\text{CSP}(\mathbb{A}) \leq_P \text{CSP}(\mathbb{A}_{fin})$.

Polymorphisms of \mathbb{A}_{fin} : Canonical Functions

Define $\equiv_{\mathcal{K}}$ as a binary relation on n -tuples such that

$$\mathbf{a} \equiv_{\mathcal{K}} \mathbf{b} \iff \mathbf{a} \text{ and } \mathbf{b} \text{ induce the same certificate } \mathbb{K} \in \mathcal{K}$$

This relation induces equality on the finite label cover instance

Polymorphisms of \mathbb{A}_{fin} : Canonical Functions

Define $\equiv_{\mathcal{K}}$ as a binary relation on n -tuples such that

$$\mathbf{a} \equiv_{\mathcal{K}} \mathbf{b} \iff \mathbf{a} \text{ and } \mathbf{b} \text{ induce the same certificate } \mathbb{K} \in \mathcal{K}$$

This relation induces equality on the finite label cover instance

Polymorphisms of \mathbb{A}_{fin} correspond to $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$: **canonical** functions

Polymorphisms of \mathbb{A}_{fin} : Canonical Functions

Define $\equiv_{\mathcal{K}}$ as a binary relation on n -tuples such that

$$\mathbf{a} \equiv_{\mathcal{K}} \mathbf{b} \iff \mathbf{a} \text{ and } \mathbf{b} \text{ induce the same certificate } \mathbb{K} \in \mathcal{K}$$

This relation induces equality on the finite label cover instance

Polymorphisms of \mathbb{A}_{fin} correspond to $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$: **canonical** functions

Proposition (Bodirsky, M.). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure with certificates \mathcal{K} . If \nexists clone homomorphism $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}}) \rightarrow \text{Pol}(\mathcal{K}_3)$, then $\text{CSP}(\mathbb{A})$ is in P.

Polymorphisms of \mathbb{A}_{fin} : Canonical Functions

Define $\equiv_{\mathcal{K}}$ as a binary relation on n -tuples such that

$$\mathbf{a} \equiv_{\mathcal{K}} \mathbf{b} \iff \mathbf{a} \text{ and } \mathbf{b} \text{ induce the same certificate } \mathbb{K} \in \mathcal{K}$$


This relation induces equality on the finite label cover instance

Polymorphisms of \mathbb{A}_{fin} correspond to $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$: **canonical** functions

Proposition (Bodirsky, M.). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure with certificates \mathcal{K} . If \nexists clone homomorphism $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}}) \rightarrow \text{Pol}(\mathcal{K}_3)$, then $\text{CSP}(\mathbb{A})$ is in P.

Question. Suppose that $\text{CSP}(\mathbb{A}, \equiv_{\mathcal{K}})$ is NP-hard.

- In which cases can we prove that $\text{CSP}(\mathbb{A})$ is NP-hard?
- What can we say otherwise?



Polymorphisms of \mathbb{A}_{fin} : Canonical Functions

Define $\equiv_{\mathcal{K}}$ as a binary relation on n -tuples such that

$$\mathbf{a} \equiv_{\mathcal{K}} \mathbf{b} \iff \mathbf{a} \text{ and } \mathbf{b} \text{ induce the same certificate } \mathbb{K} \in \mathcal{K}$$

This relation induces equality on the finite label cover instance

Polymorphisms of \mathbb{A}_{fin} correspond to $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$: **canonical** functions


Proposition (Bodirsky, M.). Let \mathbb{A} be a reduct of a finitely bounded homogeneous structure with certificates \mathcal{K} . If \nexists clone homomorphism $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}}) \rightarrow \text{Pol}(K_3)$, then $\text{CSP}(\mathbb{A})$ is in P.

Question. Suppose that $\text{CSP}(\mathbb{A}, \equiv_{\mathcal{K}})$ is NP-hard.

- In which cases can we prove that $\text{CSP}(\mathbb{A})$ is NP-hard?
- What can we say otherwise?

Similar questions are interesting for "hardnesses" other than NP-hardness:

- (un)bounded width,
- (un)bounded linear width,
- (in)solvability in particular logics



Hardness of Monochromatic triangle

Problem (Monochromatic triangle). Following decision problem:

Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?

Problem (Monochromatic triangle). Following decision problem:

Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?

Garey & Johnson:

[GT6] **MONOCHROMATIC TRIANGLE**

INSTANCE: Graph $G = (V, E)$.

QUESTION: Is there a partition of E into two disjoint sets E_1, E_2 such that neither $G_1 = (V, E_1)$ nor $G_2 = (V, E_2)$ contains a triangle?

Reference: [Burr, 1976]. Transformation from 3SAT.

Problem (Monochromatic triangle). Following decision problem:

Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?

Garey & Johnson:

[GT6] **MONOCHROMATIC TRIANGLE**

INSTANCE: Graph $G = (V, E)$.

QUESTION: Is there a partition of E into two disjoint sets E_1, E_2 such that neither $G_1 = (V, E_1)$ nor $G_2 = (V, E_2)$ contains a triangle?

Reference: [Burr, 1976]. Transformation from 3SAT.


BURR, S. [1976], private communication.

Hardness of Monochromatic triangle

Problem (Monochromatic triangle). Following decision problem:

Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?




Hardness of Monochromatic triangle

Problem (Monochromatic triangle). Following decision problem:

Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?




Hardness of Monochromatic triangle

Problem (Monochromatic triangle). Following decision problem:


Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?

First attempt:



\rightsquigarrow



Instance of Not-All-Equal

Instance of monochromatic triangle


Hardness of Monochromatic triangle

Problem (Monochromatic triangle). Following decision problem:


Input A finite undirected graph $G = (V, E)$

Question Does there exist a coloring $\chi: E \rightarrow \{R, B\}$ such that $\chi^{-1}(c)$ is triangle-free for every $c \in \{R, B\}$?

First attempt:



~~~



Where to put the edge for  $x_8$ ?

Instance of Not-All-Equal

Instance of monochromatic triangle


# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:

**Input** A finite undirected graph  $G = (V, E)$


**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?

First attempt:



Instance of Not-All-Equal

$\rightsquigarrow$



Where to put the edge for  $x_8$ ?

Instance of monochromatic triangle

The difference between NAE and Monochromatic Triangle:

- In NAE: unrestricted use of ternary constraints
- In Monochromatic triangle: ternary constraints (on edges) can only be imposed on triangles, and are **automatically** imposed on triangles


# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:

**Input** A finite undirected graph  $G = (V, E)$

**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?

Second attempt:



Instance of Not-All-Equal

The difference between NAE and Monochromatic Triangle:

- In NAE: unrestricted use of ternary constraints
- In Monochromatic triangle: ternary constraints (on edges) can only be imposed on triangles, and are **automatically** imposed on triangles



# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:

**Input** A finite undirected graph  $G = (V, E)$

**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?

Second attempt:



Instance of Not-All-Equal

The difference between NAE and Monochromatic Triangle:

- In NAE: unrestricted use of ternary constraints
- In Monochromatic triangle: ternary constraints (on edges) can only be imposed on triangles, and are **automatically** imposed on triangles



# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:


**Input** A finite undirected graph  $G = (V, E)$

**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?

Second attempt:



Instance of Not-All-Equal





# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:

**Input** A finite undirected graph  $G = (V, E)$

**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?


Second attempt:



Instance of Not-All-Equal

Intuition:

- In the finite template, equality comes for free
- equality allows one to use "structural" constraints as "normal" constraints





# Hardness of Monochromatic triangle

**Problem** (Monochromatic triangle). Following decision problem:

**Input** A finite undirected graph  $G = (V, E)$


**Question** Does there exist a coloring  $\chi: E \rightarrow \{R, B\}$  such that  $\chi^{-1}(c)$  is triangle-free for every  $c \in \{R, B\}$ ?

Second attempt:



Instance of Not-All-Equal

**Proposition** (Barsukov, M., Perinti). Monochromatic Triangle remains hard on  $K_4$ -free graphs  
(The gadget is beautiful but too large to fit on this slide)




## Why equality is a problem

**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.


## Why equality is a problem

**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.




## Why equality is a problem

**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.



## Why equality is a problem


**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.



## Why equality is a problem

**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.


This is Betweenness



## Why equality is a problem

**Observation.**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is the class of linear orders.


This is Betweenness



**Proposition (Pinckner).**  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard whenever  $\mathcal{K}$  is a class of linearly ordered structures.


## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.




## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard




## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$




## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$




# The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint




## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation




## The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation




# The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation
- No more paths of even length between the potatoes




# The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation
- No more paths of even length between the potatoes




# The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation
- No more paths of even length between the potatoes



# The Loop Lemma(s) of Smooth Approximations

- $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard if there exist pp-definable  $O, Z \subseteq \mathbb{A}_{fin}$  and  $N \subseteq (O \cup Z)^3$  representing NAE.
- If we can define those relations in  $\mathbb{A}$ , then  $\text{CSP}(\mathbb{A})$  is NP-hard
- Consider the relation generated by a single undirected edge  $\{a, b\}$  under  $\text{Pol}(\mathbb{A})$
- If there exists a path of even length between  $Z$  and  $O$ , reduce to the neighborhood of midpoint
- Finite CSP must be hard on those new subsets  
~~ existence of a new “NAE” like relation
- No more paths of even length between the potatoes
- This defines (in  $\mathbb{A}$ ) an equivalence relation that refines the original partition



# The Loop Lemma(s) of Smooth Approximations

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$

# The Loop Lemma(s) of Smooth Approximations

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$

**Theorem** (M., Nagy, Pinsker, Wrona). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is **not** NP-hard.

For every **minimal subfactor** of  $\text{Pol}(\mathbb{A}_{fin})$ , one of the following holds:

- $\exists$  pp-definable  $\sim$  that approximates the subfactor
- Every pp-definable cyclic relation on the subfactor contains a pseudoloop.

# The Fundamental Theorem of Smooth Approximations

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$

**Theorem** (M., Pinsker). Suppose

- $\{Z, O\}$  has a **smooth approximation**  $\sim$  definable in  $\mathbb{A}$
- $\text{Pol}(\mathbb{A})$  **locally interpolates**  $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$ .

Then  $\exists$  uniformly continuous minion homomorphism  $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{K}_3)$ .

In particular **CSP( $\mathbb{A}$ ) is NP-hard**.

# The Fundamental Theorem of Smooth Approximations

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$

**Theorem** (M., Pinsker). Suppose

- $\{Z, O\}$  has a **smooth approximation**  $\sim$  definable in  $\mathbb{A}$
- $\text{Pol}(\mathbb{A})$  **locally interpolates**  $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$ .

Then  $\exists$  uniformly continuous minion homomorphism  $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{K}_3)$ .

In particular **CSP( $\mathbb{A}$ ) is NP-hard**.

- **Smooth** means that  $\sim$  is not finer than  $\equiv_{\mathcal{K}}$ , at least on some interesting subset
- Need some assumptions on  $\text{Aut}(\mathbb{A})$  to go from "normal" approximation to "smooth"  
**No algebraicity and primitivity**

# The Fundamental Theorem of Smooth Approximations

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$

**Theorem** (M., Pinsker). Suppose

- $\{Z, O\}$  has a **smooth approximation**  $\sim$  definable in  $\mathbb{A}$
- $\text{Pol}(\mathbb{A})$  **locally interpolates**  $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$ .

Then  $\exists$  uniformly continuous minion homomorphism  $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{K}_3)$ .

In particular **CSP( $\mathbb{A}$ ) is NP-hard**.

- **Smooth** means that  $\sim$  is not finer than  $\equiv_{\mathcal{K}}$ , at least on some interesting subset
- Need some assumptions on  $\text{Aut}(\mathbb{A})$  to go from "normal" approximation to "smooth"  
**No algebraicity and primitivity**
- "local interpolation" means every  $f \in \text{Pol}(\mathbb{A})$  "looks like" some  $\tilde{f} \in \text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}})$  on every finite set

## Consequence of loops

**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$



Implies  $\exists f \in \text{Pol}(\mathbb{A})$  such that if  $f(a, b), f(b, a) \in Z \cup O$

$$f(a, b) \in Z \Leftrightarrow f(b, a) \in Z$$


for all  $a, b$  disjoint

## Consequence of loops


**Theorem** (M., Pinsker). Suppose  $\text{CSP}(\mathbb{A}_{fin})$  is NP-hard.

There exists an NP-hardness "certificate"  $\{Z, O\}$  such that one of the following holds:

- $\exists$  pp-definable  $\sim$  that refines  $\{Z, O\}$   
 $\sim$  is an **approximation** of  $\{Z, O\}$ .
- Every pp-definable binary relation with an edge between  $Z$  and  $O$  contains a pair  $(a, \alpha(a))$  for some  $\alpha \in \text{Aut}(\mathbb{A})$



## Connections with Promise CSPs




**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .  
There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

# Infinite CSPs as (infinite, finite)-PCSPs

**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .  
 There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod } \text{Aut}(\mathbb{B}^*)$$



# Infinite CSPs as (infinite, finite)-PCSPs


**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .

There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod Aut}(\mathbb{B}^*)$$

**Corollary** (Barto, Bulín, Krokhin, Opršal). If  $\text{Pol}(\mathbb{B})$  does not contain pseudo-Olšák operations, then approximate hypergraph coloring reduces to  $\text{CSP}(\mathbb{B})$ .

$$e_1 f(x, x, y, y, y, x) \approx e_2 f(x, y, x, y, x, y) \approx e_3 f(y, x, x, x, y, y)$$



# Infinite CSPs as (infinite, finite)-PCSPs


**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .

There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod Aut}(\mathbb{B}^*)$$

**Corollary** (Atserias, Dalmau). If  $\text{Pol}(\mathbb{B})$  does not contain a pseudo-WNU of some arity  $k \geq 3$ , then  $\mathbb{B}$  does not have bounded width.

$$e_1 w(x, \dots, x, x, y) \approx e_2 w(x, \dots, x, y, x) \approx \dots \approx e_k w(y, x, \dots, x)$$



# Infinite CSPs as (infinite, finite)-PCSPs


**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .

There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod Aut}(\mathbb{B}^*)$$

**Corollary** (Barto, Bulín, Krokhin, Opršal). If  $\text{Pol}(\mathbb{B})$  does not contain a pseudo-Siggers operation, then  $\text{PCSP}(\mathbb{K}_3, \mathbb{K}_r)$  reduces to  $\text{CSP}(\mathbb{B})$ .

$$u \circ s(x, y, x, z, y, z) \approx v \circ s(y, x, z, x, z, y)$$



# Infinite CSPs as (infinite, finite)-PCSPs

**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .


There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .

$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod Aut}(\mathbb{B}^*)$$

**Theorem (M.).** Let  $\mathbb{B}$  be a maximally tractable temporal CSP,  
 $\mathbb{A}$  as above.

There exists  $\mathbb{A}' \subseteq \mathbb{A}$  such that  $\text{PCSP}(\mathbb{A}', \mathbb{B}_{fin})$  is not finitely  
tractable:

$$\mathbb{A}' \rightarrow \mathbb{C} \rightarrow \mathbb{B}_{fin} \text{ and } \mathbb{C} \text{ finite} \implies \text{CSP}(\mathbb{C}) \text{ is NP-hard}$$



# Infinite CSPs as (infinite, finite)-PCSPs

**Theorem (M.).** Let  $\mathbb{B}$  be a reduct of a finitely bounded homogeneous structure  $\mathbb{B}^*$ .

There exists  $\mathbb{A}$  (infinite) s.t.  $\text{CSP}(\mathbb{B})$  is ptime equivalent to  $\text{PCSP}(\mathbb{A}, \mathbb{B}_{fin})$ .


$$(\forall \mathbb{A}' \subset \mathbb{A} \text{ finite} : \text{Pol}(\mathbb{A}', \mathbb{B}_{fin}) \models \Sigma) \iff \text{Pol}(\mathbb{B}) \models \Sigma \text{ mod Aut}(\mathbb{B}^*)$$

**Theorem (M.).** Let  $\mathbb{B}$  be a maximally tractable temporal CSP,  
 $\mathbb{A}$  as above.

There exists  $\mathbb{A}' \subseteq \mathbb{A}$  such that  $\text{PCSP}(\mathbb{A}', \mathbb{B}_{fin})$  is not finitely  
tractable:

$$\mathbb{A}' \rightarrow \mathbb{C} \rightarrow \mathbb{B}_{fin} \text{ and } \mathbb{C} \text{ finite} \implies \text{CSP}(\mathbb{C}) \text{ is NP-hard}$$

**Question.** Do these finite and tractable PCSP templates need new algorithms?



Finite minions!

## Temporal CSPs revisited

Temporal CSPs / ordering CSPs:

- certificate class  $\mathcal{K}$ : linear orders
- $\text{CSP}(\mathbb{Q}; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable using  $<$


**Theorem** (Bodirsky, Kára). For every temporal  $\text{CSP}(\mathbb{B})$ , if  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(\mathbb{K}_3)$ , then  $\text{CSP}(\mathbb{B})$  is in P.

# Temporal CSPs revisited

Temporal CSPs / ordering CSPs:

- certificate class  $\mathcal{K}$ : linear orders
- $\text{CSP}(\mathbb{Q}; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable using  $<$

**Theorem** (Bodirsky, Kára). For every temporal  $\text{CSP}(\mathbb{B})$ , if  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(\mathbb{K}_3)$ , then  $\text{CSP}(\mathbb{B})$  is in P.



# Temporal CSPs revisited


Temporal CSPs / ordering CSPs:

- certificate class  $\mathcal{K}$ : linear orders
- $\text{CSP}(\mathbb{Q}; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable using  $<$

**Theorem** (Bodirsky, Kára). For every temporal  $\text{CSP}(\mathbb{B})$ , if  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(\mathbb{K}_3)$ , then  $\text{CSP}(\mathbb{B})$  is in P.

Non-satisfactory parts of the proof:

- 4 ad-hoc algorithms, yet with a "finite flavor"
- Algorithms need polymorphisms to work
- Any work building on this must essentially follow the same case distinction
- Hard to use as a blackbox



- Original idea: iteratively decompose instance by finding a non-empty set of variables than can be minimal in a solution: **free set** of variables (Bodirsky, Kára)
- Deciding the existence of such a free set is a boolean CSP with domain {min, not min} (M., Pinsker)
- Tractable boolean CSPs are solvable by BLP+AIP (Brakensiek, Guruswami, Wrochna, Živný)

- Original idea: iteratively decompose instance by finding a non-empty set of variables than can be minimal in a solution: **free set** of variables (Bodirsky, Kára)
- Deciding the existence of such a free set is a boolean CSP with domain  $\{\min, \text{not min}\}$  (M., Pinsker)
- Tractable boolean CSPs are solvable by BLP+AIP (Brakensiek, Guruswami, Wrochna, Živný)



**Observation.** On input with  $n$  variables,  $\text{CSP}(\mathbb{B})$  reduces (trivially) to  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  where  $\mathbb{A}$  has domain  $\{1, \dots, n\}$ .

- Original idea: iteratively decompose instance by finding a non-empty set of variables than can be minimal in a solution: **free set** of variables (Bodirsky, Kára)
- Deciding the existence of such a free set is a boolean CSP with domain  $\{\min, \text{not min}\}$  (M., Pinsker)
- Tractable boolean CSPs are solvable by BLP+AIP (Brakensiek, Guruswami, Wrochna, Živný)

 **Observation.** On input with  $n$  variables,  $\text{CSP}(\mathbb{B})$  reduces (trivially) to  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  where  $\mathbb{A}$  has domain  $\{1, \dots, n\}$ .

**Theorem** (M.). Let  $\mathbb{B}$  be a temporal structure with tractable CSP,  $\mathbb{A} \subset \mathbb{B}$  finite.

$k \geq 1$  large enough (depending on arity of relations).

Let  $\mathbb{X}$  be an instance that has a ( $k$ th level) singleton BLP/AIP solution w.r.t.  $\mathbb{A}$ .

Then  $\mathbb{X}$  has a non-empty free set  $F$  s.t.  $\mathbb{X} \wedge "F \text{ is minimal}"$  still has a ( $k$ th level) singleton BLP/AIP solution.


**Theorem (M.).**  $\mathbb{B}$  temporal structure such that  $\nexists \text{Pol}(\mathbb{B}, \mathbf{c}) \rightarrow \text{Pol}(K_3)$ ,  $\mathbb{A}$  finite structure.  
Then  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  is solvable by some level of singleton BLP+AIP.

**Theorem (M.).**  $\mathbb{B}$  temporal structure such that  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ ,  $\mathbb{A}$  finite structure.  
Then PCSP( $\mathbb{A}, \mathbb{B}$ ) is solvable by some level of singleton BLP+AIP.

Singleton Arc Consistency

Singleton AIP

Width  $\leq r + 1$




**Theorem (M.).**  $\mathbb{B}$  temporal structure such that  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ ,  $\mathbb{A}$  finite structure.  
Then  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  is solvable by some level of singleton BLP+AIP.

Singleton Arc Consistency

Singleton AIP

Width  $\leq r + 1$




**Theorem (Bodirsky, Kára).**  $\exists$  temporal CSP templates with those polymorphisms and without bounded width.

**Theorem (M.).**  $\mathbb{B}$  temporal structure such that  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ ,  $\mathbb{A}$  finite structure.  
 Then  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  is solvable by some level of singleton BLP+AIP.

Singleton Arc Consistency

Singleton AIP

Width  $\leq r + 1$



**Theorem (Bodirsky, Kára).**  $\exists$  temporal CSP templates with those polymorphisms and without bounded width.

**Corollary (M.).** There exist finite PCSP templates  $(\mathbb{A}, \mathbb{B})$  such that:


- $\text{PCSP}(\mathbb{A}, \mathbb{B})$  has bounded width,
- $\text{PCSP}(\mathbb{A}, \mathbb{B})$  not solvable by BLP+AIP,
- Not finitely tractable: every finite  $\mathbb{C}$  such that  $\mathbb{A} \rightarrow \mathbb{C} \rightarrow \mathbb{B}$  has an NP-hard CSP. In particular, no finite  $\mathbb{C}$  with bounded width.

**Theorem (M.).**  $\mathbb{B}$  temporal structure such that  $\nexists \text{Pol}(\mathbb{B}, c) \rightarrow \text{Pol}(K_3)$ ,  $\mathbb{A}$  finite structure.  
 Then  $\text{PCSP}(\mathbb{A}, \mathbb{B})$  is solvable by some level of singleton BLP+AIP.

Singleton Arc Consistency

Singleton AIP

Width  $\leq r + 1$



**Theorem (Bodirsky, Kára).**  $\exists$  temporal CSP templates with those polymorphisms and without bounded width.

**Corollary (M.).** There exist finite PCSP templates  $(\mathbb{A}, \mathbb{B})$  such that:

- $\text{PCSP}(\mathbb{A}, \mathbb{B})$  solvable by singleton AIP,
- $\text{PCSP}(\mathbb{A}, \mathbb{B})$  not FPC-solvable,
- Not finitely tractable: every finite  $\mathbb{C}$  such that  $\mathbb{A} \rightarrow \mathbb{C} \rightarrow \mathbb{B}$  has an NP-hard CSP. In particular, no finite  $\mathbb{C}$  with bounded width.


## CSPs over Hypergraphs:

- Certificate class  $\mathcal{K}$ : all hypergraphs, or all  $K_n^r$ -free hypergraphs ( $n > r$ )
- $\text{CSP}(V; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable over certain homogeneous hypergraphs

# CSPs over Hypergraphs revisited






## CSPs over Hypergraphs:

- Certificate class  $\mathcal{K}$ : all hypergraphs, or all  $K_n^r$ -free hypergraphs ( $n > r$ )
- $\text{CSP}(V; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable over certain homogeneous hypergraphs







$\iff$

$$\begin{array}{ll} \bullet = \bullet & \bullet \neq \bullet \\ \# = \# & \parallel \parallel \\ \bullet = \bullet & \bullet \neq \bullet \end{array}$$



instance  
with 4  
constraints



CSPs over Hypergraphs:

- Certificate class  $\mathcal{K}$ : all hypergraphs, or all  $K_n^r$ -free hypergraphs ( $n > r$ )
- $\text{CSP}(V; R_1, \dots, R_k)$  where  $R_1, \dots, R_k$  definable over certain homogeneous hypergraphs

**Theorem** (M., Nagy, Pinsker). The conjecture is true for such CSPs.

Our proof has interesting features:

- It is not true that "good polymorphisms" preserve  $\equiv_{\mathcal{K}}$  (like for ordered structures)
- It is not true that "good polymorphisms" are completely wild (unlike for ordered structures)
- Algorithm loosely based on Zhuk's approach (binary absorption into affine algebra)


# CSPs over Hypergraphs revisited

**Theorem** (M., Nagy). BLP+AIP solves all the tractable cases.

**Theorem** (M., Nagy). BLP+AIP solves all the tractable cases.

- Label cover instance with **ordered** hypergraphs


all 27 types of **ordered**  
3-hypergraphs on  $\leq 3$   
vertices



**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.

- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance


all 27 types of **ordered** 3-hypergraphs on  $\leq 3$  vertices



**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.


- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance
- $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}_<})$  has injective operation

all 27 types of **ordered** 3-hypergraphs on  $\leq 3$  vertices




**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.

- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance
- $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}_<})$  has injective operation




**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.

- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance
- $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}_<})$  has injective operation
- Get a BLP+AIP distribution for label cover instance over **unordered** hypergraphs



**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.


- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance
- $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}_<})$  has injective operation
- Get a BLP+AIP distribution for label cover instance over **unordered** hypergraphs
- This is a Boolean CSP with non-trivial polymorphisms, so there is a solution



# CSPs over Hypergraphs revisited

**Theorem (M., Nagy).** BLP+AIP solves all the tractable cases.

- Label cover instance with **ordered** hypergraphs
- Find BLP+AIP solution for that instance
- $\text{Pol}(\mathbb{A}, \equiv_{\mathcal{K}_<})$  has injective operation
- Get a BLP+AIP distribution for label cover instance over **unordered** hypergraphs
- This is a Boolean CSP with non-trivial polymorphisms, so there is a solution



- Decision: basically done with the "Schaefer" case  
Next: progress on uniform algorithm for finite CSPs could help solving the infinite conjecture as well
- Optimisation: valued CSPs classified for order CSPs (Bodirsky, Bonnet, Semanišinová)
- Approximation resistance:
  - known for order CSPs (Guruswami, Håstad, Manokaran, Raghavendra, Charikar)
  - phylogenetic CSPs (Chatziafratis, Makarychev)
- Parametrized versions (Jonsson, Pilipczuk, Osipov, Wahlström, ...)