

MMSNP: An algebraic proof of the dichotomy

Manuel Bodirsky, Florent Madelaine, **Antoine Mottet**


September 25, 2018

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity:

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:




- ▶ Complexity: NP-complete.

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

-
- ▶ Input: undirected graph G
 - ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: in P.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg($$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x})) \quad)$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

\rightsquigarrow MMSNP is the logic of FPP.

Dichotomy: “every problem is easy or hard.”

Dichotomy: “every problem is easy or hard.”

Theorem (Bulatov, Zhuk '17)

Every finite-domain CSP is in P or NP -complete.

Dichotomy: “every problem is easy or hard.”

Theorem (Bulatov, Zhuk '17)

Every finite-domain CSP is in P or NP-complete.

Theorem (Feder-Vardi '93, Kun '13)

MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.

Dichotomy: “every problem is easy or hard.”

Theorem (Bulatov, Zhuk '17)

Every finite-domain CSP is in P or NP-complete.

Theorem (Feder-Vardi '93, Kun '13)

MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.

The End.

Dichotomy: “every problem is easy or hard.”

Theorem (Bulatov, Zhuk '17)

Every finite-domain CSP is in P or NP-complete.

Theorem (Feder-Vardi '93, Kun '13)

MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.

The End?

- ▶ Finite-domain dichotomy is **universal**-algebraic.
Feder-Vardi-Kun's reduction is not.

- ▶ Finite-domain dichotomy is **universal**-algebraic.
Feder-Vardi-Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs.
The conjecture for infinite domains is still open.

- ▶ Finite-domain dichotomy is **universal**-algebraic. Feder-Vardi-Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs. The conjecture for infinite domains is still open.

Theorem

Let \mathcal{A} be such that $\text{CSP}(\mathcal{A})$ is in MMSNP.

Exactly one of the following holds:

- ▶ *There is a uniformly continuous homomorphism $\text{Pol}(\mathcal{A}) \rightarrow \text{Pol}(\text{SAT})$, and $\text{CSP}(\mathcal{A})$ is **NP-complete**,*
- ▶ *$\text{Pol}(\mathcal{A})$ contains a cyclic operation and $\text{CSP}(\mathcal{A})$ is in **P**.*

- ▶ Finite-domain dichotomy is **universal**-algebraic. Feder-Vardi-Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs. The conjecture for infinite domains is still open.

Theorem

Let \mathcal{A} be such that $\text{CSP}(\mathcal{A})$ is in MMSNP.

Exactly one of the following holds:

- ▶ *There is a uniformly continuous homomorphism $\text{Pol}(\mathcal{A}) \rightarrow \text{Pol}(\text{SAT})$, and $\text{CSP}(\mathcal{A})$ is **NP-complete**,*
- ▶ *$\text{Pol}(\mathcal{A})$ contains a cyclic operation and $\text{CSP}(\mathcal{A})$ is in **P**.*

\rightsquigarrow new proof of the Feder-Vardi-Kun theorem,

- ▶ Finite-domain dichotomy is **universal**-algebraic. Feder-Vardi-Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs. The conjecture for infinite domains is still open.

Theorem

Let \mathcal{A} be such that $\text{CSP}(\mathcal{A})$ is in MMSNP.

Exactly one of the following holds:

- ▶ *There is a uniformly continuous homomorphism $\text{Pol}(\mathcal{A}) \rightarrow \text{Pol}(\text{SAT})$, and $\text{CSP}(\mathcal{A})$ is **NP-complete**,*
- ▶ *$\text{Pol}(\mathcal{A})$ contains a cyclic operation and $\text{CSP}(\mathcal{A})$ is in **P**.*

\rightsquigarrow new proof of the Feder-Vardi-Kun theorem,

\rightsquigarrow the infinite-domain conjecture for CSPs in MMSNP is true.

Definition

$\mathcal{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathcal{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathcal{A} ,
- ▶ **Question:** is there a homomorphism $\mathcal{A} \rightarrow \mathcal{B}$?

Definition

$\mathcal{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathcal{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathcal{A} ,
- ▶ **Question:** is there a homomorphism $\mathcal{A} \rightarrow \mathcal{B}$?

Examples:

- ▶ $\text{CSP}(K_3)$: 3-colouring problem,
- ▶ $\text{CSP}(\mathbb{Q}, <)$: digraph acyclicity,
- ▶ In general, we don't restrict to graphs.
 $\text{CSP}(\mathbb{Z}; +, \times, 1)$: satisfiability of diophantine equations.

Definition

$\mathcal{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathcal{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathcal{A} ,
- ▶ **Question:** is there a homomorphism $\mathcal{A} \rightarrow \mathcal{B}$?

Examples:

- ▶ $\text{CSP}(K_3)$: 3-colouring problem,
- ▶ $\text{CSP}(\mathbb{Q}, <)$: digraph acyclicity,
- ▶ In general, we don't restrict to graphs.
 $\text{CSP}(\mathbb{Z}; +, \times, 1)$: satisfiability of diophantine equations.

If \mathcal{B} is finite, $\text{CSP}(\mathcal{B})$ is in P or NP-complete.

- For a nice structure \mathcal{B} ,

complexity of $\text{CSP}(\mathcal{B}) \leftrightarrow$ algebraic properties of $\text{Pol}(\mathcal{B})$

- ▶ For a nice structure \mathcal{B} ,

complexity of $\text{CSP}(\mathcal{B}) \leftrightarrow$ algebraic properties of $\text{Pol}(\mathcal{B})$

- ▶ One measures how “trivial” $\text{Pol}(\mathcal{B})$ is by the equations satisfiable in it.

- ▶ For a nice structure \mathcal{B} ,

complexity of $\text{CSP}(\mathcal{B}) \leftrightarrow$ algebraic properties of $\text{Pol}(\mathcal{B})$

- ▶ One measures how “trivial” $\text{Pol}(\mathcal{B})$ is by the equations satisfiable in it.
- ▶ Trivial-est of all: $\text{Pol}(\text{SAT})$, which only contains projections.

- ▶ For a nice structure \mathcal{B} ,

complexity of $\text{CSP}(\mathcal{B}) \leftrightarrow$ algebraic properties of $\text{Pol}(\mathcal{B})$

- ▶ One measures how “trivial” $\text{Pol}(\mathcal{B})$ is by the equations satisfiable in it.
- ▶ Trivial-est of all: $\text{Pol}(\text{SAT})$, which only contains projections.
- ▶ A **homomorphism** $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\mathcal{A})$ induces a log-space reduction $\text{CSP}(\mathcal{A}) \leq \text{CSP}(\mathcal{B})$.

- ▶ For a nice structure \mathcal{B} ,

complexity of $\text{CSP}(\mathcal{B}) \leftrightarrow$ algebraic properties of $\text{Pol}(\mathcal{B})$

- ▶ One measures how “trivial” $\text{Pol}(\mathcal{B})$ is by the equations satisfiable in it.
- ▶ Trivial-est of all: $\text{Pol}(\text{SAT})$, which only contains projections.
- ▶ A **homomorphism** $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\mathcal{A})$ induces a log-space reduction $\text{CSP}(\mathcal{A}) \leq \text{CSP}(\mathcal{B})$.
- ▶ **Algebraic hardness:** $\text{Pol}(\mathcal{A}) \rightarrow \text{Pol}(\text{SAT})$ implies that $\text{CSP}(\mathcal{A})$ is NP-hard.

Theorem (Cherlin-Shelah-Shi, '99)

*For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.*

Theorem (Cherlin-Shelah-Shi, '99)


*For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.*

Obstructions \mathcal{F} : 

Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable *universal \mathcal{F} -free* graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.


Obstructions \mathcal{F} : 



Theorem (Cherlin-Shelah-Shi, '99)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists a countable **universal \mathcal{F} -free** graph.

Obstructions \mathcal{F} : 



"Algebraic hardness is the only source of hardness":

Conjecture (Bodirsky-Pinsker, '11 (rephrased))

*Let \mathcal{B} be a reduct of a finitely bounded homogeneous structure.
If there is no uniformly continuous homomorphism
 $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathcal{B})$ is in P .*

"Algebraic hardness is the only source of hardness":

Conjecture (Bodirsky-Pinsker, '11 (rephrased))

*Let \mathcal{B} be a reduct of a finitely bounded homogeneous structure.
If there is no uniformly continuous homomorphism
 $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathcal{B})$ is in P.*

Theorem

The conjecture is true for \mathcal{B} such that $\text{CSP}(\mathcal{B})$ is in MMSNP.

Definition


$f : B^k \rightarrow B$. f is **canonical** if

“the colour of the output only depends on the colours of the inputs”.


Definition

$f : B^k \rightarrow B$. f is **canonical** if


“the colour of the output only depends on the colours of the inputs”.




Every canonical function determines a **behaviour** on colours.



Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.


Only **finitely many** behaviours of a given arity.

Theorem (Bodirsky-M, '16)

Let \mathcal{B} be in the class (reduct of...).

If there is no homomorphism $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathcal{B})$ is in P .

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.
Only **finitely many** behaviours of a given arity.

Theorem (Bodirsky-M, '16)

Let \mathcal{B} be in the class (reduct of...).

If there is no homomorphism $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathcal{B})$ is in P .

\leadsto the BP conjecture is true when $\text{Pol}(\mathcal{B})_{\text{can}} = \text{Pol}(\mathcal{B})$.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.
- ▶ Challenge: extending $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$ to $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.
 - ▶ Challenge: extending $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$ to $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$.
1. We suppose that there is a clone homomorphism $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.
 - ▶ Challenge: extending $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$ to $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$.
1. We suppose that there is a clone homomorphism $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$.
 2. Prove that $\text{CSP}(\mathcal{B})$ is algebraically hard iff $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ is.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.
 - ▶ Challenge: extending $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$ to $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$.
1. We suppose that there is a clone homomorphism $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$.
 2. Prove that $\text{CSP}(\mathcal{B})$ is algebraically hard iff $\text{CSP}(\mathcal{B}, \bullet, \circ)$ is.
 3. $\text{Pol}(\mathcal{B}, \bullet, \circ)_{\text{can}}$ is an **idempotent** clone.
 \leadsto **subfactor** of $\text{Pol}(\mathcal{B}, \bullet, \circ)_{\text{can}}$ isomorphic to $\text{Pol}(\text{SAT})$.

- ▶ $\text{Pol}(\mathcal{B})_{\text{can}}$ is not sufficient in general.
 - ▶ Challenge: extending $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$ to $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$.
1. We suppose that there is a clone homomorphism $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$.
 2. Prove that $\text{CSP}(\mathcal{B})$ is algebraically hard iff $\text{CSP}(\mathcal{B}, \bullet, \circ)$ is.
 3. $\text{Pol}(\mathcal{B}, \bullet, \circ)_{\text{can}}$ is an **idempotent** clone.
 \leadsto **subfactor** of $\text{Pol}(\mathcal{B}, \bullet, \circ)_{\text{can}}$ isomorphic to $\text{Pol}(\text{SAT})$.
 4. Define a homomorphism $\text{Pol}(\mathcal{B}, \bullet, \circ) \rightarrow \text{Pol}(\text{SAT})$ by **canonizing** and composing with ξ .

Summing up:

Theorem

Let \mathcal{B} be an MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$,*

and $\text{CSP}(\mathcal{B})$ is in P , or $\text{CSP}(\mathcal{B})$ is NP-complete.

Summing up:

Theorem

Let \mathcal{B} be an MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$,*
- 2. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$,*

and $\text{CSP}(\mathcal{B})$ is in P , or $\text{CSP}(\mathcal{B})$ is NP-complete.

Summing up:

Theorem

Let \mathcal{B} be an MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$,*
- 2. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$,*

and $\text{CSP}(\mathcal{B})$ is in P , or $\text{CSP}(\mathcal{B})$ is NP-complete.

Item 2. can be checked effectively given an MMSNP sentence.

Summing up:

Theorem

Let \mathcal{B} be an MMSNP structure.

Then either the following equivalent statements hold:

- 1. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$,*
- 2. there is no uniformly continuous homomorphism $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \text{Pol}(\text{SAT})$,*

and $\text{CSP}(\mathcal{B})$ is in P , or $\text{CSP}(\mathcal{B})$ is NP-complete.

Item 2. can be checked effectively given an MMSNP sentence.

The End.

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, '15)


\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?


A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, '15)


\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?


Rephrased: do $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ and $\text{CSP}(\mathcal{B})$ have same complexity?





- ▶ Obstructions \mathcal{F} : 
- ▶ x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:




- ▶ Obstructions \mathcal{F} : 
- ▶ x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:





- Obstructions \mathcal{F} : 
- x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:







- x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:








- x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:



- Obstructions \mathcal{F} : 
- x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \bullet)$:



- Obstructions \mathcal{F} : 
- x precoloured in the instance of $\text{CSP}(\mathcal{B}, \bullet, \circ)$:



- The input precoloured graph is colourable iff the graph obtained by adding the gadgets is colourable.

- ▶ σ : set of colour symbols.
- ▶ A **trivial subfactor** of \mathcal{C} is a partition $S \uplus T \subseteq \sigma$ such that \mathcal{C}/\sim is isomorphic to $\text{Pol}(\text{SAT})$.

Proposition

S, T trivial subfactor of \mathcal{C} . $\exists(\sigma, E)$ bipartite undirected graph s.t.:

- ▶ (σ, E) contains an edge from S to T ;
- ▶ (σ, E) has no path of even length from S to T ;
- ▶ E is preserved by $\text{Pol}(\mathcal{B})$.

Theorem (Hubička-Nešetřil, 2016)

Let \mathcal{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathcal{B}, <)$ is ω -categorical and Ramsey.

Theorem (Hubička-Nešetřil, 2016)

Let \mathcal{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathcal{B}, <)$ is ω -categorical and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Theorem (Hubička-Nešetřil, 2016)

Let \mathcal{B} be an MMSNP structure. Then there is a linear order $<$ on B such that $(\mathcal{B}, <)$ is ω -categorical and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Given $f \in \text{Pol}(\mathcal{B})$, and $\xi: \mathcal{C} \rightarrow \text{Pol}(\text{SAT})$ given by subfactor $\{S, T\}$, define $\phi(f) := \xi(g)$ where g is an arbitrary function in $\mathcal{C} \cap \overline{\text{Aut}(\mathcal{B}, <)f\text{Aut}(\mathcal{B}, <)}$.

Proposition

$\phi: \text{Pol}(\mathcal{B}) \rightarrow \text{Pol}(\text{SAT})$ is a well-defined uniformly continuous homomorphism.