


Theory of Constraint Satisfaction Problems

Antoine Wiehe

Colloquium Computational Discrete Mathematics
TU Graz



Instance:

- a **domain** A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

Instance:

- a **domain** A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

Examples. Boolean Satisfiability

Is $(x \vee y \vee z) \wedge (\neg z \vee t \vee p) \wedge \dots$ satisfiable?

Instance:

- a **domain** A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

Examples. Boolean Satisfiability

Is $(x \vee y \vee z) \wedge (\neg z \vee t \vee p) \wedge \dots$ satisfiable?

Solving Linear Equations

$$\text{Is } \begin{cases} 2x & -y & & = & 0 \\ & y & +z & = & 1 \\ x & & -z & = & 2 \end{cases} \text{ satisfiable?}$$

Instance:

- a **domain** A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

Examples. Boolean Satisfiability

Is $(x \vee y \vee z) \wedge (\neg z \vee t \vee p) \wedge \dots$ satisfiable?

NP-complete

Solving Linear Equations

$$\text{Is } \begin{cases} 2x & -y & & = & 0 \\ & y & +z & = & 1 \\ x & & -z & = & 2 \end{cases} \text{ satisfiable?}$$

Solvable in polynomial time

Instance:

- a domain A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

Examples. Boolean Satisfiability

Is $(x \vee y \vee z) \wedge (\neg z \vee t \vee p) \wedge \dots$ satisfiable?

NP-complete


Solving Linear Equations

Is $\begin{cases} 2x & -y & & = & 0 \\ & y & +z & = & 1 \\ x & & -z & = & 2 \end{cases}$ satisfiable?

Solvable in polynomial time

Linear Programming

Is $\mathbf{Ax} = \mathbf{b}$ satisfiable in $\mathbb{R}_{\geq 0}$?



Instance:

- a **domain** A
- variables x, y, z, \dots
- constraints: $x \vee y \vee z, z \leq x, 2x + y = 0, \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

- **Non-uniform** CSP: domain is **fixed** and only certain types of constraints are allowed

Constraint with r variables \leftrightarrow relation $R \subseteq A^r$ listing all valid assignments

Constraint language / **template**: $\mathbb{A} = (A; R_1, \dots, R_k)$

Instance:

- a ~~domain~~ A
- variables x, y, z, \dots
- constraints: $R_1(x, y, z), R_2(z, x), R_3(x, y), R_2(z, y), \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

- **Non-uniform** CSP: domain is **fixed** and only certain types of constraints are allowed

Constraint with r variables \leftrightarrow relation $R \subseteq A^r$ listing all valid assignments

Constraint language / **template**: $\mathbb{A} = (A; R_1, \dots, R_k)$

Instance:

- a ~~domain~~ A
- variables x, y, z, \dots
- constraints: $R_1(x, y, z), R_2(z, x), R_3(x, y), R_2(z, y), \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

- **Non-uniform** CSP: domain is **fixed** and only certain types of constraints are allowed

Constraint with r variables \leftrightarrow relation $R \subseteq A^r$ listing all valid assignments

Constraint language / **template**: $\mathbb{A} = (A; R_1, \dots, R_k)$

1-IN-3-SAT : $(\{0, 1\}; \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\})$

n -COLORING : $(\{1, \dots, n\}, \neq)$

ACYCLICITY : $(\mathbb{Q}; <)$

Instance:

- a ~~domain~~ A
- variables x, y, z, \dots
- constraints: $R_1(x, y, z), R_2(z, x), R_3(x, y), R_2(z, y), \dots$

Question: does there exist $h: \text{Vars} \rightarrow A$ satisfying every constraint?

- **Non-uniform** CSP: domain is **fixed** and only certain types of constraints are allowed

Constraint with r variables \leftrightarrow relation $R \subseteq A^r$ listing all valid assignments

Constraint language / **template**: $\mathbb{A} = (A; R_1, \dots, R_k)$

1-IN-3-SAT : $(\{0, 1\}; \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\})$


n -COLORING : $(\{1, \dots, n\}, \neq)$

ACYCLICITY : $(\mathbb{Q}; <)$

Theorem (Bulatov // Zhuk '17). For every template \mathbb{A} with a **finite** domain, $\text{CSP}(\mathbb{A})$ is solvable in polynomial time or NP-complete.


Finite-domain dichotomy result built on **polymorphisms** $\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$

- **Clones**: closed under composition, contain **projections**
- Standard object in **universal algebra**
- Clone **actions** of $\text{Pol}(\mathbb{A})$ tell us about the complexity of $\text{CSP}(\mathbb{A})$




Finite-domain dichotomy result built on **polymorphisms**

$$\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$$



Finite-domain dichotomy result built on **polymorphisms**

$$\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$$




- $\text{Pol}(\mathbb{A}) = \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{A}) \equiv \text{CSP}(\mathbb{B})$

(Bulatov, Jeavons, Krokhin)

Finite-domain dichotomy result built on **polymorphisms**


$$\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$$



- $\text{Pol}(\mathbb{A}) \subseteq \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{B}) \leq \text{CSP}(\mathbb{A})$

(Bulatov, Jeavons, Krokhin)

Finite-domain dichotomy result built on **polymorphisms** $\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$




- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{B}) \leq \text{CSP}(\mathbb{A})$

(Barto, Opršal, Pinsker)

Finite-domain dichotomy result built on **polymorphisms**


$$\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$$



- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{B}) \leq \text{CSP}(\mathbb{A})$
- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\text{SAT})$ implies $\text{CSP}(\mathbb{A})$ is NP-hard


(Barto, Opršal, Pinsker)

Finite-domain dichotomy result built on **polymorphisms** $\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$




- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{B}) \leq \text{CSP}(\mathbb{A})$ (Barto, Opršal, Pinsker)
- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\text{SAT})$ implies $\text{CSP}(\mathbb{A})$ is NP-hard
- If $\text{Pol}(\mathbb{A}) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time (Bulatov // Zhuk)

Finite-domain dichotomy result built on **polymorphisms** $\text{Pol}(\mathbb{A}) = \{f: \mathbb{A}^n \rightarrow \mathbb{A}, n \geq 1\}$



- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\mathbb{B})$ implies $\text{CSP}(\mathbb{B}) \leq \text{CSP}(\mathbb{A})$ (Barto, Opršal, Pinsker)
- $\text{Pol}(\mathbb{A}) \rightarrow \text{Pol}(\text{SAT})$ implies $\text{CSP}(\mathbb{A})$ is NP-hard
- If $\text{Pol}(\mathbb{A}) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time (Bulatov // Zhuk)
- Can the **algebraic approach** be leveraged for other computational problems?

Research in **infinite-domain** CSPs



- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

AND/OR scheduling

Betweenness

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\left\{ \begin{array}{lcl} x & < & y \\ y & < & z \\ z & < & t \\ t & < & x \end{array} \right.$$

AND/OR scheduling

Betweenness

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\left\{ \begin{array}{lcl} x & < & y \\ y & < & z \\ z & < & t \\ t & < & x \end{array} \right.$$

In P: check if constraint graph
is acyclic

AND/OR scheduling

Betweenness

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x < y \\ y < z \\ z < t \\ t < x \end{cases}$$

In P: check if constraint graph
is acyclic

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

Betweenness

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \iff constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x < y \\ y < z \\ z < t \\ t < x \end{cases}$$

In P: check if constraint graph
is acyclic

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\} \iff$ constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x < y \\ y < z \\ z < t \\ t < x \end{cases}$$

In P: check if constraint graph is acyclic

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

$$\begin{cases} x < y < z \vee z < y < x \\ y < z < t \vee t < z < y \\ x < z < t \vee t < z < y \end{cases}$$

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \iff constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x < y \\ y < z \\ z < t \\ t < x \end{cases}$$

In P: check if constraint graph is acyclic

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

$$\begin{cases} x < y < z \vee z < y < x \\ y < z < t \vee t < z < y \\ x < z < t \vee t < z < y \end{cases}$$

NP-complete

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \Leftrightarrow constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x \leq y + 1 \\ y \leq z - 1 \\ z \leq t + 3 \\ t \leq x - 4 \end{cases}$$

In P: check if constraint graph has negative cycles

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

$$\begin{cases} x < y < z \vee z < y < x \\ y < z < t \vee t < z < y \\ x < z < t \vee t < z < y \end{cases}$$

NP-complete

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\} \iff$ constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x \leq y + 1 \\ y \leq z - 1 \\ z \leq t + 3 \\ t \leq x - 4 \end{cases}$$

In P: check if constraint graph
has negative cycles

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

$$\begin{cases} x < y < z \vee z < y < x \\ y < z < t \vee t < z < y \\ x < z < t \vee t < z < y \end{cases}$$

NP-complete

Temporal template: domain is \mathbb{Z} , relations are definable by boolean combinations of $<$ and $y = x + c$

- Reasoning: given a collection Φ of facts and an atom ψ , does Φ imply ψ ?
- Same as unsatisfiability of $\Phi \cup \{\neg\psi\}$ \iff constraint satisfaction problem
- In spatial/temporal reasoning, variables are coordinates and the facts/atom are of the form $x < y, y = x + 5, \dots$

Examples.

Ordering constraints

$$\begin{cases} x \leq y + 1 \\ y \leq z - 1 \\ z \leq t + 3 \\ t \leq x - 4 \end{cases}$$

In P: check if constraint graph has negative cycles

AND/OR scheduling

$$\begin{cases} x < \max(y, z) \\ y < t \\ t < \max(y, x) \end{cases}$$

In P: essentially **arc consistency**

Betweenness

$$\begin{cases} x < y < z \vee z < y < x \\ y < z < t \vee t < z < y \\ x < z < t \vee t < z < y \end{cases}$$

NP-complete

Temporal template: domain is \mathbb{Z} , relations are definable by boolean combinations of $<$ and $y = x + c$

Theorem (Bodirsky-Martin-M., *J. ACM*'18). Let \mathbb{A} be a temporal template. Then $\text{CSP}(\mathbb{A})$ is in P or NP-complete. Given \mathbb{A} , it is **decidable** which case applies.

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \text{ (red) }, \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \text{ (green) }, \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \text{ (blue) } \right\}$

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\}$

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\} \rightsquigarrow \text{no-monochromatic-triangle problem}$

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\} \rightsquigarrow \text{no-monochromatic-triangle problem}$

Observation. There is \mathbb{A} such that $\text{CSP}(\mathbb{A})$ is the no-mono-triangle problem, but no **finite** \mathbb{A} .

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\} \rightsquigarrow \text{no-monochromatic-triangle problem}$

Observation. There is \mathbb{A} such that $\text{CSP}(\mathbb{A})$ is the no-mono-triangle problem, but no **finite** \mathbb{A} .

Theorem (Bodirsky-Madelaine-M., *SIAM J. Comp.* '21). Every vertex partitioning problem is in P or NP-complete. Given \mathcal{F} , it is **decidable** which case applies.

The proof uses techniques from infinite-domain constraint satisfaction.

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\} \rightsquigarrow \text{no-monochromatic-triangle problem}$

Observation. There is \mathbb{A} such that $\text{CSP}(\mathbb{A})$ is the no-mono-triangle problem, but no **finite** \mathbb{A} .

Theorem (M.-Nagy-Pinsker-Wrona, *SIAM J. Comp.* '24). Characterization of the \mathcal{F} that have bounded treewidth duality.

The proof uses techniques from infinite-domain constraint satisfaction.

Definition. Fix a set \mathcal{F} of **vertex-colored** graphs. The **vertex partitioning problem** for \mathcal{F} is:

Input: non-colored graph G ,

Question: does there exist a vertex-coloring χ such that (G, χ) contains no element of \mathcal{F} .

Examples. $\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \right\} \right\} \rightsquigarrow \text{3-coloring problem}$

$\mathcal{F} = \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} , \left\{ \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \end{array} \right\} \right\} \rightsquigarrow \text{no-monochromatic-triangle problem}$

Observation. There is \mathbb{A} such that $\text{CSP}(\mathbb{A})$ is the no-mono-triangle problem, but no **finite** \mathbb{A} .

Theorem (Bodirsky-Madelaine-M., *SIAM J. Comp.* '21). Every vertex partitioning problem is in P or NP-complete. Given \mathcal{F} , it is **decidable** which case applies.

The proof uses techniques from infinite-domain constraint satisfaction.

Project (with D. Perinti). Understand the complexity of edge partitioning problems.


Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Example (1-in-3 Satisfiability). Given a set of constraints $(x, y, z), (y, z, t), \dots$, decide if there is a map $h: V \rightarrow \{0, 1\}$ such that for every constraint (a, b, c) , exactly one of $h(a), h(b), h(c)$ is 1.

Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Example (1-in-3 Satisfiability). Given a set of constraints $(x, y, z), (y, z, t), \dots$, decide if there is a map $h: V \rightarrow \{0, 1\}$ such that for every constraint (a, b, c) , exactly one of $h(a), h(b), h(c)$ is 1.


Instance




Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Example (1-in-3 Satisfiability). Given a set of constraints $(x, y, z), (y, z, t), \dots$, decide if there is a map $h: V \rightarrow \{0, 1\}$ such that for every constraint (a, b, c) , exactly one of $h(a), h(b), h(c)$ is 1.

Instance




Possible certificates



Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Example (Betweenness). Given a set of constraints $(x, y, z), (y, z, t), \dots$, decide if there is a linear order on the variables such that for every constraint (a, b, c) , either $a < b < c$ or $c < b < a$.


Instance




Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Example (Betweenness). Given a set of constraints $(x, y, z), (y, z, t), \dots$, decide if there is a linear order on the variables such that for every constraint (a, b, c) , either $a < b < c$ or $c < b < a$.

Instance



Possible certificates




Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Conjecture (2011). Let \mathbb{A} be s.t. $\text{CSP}(\mathbb{A})$ has combinatorial certificates and $\text{Pol}(\mathbb{A}) \not\rightarrow \text{Pol}(\text{SAT})$.
Then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.

2011

Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Conjecture (2011). Let \mathbb{A} be s.t. $\text{CSP}(\mathbb{A})$ has combinatorial certificates and $\text{Pol}(\mathbb{A}) \not\rightarrow \text{Pol}(\text{SAT})$. Then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.




Early proofs:

- Proofs by case distinctions
- Victim of combinatorial explosion
- *Ad hoc* arguments
- Limited theoretical understanding

Algebraic approach works whenever $\text{CSP}(\mathbb{A})$ has **combinatorial certificates**.

Conjecture (2011). Let \mathbb{A} be s.t. $\text{CSP}(\mathbb{A})$ has combinatorial certificates and $\text{Pol}(\mathbb{A}) \not\rightarrow \text{Pol}(\text{SAT})$. Then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.



Early proofs:

- Proofs by case distinctions
- Victim of combinatorial explosion
- *Ad hoc* arguments
- Limited theoretical understanding

Modern proofs:

- Unifying algebraic theory
- No combinatorial explosion
- Generic reduction to finite-domain CSPs
- Generic uniform algorithms
- Finer understanding than P vs. NP

Theorem (Bodirsky-M. LICS'16). For every \mathbb{A} with combinatorial certificates, there exists an equivalence relation \equiv such that if $\text{Pol}(\mathbb{A}, \equiv) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.


Theorem (Bodirsky-M. LICS'16). For every \mathbb{A} with combinatorial certificates, there exists an equivalence relation \equiv such that if $\text{Pol}(\mathbb{A}, \equiv) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.

- \equiv only depends on the type of combinatorial certificates
- Uses a reduction to a finite-domain CSP combined with the Bulatov-Zhuk theorem
- $\text{Pol}(\mathbb{A}, \equiv) \subseteq \text{Pol}(\mathbb{A})$

Theorem (Bodirsky-M. LICS'16). For every \mathbb{A} with combinatorial certificates, there exists an equivalence relation \equiv such that if $\text{Pol}(\mathbb{A}, \equiv) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.

- \equiv only depends on the type of combinatorial certificates
- Uses a reduction to a finite-domain CSP combined with the Bulatov-Zhuk theorem
- $\text{Pol}(\mathbb{A}, \equiv) \subseteq \text{Pol}(\mathbb{A})$


“Only” remaining case to understand:



Theorem (Bodirsky-M. LICS'16). For every \mathbb{A} with combinatorial certificates, there exists an equivalence relation \equiv such that if $\text{Pol}(\mathbb{A}, \equiv) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.

- \equiv only depends on the type of combinatorial certificates
- Uses a reduction to a finite-domain CSP combined with the Bulatov-Zhuk theorem
- $\text{Pol}(\mathbb{A}, \equiv) \subseteq \text{Pol}(\mathbb{A})$

“Only” remaining case to understand:




2020–2025: **algebraic theory** to handle this situation in generality

Theorem (Bodirsky-M. LICS'16). For every \mathbb{A} with combinatorial certificates, there exists an equivalence relation \equiv such that if $\text{Pol}(\mathbb{A}, \equiv) \not\rightarrow \text{Pol}(\text{SAT})$, then $\text{CSP}(\mathbb{A})$ is solvable in polynomial time.

- \equiv only depends on the type of combinatorial certificates
- Uses a reduction to a finite-domain CSP combined with the Bulatov-Zhuk theorem
- $\text{Pol}(\mathbb{A}, \equiv) \subseteq \text{Pol}(\mathbb{A})$


“Only” remaining case to understand:



2020–2025: **algebraic theory** to handle this situation in generality

Invited tutorial from
2025 Dagstuhl seminar:







Thank you for your attention!



Slides




Research in Quantum Constraint Satisfaction



Verifier

Alice




Bob

Verifier

1. selects $e = (v, w)$ following distribution π


Alice



Bob


Verifier

- 1. selects $e = (v, w)$ following distribution π
- 2. sends e to Alice
- 3. sends v to Bob



Alice

- 4. Answers $(a, b) \in [k]^2$ with $a \neq b$ following distribution p_e




Bob



- 4. Answers $c \in [k]$ following distribution p_v


Verifier

- 1. selects $e = (v, w)$ following distribution π
- 2. sends e to Alice
- 3. sends v to Bob




Alice

- 4. Answers $(a, b) \in [k]^2$ with $a \neq b$ following distribution p_e




Bob

- 4. Answers $c \in [k]$ following distribution p_v




Verifier

- 1. selects $e = (v, w)$ following distribution π
- 2. sends e to Alice
- 3. sends v to Bob



Alice

- 4. Answers $(a, b) \in [k]^2$ with $a \neq b$ following distribution p_e



Bob

- 4. Answers $c \in [k]$ following distribution p_v



- Strategy for Alice: for each $e \in E$, a probability distribution p_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a probability distribution p_v on $[k]$
- Winning probability of a strategy: $\Pr_{e \sim \pi, (a, b) \sim p_e, c \sim p_v}[a = c]$

Observation. Alice and Bob win with probability 1 if, and only if, G is k -colorable.


- Strategy for Alice: for each $e \in E$, a probability distribution p_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a probability distribution p_v on $[k]$
- Winning probability of a strategy: $\Pr_{e \sim \pi, (a,b) \sim p_e, c \sim p_v}[a = c]$

Observation. Alice and Bob win with probability 1 if, and only if, G is k -colorable.

What if Alice and Bob share an entangled state $|q\rangle$?

- Strategy for Alice: for each $e \in E$, a probability distribution p_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a probability distribution p_v on $[k]$
- Winning probability of a strategy: $\Pr_{e \sim \pi, (a,b) \sim p_e, c \sim p_v}[a = c]$


Observation. Alice and Bob win with probability 1 if, and only if, G is k -colorable.
What if Alice and Bob share an entangled state $|q\rangle$?



- Strategy for Alice: for each $e \in E$, a probability distribution p_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a probability distribution p_v on $[k]$
- Winning probability of a strategy: $\Pr_{e \sim \pi, (a,b) \sim p_e, c \sim p_v}[a = c]$

Observation. Alice and Bob win with probability 1 if, and only if, G is k -colorable.

What if Alice and Bob share an entangled state $|q\rangle$?



How Alice performs a measurement:


- Choose projectors $Q_{a,b}$ summing to id
- Output of measurement is (a, b) with probability $\langle q|Q_{a,b}|q\rangle$

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.



Quantum 4-colorable graph with chromatic number 5
[Mańćinska-Roberson '18]

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.

Theorem (Culf-Mastel, FOCS'25). There exists $s \in (0, 1)$ such that it is undecidable to distinguish graphs that are quantum **3**-colorable from those where every strategy wins with probability $< s$.

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.

Theorem (Culf-Mastel, FOCS'25). There exists $s \in (0, 1)$ such that it is undecidable to distinguish graphs that are quantum **3**-colorable from those where every strategy wins with probability $< s$.

- Similar flavor to the **PCP theorem** in classical complexity
- Hardness of approximation result in the quantum setting
- Unlike in classical complexity, many reductions still unknown
- Relies on **commutativity gadgets** for 3-coloring (Ji, 2013)

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.

Theorem (Culf-Mastel, FOCS'25). There exists $s \in (0, 1)$ such that it is undecidable to distinguish graphs that are quantum **3**-colorable from those where every strategy wins with probability $< s$.

- Similar flavor to the **PCP theorem** in classical complexity
- Hardness of approximation result in the quantum setting
- Unlike in classical complexity, many reductions still unknown
- Relies on **commutativity gadgets** for 3-coloring (Ji, 2013)

Theorem (Ciardo-M., 2025+). k -coloring has commutativity gadgets for all $k \geq 3$.

Algebraic characterization of the existence of commutativity gadgets using **polymorphisms**.

- Strategy for Alice: for each $e \in E$, a **measurement** Q_e on $\{(a, b) \in [k]^2 \mid a \neq b\}$
- Strategy for Bob: for each $v \in V$, a **measurement** Q_v on $[k]$
- Winning probability of a strategy: $\sum_e \sum_{(a,b)} \pi(e) \langle q | Q_{e,ab} \otimes Q_{v,a} | q \rangle$

Definition. Say G is **quantum** k -colorable if there is a quantum strategy that wins with probability 1.

Theorem (Culf-Mastel, FOCS'25). There exists $s \in (0, 1)$ such that it is undecidable to distinguish graphs that are quantum **3**-colorable from those where every strategy wins with probability $< s$.


- Similar flavor to the **PCP theorem** in classical complexity
- Hardness of approximation result in the quantum setting
- Unlike in classical complexity, many reductions still unknown
- Relies on **commutativity gadgets** for 3-coloring (Ji, 2013)

Theorem (Ciardo-M., 2025+). k -coloring has commutativity gadgets for all $k \geq 3$.

Algebraic characterization of the existence of commutativity gadgets using **polymorphisms**.

Project (with L. Ciardo and G. Joubert). Understand the complexity landscape for entangled CSPs.

Research in Promise Constraint Satisfaction



Solving CSPs under structural assumptions on the **instances**:

Definition. $\text{PCSP}(\mathbb{A}, \mathbb{B})$ is the problem to find a satisfying assignment in \mathbb{B} , under the **promise** that there exists one in \mathbb{A} .

Solving CSPs under structural assumptions on the **instances**:

Definition. $\text{PCSP}(\mathbb{A}, \mathbb{B})$ is the problem to find a satisfying assignment in \mathbb{B} , under the **promise** that there exists one in \mathbb{A} .

Example. $\text{PCSP}(\mathbb{K}_3, \mathbb{K}_6)$

Given an undirected **3-colorable** graph G , find a 6-coloring.

Unknown complexity, related to the existence of a **2-approximation algorithm** for the chromatic number.

Solving CSPs under structural assumptions on the **instances**:

Definition. $\text{PCSP}(\mathbb{A}, \mathbb{B})$ is the problem to find a satisfying assignment in \mathbb{B} , under the **promise** that there exists one in \mathbb{A} .

Example. $\text{PCSP}(\mathbb{K}_3, \mathbb{K}_6)$

Given an undirected **3-colorable** graph G , find a 6-coloring.

Unknown complexity, related to the existence of a **2-approximation algorithm** for the chromatic number.

- Massive framework for the study of approximation problems
- Existence of a suitable generalization to the algebraic approach

Solving CSPs under structural assumptions on the **instances**:

Definition. $\text{PCSP}(\mathbb{A}, \mathbb{B})$ is the problem to find a satisfying assignment in \mathbb{B} , under the **promise** that there exists one in \mathbb{A} .

Example. $\text{PCSP}(\mathbb{K}_3, \mathbb{K}_6)$


Given an undirected **3-colorable** graph G , find a 6-coloring.

Unknown complexity, related to the existence of a **2-approximation algorithm** for the chromatic number.


- Massive framework for the study of approximation problems
- Existence of a suitable generalization to the algebraic approach
- Surprising connection between **finite-domain** PCSPs and **combinatorial** CSPs:

Theorem (M. LICS'25). The following hold:


- Combinatorial CSP $\simeq \text{PCSP}(\text{combinatorial}, \text{finite})$
- There are finite PCSPs whose tractability can be shown by combinatorial CSPs and not by finite CSPs.
- There exists a uniform algorithm for temporal CSPs obtained by a reduction to finite PCSPs.



Research in Optimization




k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$



k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$


Value of ℓ : $\#\{(u, v) \in E \mid \ell(u) < \ell(v)\}$



k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$

Value of ℓ : $\#\{(u, v) \in E \mid \ell(u) < \ell(v)\}$

Note: every layering with value ρ is an acyclic subgraph with ρ edges




k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$


Value of ℓ : $\#\{(u, v) \in E \mid \ell(u) < \ell(v)\}$

Note: every layering with value ρ is an acyclic subgraph with ρ edges

Approximation diagram for
Maximum Acyclic Subgraph



Given a digraph with an acyclic subgraph of weight $\geq c|E|$,
find an acyclic subgraph with weight $\geq s|E|$.




k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$


Value of ℓ : $\#\{(u, v) \in E \mid \ell(u) < \ell(v)\}$

Note: every layering with value ρ is an acyclic subgraph with ρ edges

Approximation diagram for
Maximum Acyclic Subgraph



Given a digraph with an acyclic subgraph of weight $\geq c|E|$,
find an acyclic subgraph with weight $\geq s|E|$.




PCSP($\text{LO}_k, \text{LO}_\infty$) (maximization version): given a directed graph G admitting a k -layering of value ρ , find an acyclic subgraph of G containing at least ρ edges.

k -layering of a digraph: map $\ell: V \rightarrow \{1, \dots, k\}$


Value of ℓ : $\#\{(u, v) \in E \mid \ell(u) < \ell(v)\}$

Note: every layering with value ρ is an acyclic subgraph with ρ edges

Approximation diagram for
Maximum Acyclic Subgraph



Given a digraph with an acyclic subgraph of weight $\geq c|E|$,
find an acyclic subgraph with weight $\geq s|E|$.



PCSP($\text{LO}_k, \text{LO}_\infty$) (maximization version): given a directed graph G admitting a k -layering of value ρ , find an acyclic subgraph of G containing at least ρ edges.

- Solvable in polynomial time for $k = 2$ (Nakajima-Živný, **M.** '25)
- NP-hard for $k \geq 4$, but unknown tractability boundary in approximation diagram
- Open complexity for $k = 3$