

Reducts of finitely bounded structures, and lifting tractability from finite-domain constraint satisfaction

Antoine Mottet

May 11, 2016, PhDs in Logic

- ▶ Relational structure: $\mathcal{A} = (A, R_1^{\mathcal{A}}, \dots, R_k^{\mathcal{A}})$ with $R_i^{\mathcal{A}} \subseteq A^{r_i}$

- ▶ Relational structure: $\mathcal{A} = (A, R_1^{\mathcal{A}}, \dots, R_k^{\mathcal{A}})$ with $R_i^{\mathcal{A}} \subseteq A^{r_i}$
- ▶ A **homomorphism** $f: \mathcal{A} \rightarrow \mathcal{B}$ is a function such that

$$\forall R_i, \forall (a_1, \dots, a_{r_i}) \in R_i^{\mathcal{A}}, (f(a_1), \dots, f(a_{r_i})) \in R_i^{\mathcal{B}}$$

- ▶ Relational structure: $\mathcal{A} = (A, R_1^{\mathcal{A}}, \dots, R_k^{\mathcal{A}})$ with $R_i^{\mathcal{A}} \subseteq A^{r_i}$
- ▶ A **homomorphism** $f: \mathcal{A} \rightarrow \mathcal{B}$ is a function such that

$$\forall R_i, \forall (a_1, \dots, a_{r_i}) \in R_i^{\mathcal{A}}, (f(a_1), \dots, f(a_{r_i})) \in R_i^{\mathcal{B}}$$

Let \mathcal{A} be a relational structure, in a fixed finite signature τ . We consider the following problem:

Definition ($\text{Hom}(\mathcal{A})$)

Input: a **finite** τ -structure \mathcal{B}

Question: decide the existence of a **homomorphism** $\mathcal{B} \rightarrow \mathcal{A}$.

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question:

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question: Is \mathcal{B} 3-colourable?

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question: Is \mathcal{B} 3-colourable?

Complexity: NP-complete

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question: Is \mathcal{B} 3-colourable?

Complexity: NP-complete

Example ($\text{Hom}(\mathbb{Z}, <)$)

Input: a finite directed graph \mathcal{B}

Question:

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question: Is \mathcal{B} 3-colourable?

Complexity: NP-complete

Example ($\text{Hom}(\mathbb{Z}, <)$)

Input: a finite directed graph \mathcal{B}

Question: Is \mathcal{B} acyclic?

Example ($\text{Hom}(K_3)$)

Input: a finite graph \mathcal{B}

Question: Is \mathcal{B} 3-colourable?

Complexity: NP-complete

Example ($\text{Hom}(\mathbb{Z}, <)$)

Input: a finite directed graph \mathcal{B}

Question: Is \mathcal{B} acyclic?

Complexity: linear time

Let again \mathcal{A} be a τ -structure.

Definition ($\text{CSP}(\mathcal{A})$)

Input: a first-order sentence ϕ of the form

$$\exists x_1 \dots \exists x_k \bigwedge_i R_i(x_{i1}, \dots, x_{is_i}) \quad (R_i \in \tau)$$

Question: $\mathcal{A} \models \phi$?

Let again \mathcal{A} be a τ -structure.

Definition ($\text{CSP}(\mathcal{A})$)

Input: a first-order sentence ϕ of the form

$$\exists x_1 \dots \exists x_k \bigwedge_i R_i(x_{i1}, \dots, x_{is_i}) \quad (R_i \in \tau)$$

Question: $\mathcal{A} \models \phi$?

Example ($\text{CSP}(\mathbb{Z}; +, \times)$)

Input: polynomials f_1, \dots, f_m

Question:

Let again \mathcal{A} be a τ -structure.

Definition ($\text{CSP}(\mathcal{A})$)

Input: a first-order sentence ϕ of the form

$$\exists x_1 \dots \exists x_k \bigwedge_i R_i(x_{i1}, \dots, x_{is_i}) \quad (R_i \in \tau)$$

Question: $\mathcal{A} \models \phi$?

Example ($\text{CSP}(\mathbb{Z}; +, \times)$)

Input: polynomials f_1, \dots, f_m

Question: do f_1, \dots, f_m have a common integer root?

Complexity:

Let again \mathcal{A} be a τ -structure.

Definition ($\text{CSP}(\mathcal{A})$)

Input: a first-order sentence ϕ of the form

$$\exists x_1 \dots \exists x_k \bigwedge_i R_i(x_{i1}, \dots, x_{is_i}) \quad (R_i \in \tau)$$

Question: $\mathcal{A} \models \phi$?

Example ($\text{CSP}(\mathbb{Z}; +, \times)$)

Input: polynomials f_1, \dots, f_m

Question: do f_1, \dots, f_m have a common integer root?

Complexity: **undecidable**.

Let again \mathcal{A} be a τ -structure.

Definition ($\text{CSP}(\mathcal{A})$)

Input: a first-order sentence ϕ of the form

$$\exists x_1 \dots \exists x_k \bigwedge_i R_i(x_{i1}, \dots, x_{is_i}) \quad (R_i \in \tau)$$

Question: $\mathcal{A} \models \phi$?

Example ($\text{CSP}(\mathbb{Z}; +, \times)$)

Input: polynomials f_1, \dots, f_m

Question: do f_1, \dots, f_m have a common integer root?

Complexity: undecidable.

$\text{CSP}(\mathcal{A})$ and $\text{Hom}(\mathcal{A})$ are equivalent.

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete.

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete. Confirmed for:

- ▶ structures of size up to 5 (Schaefer '78, Bulatov '06, Zhuk '16)

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete. Confirmed for:

- ▶ structures of size up to 5 (Schaefer '78, Bulatov '06, Zhuk '16)
- ▶ undirected graphs (Hell-Nešetřil '90)

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete. Confirmed for:

- ▶ structures of size up to 5 (Schaefer '78, Bulatov '06, Zhuk '16)
- ▶ undirected graphs (Hell-Nešetřil '90)
- ▶ smooth digraphs (Barto-Kozik-Niven '09)

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete. Confirmed for:

- ▶ structures of size up to 5 (Schaefer '78, Bulatov '06, Zhuk '16)
- ▶ undirected graphs (Hell-Nešetřil '90)
- ▶ smooth digraphs (Barto-Kozik-Niven '09)
- ▶ ...

Conjecture (Feder-Vardi '93): $\text{CSP}(\mathcal{A})$ for a structure \mathcal{A} with finite domain is in P or NP-complete. Confirmed for:

- ▶ structures of size up to 5 (Schaefer '78, Bulatov '06, Zhuk '16)
- ▶ undirected graphs (Hell-Nešetřil '90)
- ▶ smooth digraphs (Barto-Kozik-Niven '09)
- ▶ ...

Main tool: universal-algebraic approach.

Definition (Polymorphism)

A k -ary **polymorphism** of \mathcal{A} is a homomorphism $\mathcal{A}^k \rightarrow \mathcal{A}$.

Definition (Polymorphism)

A k -ary **polymorphism** of \mathcal{A} is a homomorphism $\mathcal{A}^k \rightarrow \mathcal{A}$.

Polymorphisms are used to **combine homomorphisms**: if $h_1, \dots, h_k: \mathcal{B} \rightarrow \mathcal{A}$ then $f(h_1, \dots, h_k): \mathcal{B} \rightarrow \mathcal{A}$ is (hopefully) a better homomorphism.

Definition (Polymorphism)

A k -ary **polymorphism** of \mathcal{A} is a homomorphism $\mathcal{A}^k \rightarrow \mathcal{A}$.

Polymorphisms are used to **combine homomorphisms**: if $h_1, \dots, h_k: \mathcal{B} \rightarrow \mathcal{A}$ then $f(h_1, \dots, h_k): \mathcal{B} \rightarrow \mathcal{A}$ is (hopefully) a better homomorphism.

Example

Suppose that \mathcal{A} has $f(x_1, \dots, x_k) = a$ as a polymorphism for some $a \in \mathcal{A}$. If $h: \mathcal{B} \rightarrow \mathcal{A}$ is a homomorphism, then $f(h, \dots, h): x \mapsto a$ is also a homomorphism.

Definition (Polymorphism)

A k -ary **polymorphism** of \mathcal{A} is a homomorphism $\mathcal{A}^k \rightarrow \mathcal{A}$.

Polymorphisms are used to **combine homomorphisms**: if $h_1, \dots, h_k: \mathcal{B} \rightarrow \mathcal{A}$ then $f(h_1, \dots, h_k): \mathcal{B} \rightarrow \mathcal{A}$ is (hopefully) a better homomorphism.

Example

Suppose that \mathcal{A} has $f(x_1, \dots, x_k) = a$ as a polymorphism for some $a \in \mathcal{A}$. If $h: \mathcal{B} \rightarrow \mathcal{A}$ is a homomorphism, then $f(h, \dots, h): x \mapsto a$ is also a homomorphism.

Concrete example: a group G , $f: x \mapsto e_G$.

Conjecture (Bulatov-Jeavons-Krokhin '05)

Let \mathcal{A} be a *finite structure*. Then:

- ▶ \mathcal{A} has a *cyclic polymorphism* – something satisfying

$$f(x_1, \dots, x_k) = f(x_2, \dots, x_k, x_1),$$

and $\text{CSP}(\mathcal{A})$ is in P ,

- ▶ or $\text{CSP}(\mathcal{A})$ is *NP-complete*.

Conjecture (Bulatov-Jeavons-Krokhin '05)

Let \mathcal{A} be a *finite structure*. Then:

- ▶ \mathcal{A} has a *cyclic polymorphism* – something satisfying

$$f(x_1, \dots, x_k) = f(x_2, \dots, x_k, x_1),$$

and $\text{CSP}(\mathcal{A})$ is in P ,

- ▶ or $\text{CSP}(\mathcal{A})$ is *NP-complete*.

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:
 - ▶ The powerful results of universal algebra only hold for **finite** structures,

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:
 - ▶ The powerful results of universal algebra only hold for **finite** structures,
 - ▶ hopefully allows to reduce the combinatorial difficulties to the “finite blackbox”.

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:
 - ▶ The powerful results of universal algebra only hold for **finite** structures,
 - ▶ hopefully allows to reduce the combinatorial difficulties to the “finite blackbox”.
- ▶ How:

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:
 - ▶ The powerful results of universal algebra only hold for **finite** structures,
 - ▶ hopefully allows to reduce the combinatorial difficulties to the “finite blackbox”.
- ▶ How:
 - ▶ structures for which the CSP is “manageable”: **finitely bounded** structures,

- ▶ Goal: if \mathcal{A} is an infinite structure, try to find a **finite** structure \mathcal{A}' such that $\text{CSP}(\mathcal{A})$ reduces to $\text{CSP}(\mathcal{A}')$.
- ▶ Why:
 - ▶ The powerful results of universal algebra only hold for **finite** structures,
 - ▶ hopefully allows to reduce the combinatorial difficulties to the “finite blackbox”.
- ▶ How:
 - ▶ structures for which the CSP is “manageable”: **finitely bounded** structures,
 - ▶ structures for which finite substructures can move around: **homogeneity**

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs.

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs. By induction, one can glue all those graphs together to obtain a countable graph: **the random graph** \mathcal{G} .

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs. By induction, one can glue all those graphs together to obtain a countable graph: **the random graph** \mathcal{G} .

All we need for the induction is the following:

Definition

A class \mathcal{K} of finite structures is said to be an **amalgamation class** if for every $\mathcal{A}, \mathcal{B}_1, \mathcal{B}_2 \in \mathcal{K}$

\mathcal{A} \mathcal{B}_1

\mathcal{B}_2

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs. By induction, one can glue all those graphs together to obtain a countable graph: **the random graph** \mathcal{G} .

All we need for the induction is the following:

Definition

A class \mathcal{K} of finite structures is said to be an **amalgamation class** if for every $\mathcal{A}, \mathcal{B}_1, \mathcal{B}_2 \in \mathcal{K}$ and every embedding $e_i: \mathcal{A} \rightarrow \mathcal{B}_i$,

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{B}_1 \\ \downarrow & & \\ & & \mathcal{B}_2 \end{array}$$

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs. By induction, one can glue all those graphs together to obtain a countable graph: **the random graph** \mathcal{G} .

All we need for the induction is the following:

Definition

A class \mathcal{K} of finite structures is said to be an **amalgamation class** if for every $\mathcal{A}, \mathcal{B}_1, \mathcal{B}_2 \in \mathcal{K}$ and every embedding $e_i: \mathcal{A} \rightarrow \mathcal{B}_i$, there exists $\mathcal{C} \in \mathcal{K}$

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{B}_1 \\ \downarrow & & \\ \mathcal{B}_2 & & \mathcal{C} \end{array}$$

Idea: build a countably infinite structures by gluing finite structures from some class \mathcal{K} .

Example

\mathcal{K} = class of finite simple graphs. By induction, one can glue all those graphs together to obtain a countable graph: **the random graph** \mathcal{G} .

All we need for the induction is the following:

Definition

A class \mathcal{K} of finite structures is said to be an **amalgamation class** if for every $\mathcal{A}, \mathcal{B}_1, \mathcal{B}_2 \in \mathcal{K}$ and every embedding $e_i: \mathcal{A} \rightarrow \mathcal{B}_i$, there exists $\mathcal{C} \in \mathcal{K}$ and embeddings $f_i: \mathcal{B}_i \rightarrow \mathcal{C}$ with $f_1 \circ e_1 = f_2 \circ e_2$.

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{B}_1 \\ \downarrow & & \downarrow \\ \mathcal{B}_2 & \dashrightarrow & \mathcal{C} \end{array}$$

Idea: only work with classes \mathcal{K} which can be finitely represented.
A Fraïssé class \mathcal{K} is **finitely bounded** if. . .

Idea: only work with classes \mathcal{K} which can be finitely represented.
A Fraïssé class \mathcal{K} is **finitely bounded** if. . . it has finitely many bounds (see board).

Idea: only work with classes \mathcal{K} which can be finitely represented.
A Fraïssé class \mathcal{K} is **finitely bounded** if. . . it has finitely many bounds (see board).

Lemma

Let \mathcal{A} be a finitely bounded homogeneous structure, and let \mathcal{B} be first-order definable in \mathcal{A} . Then $\text{CSP}(\mathcal{B})$ is in NP.

Definition

A function $f: A^k \rightarrow A$ is **canonical with respect to \mathcal{A}** if for every finite $S \subset A$ and $\alpha_1, \dots, \alpha_k \in \text{Aut}(\mathcal{A})$, there exists $\beta \in \text{Aut}(\mathcal{A})$ such that

$$\forall x_1, \dots, x_k \in S, f(\alpha_1 x_1, \dots, \alpha_k x_k) = \beta f(x_1, \dots, x_k).$$

Definition

A function $f: A^k \rightarrow A$ is **canonical with respect to \mathcal{A}** if for every finite $S \subset A$ and $\alpha_1, \dots, \alpha_k \in \text{Aut}(\mathcal{A})$, there exists $\beta \in \text{Aut}(\mathcal{A})$ such that

$$\forall x_1, \dots, x_k \in S, f(\alpha_1 x_1, \dots, \alpha_k x_k) = \beta f(x_1, \dots, x_k).$$

Example

Let \mathcal{G} be the countable random graph. There is an infinite clique in \mathcal{G} , say K_∞ . Then an injection $f: \mathcal{G} \rightarrow K_\infty$ is canonical with respect to \mathcal{G} .

Theorem (Bodirsky-M, LICS '16)

Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} .

Theorem (Bodirsky-M, LICS '16)

*Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a **finite** structure $T_{\mathcal{B}}(\mathcal{A})$ such that:*

Theorem (Bodirsky-M, LICS '16)

*Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a **finite** structure $T_{\mathcal{B}}(\mathcal{A})$ such that:*

- ▶ *if f is a polymorphism of \mathcal{A} which is canonical with respect to \mathcal{B} , it induces a polymorphism of $T_{\mathcal{B}}(\mathcal{A})$,*

Theorem (Bodirsky-M, LICS '16)

*Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a **finite** structure $T_{\mathcal{B}}(\mathcal{A})$ such that:*

- ▶ *if f is a polymorphism of \mathcal{A} which is canonical with respect to \mathcal{B} , it induces a polymorphism of $T_{\mathcal{B}}(\mathcal{A})$,*
- ▶ *there is a polynomial-time reduction from $\text{CSP}(\mathcal{A})$ to $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$.*

Theorem (Bodirsky-M, LICS '16)

*Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a **finite** structure $T_{\mathcal{B}}(\mathcal{A})$ such that:*

- ▶ *if f is a polymorphism of \mathcal{A} which is canonical with respect to \mathcal{B} , it induces a polymorphism of $T_{\mathcal{B}}(\mathcal{A})$,*
- ▶ *there is a polynomial-time reduction from $\text{CSP}(\mathcal{A})$ to $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$.*

Question: Is $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ reducible to $\text{CSP}(\mathcal{A})$?

Theorem (Bodirsky-M, LICS '16)

Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a *finite* structure $T_{\mathcal{B}}(\mathcal{A})$ such that:

- ▶ if f is a polymorphism of \mathcal{A} which is canonical with respect to \mathcal{B} , it induces a polymorphism of $T_{\mathcal{B}}(\mathcal{A})$,
- ▶ there is a polynomial-time reduction from $\text{CSP}(\mathcal{A})$ to $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$.

Question: Is $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ reducible to $\text{CSP}(\mathcal{A})$?

Proposition

No

Theorem (Bodirsky-M, LICS '16)

Let \mathcal{A} be first-order definable in a finitely bounded homogeneous structure \mathcal{B} . There exists a *finite* structure $T_{\mathcal{B}}(\mathcal{A})$ such that:

- ▶ if f is a polymorphism of \mathcal{A} which is canonical with respect to \mathcal{B} , it induces a polymorphism of $T_{\mathcal{B}}(\mathcal{A})$,
- ▶ there is a polynomial-time reduction from $\text{CSP}(\mathcal{A})$ to $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$.

Question: Is $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ reducible to $\text{CSP}(\mathcal{A})$?

Proposition

No (in general).

Question: find the biggest class of structures for which $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are equivalent. This class contains (a posteriori):

- ▶ equality constraints,

Question: find the biggest class of structures for which $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are equivalent. This class contains (a posteriori):

- ▶ equality constraints,
- ▶ structures definable over the random graph (in fact, over *any* homogeneous undirected graph),

Question: find the biggest class of structures for which $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are equivalent. This class contains (a posteriori):

- ▶ equality constraints,
- ▶ structures definable over the random graph (in fact, over *any* homogeneous undirected graph),
- ▶ some structures definable over the random partially-ordered set,

Question: find the biggest class of structures for which $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are equivalent. This class contains (a posteriori):

- ▶ equality constraints,
- ▶ structures definable over the random graph (in fact, over *any* homogeneous undirected graph),
- ▶ some structures definable over the random partially-ordered set,
- ▶ ...

Theorem (Bodirsky-M, LICS '16)

Let \mathcal{A} be a finite-signature structure that is first-order definable in $\mathcal{B} = (\mathbb{N}; 0, 1, \dots)$. Then $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are polynomial-time equivalent.

Theorem (Bodirsky-M, LICS '16)

Let \mathcal{A} be a finite-signature structure that is first-order definable in $\mathcal{B} = (\mathbb{N}; 0, 1, \dots)$. Then $\text{CSP}(\mathcal{A})$ and $\text{CSP}(T_{\mathcal{B}}(\mathcal{A}))$ are polynomial-time equivalent.

Corollary

The *algebraic conjecture* for finite-domain CSPs is *equivalent* to the statement: if \mathcal{A} is definable in $(\mathbb{N}, 0, 1, \dots)$, then:

- ▶ if \mathcal{A} has a cyclic polymorphism modulo endomorphisms, then $\text{CSP}(\mathcal{A})$ is in P ,
- ▶ or $\text{CSP}(\mathcal{A})$ is NP -hard.

Proof based on *clone homomorphisms*.

Open problem: can we lift hardness for all structures \mathcal{A} that are first-order interpretable over $(\mathbb{N}; 0, 1, \dots)$? So-called structures definable with atoms.

Open problem: can we **lift hardness** for all structures \mathcal{A} that are first-order **interpretable** over $(\mathbb{N}; 0, 1, \dots)$? So-called **structures definable with atoms**. In fact we make the following conjecture:

Conjecture

*Let \mathcal{A} be a finite-signature structure such that $\text{CSP}(\mathcal{A})$ is definable in **MMSNP**. Then there is a polynomial-time reduction from $\text{CSP}(T(\mathcal{A}))$ to $\text{CSP}(\mathcal{A})$.*

Let \mathcal{A} be a structure. There is a natural partition of A^n for all n :



Let \mathcal{A} be a structure. There is a natural partition of A^n for all n :



Example

Let $\mathcal{A} = (\mathbb{Q}; <)$. $\text{Aut}(\mathcal{A})$ is the group of increasing bijections on \mathbb{Q} .

- ▶ $n = 1$: only one orbit
- ▶ $n = 2$: three orbits given by $x < y$, $x = y$, $x > y$.

The **domain** of $T_{\mathcal{B}}(\mathcal{A})$ is the set of orbits of A^m under $\text{Aut}(\mathcal{B})$.

The **relations** of $T_{\mathcal{B}}(\mathcal{A})$ are:

- ▶ “the tuples of the orbit x is in the relation R of \mathcal{A} ”, for each relation of \mathcal{A} ,
- ▶ “the tuples of the orbits x and y are in the same orbit, if we restrict them to I and J ”, for each $I, J \subseteq \{1, \dots, m\}$.