

# MMSNP: An algebraic proof of the dichotomy

Manuel Bodirsky, **Antoine Mottet**

October 12, 2018

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity:

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:




- ▶ Complexity: NP-complete.

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.
- 

- ▶ Input: undirected graph  $G$
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity: in P.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg($$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x})) \quad )$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ ,



In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ ,
- ▶  $\beta_i$  is a conjunction of literals.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ , (**Monotone**)
- ▶  $\beta_i$  is a conjunction of literals.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates, (**Monadic**)
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ , (**Monotone**)
- ▶  $\beta_i$  is a conjunction of literals.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates, (**Monadic**)
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ , (**Monotone**)
- ▶  $\beta_i$  is a conjunction of literals.

(**Strict NP**)

## Theorem (Bulatov, Zhuk '17)

*Finite-domain CSPs have a complexity dichotomy.*

**Theorem (Bulatov, Zhuk '17)**

*Finite-domain CSPs have a complexity dichotomy.*

**Theorem (Feder-Vardi '93, Kun '13)**

*MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.*

**Theorem (Bulatov, Zhuk '17)**

*Finite-domain CSPs have a complexity dichotomy.*

**Theorem (Feder-Vardi '93, Kun '13)**

*MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.*

**Proof.**

- ▶  $\text{finite CSP} \subseteq \text{MMSNP}$ ,
- ▶ randomized reduction (Feder-Vardi) from MMSNP to CSP,
- ▶ derandomization by Kun (expander structures).



- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.



- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.

## Theorem

*Let  $\mathcal{A}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{A})$  is in MMSNP. Then one of the following holds:*

- ▶ *there is a uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{A}) \rightarrow \mathcal{P}$ , and  $\text{CSP}(\mathcal{A})$  is NP-complete,*
- ▶  *$\mathcal{A}$  has a pseudo-Siggers polymorphism and  $\text{CSP}(\mathcal{A})$  is in P.*

- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.

## Theorem

*Let  $\mathcal{A}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{A})$  is in MMSNP. Then one of the following holds:*

- ▶ *there is a uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{A}) \rightarrow \mathcal{P}$ , and  $\text{CSP}(\mathcal{A})$  is NP-complete,*
- ▶  *$\mathcal{A}$  has a pseudo-Siggers polymorphism and  $\text{CSP}(\mathcal{A})$  is in P.*

In particular, this confirms the infinite-domain conjecture for CSPs in MMSNP.

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

## Definition

$\mathcal{B} = (B; E)$  a **graph**.

$\text{CSP}(\mathcal{B})$  is the problem:

- ▶ **Input:** a **finite** graph  $\mathcal{A}$ ,
- ▶ **Question:** is there a homomorphism  $\mathcal{A} \rightarrow \mathcal{B}$ ?

## Definition

$\mathcal{B} = (B; E)$  a **graph**.

$\text{CSP}(\mathcal{B})$  is the problem:

- ▶ **Input:** a **finite** graph  $\mathcal{A}$ ,
- ▶ **Question:** is there a homomorphism  $\mathcal{A} \rightarrow \mathcal{B}$ ?

In general, the forbidden patterns problem (FPP) for  $\mathcal{F}$  is **not a CSP**, but a **finite union** of CSPs.

## Proposition

*Every FPP reduces in polynomial-time to a finite number of FPP of connected structures.*

**Theorem (Cherlin-Shelah-Shi, '99)**

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

**Theorem (Cherlin-Shelah-Shi, '99)**

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

Construct a new  $\mathcal{B}$  by:

- ▶ deleting the uncoloured elements in  $\mathcal{B}^*$ ,
- ▶ forgetting about the colours.

For this talk: we call  $\mathcal{B}$  an **MMSNP structure**.

### Theorem (Cherlin-Shelah-Shi, '99)

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

Construct a new  $\mathcal{B}$  by:

- ▶ deleting the uncoloured elements in  $\mathcal{B}^*$ ,
- ▶ forgetting about the colours.

For this talk: we call  $\mathcal{B}$  an **MMSNP structure**.

### Proposition (Bodirsky-Dalmau, '06)

$$\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F}).$$

Moreover,  $\mathcal{B}$  belongs to the class of **reducts of finitely bounded homogeneous structures**.




Obstructions  $\mathcal{F}$ :

Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :



Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :






Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :



Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

- ▶  $\text{Pol}(\mathcal{B})$ : clone of homomorphisms  $\mathcal{B}^k \rightarrow \mathcal{B}$ ,

- ▶  $\text{Pol}(\mathcal{B})$ : clone of homomorphisms  $\mathcal{B}^k \rightarrow \mathcal{B}$ ,
- ▶ Clonoid homomorphism:  
$$f(x_{i_1}, \dots, x_{i_k}) \approx g(x_{j_1}, \dots, x_{j_l}) \Rightarrow \xi(f)(\dots) \approx \xi(g)(\dots),$$

- ▶  $\text{Pol}(\mathcal{B})$ : clone of homomorphisms  $\mathcal{B}^k \rightarrow \mathcal{B}$ ,
- ▶ Clonoid homomorphism:  
$$f(x_{i_1}, \dots, x_{i_k}) \approx g(x_{j_1}, \dots, x_{j_l}) \Rightarrow \xi(f)(\dots) \approx \xi(g)(\dots),$$
- ▶  $\mathcal{A}$  finitely bounded homogeneous structure  $\Leftrightarrow$  exists finite universal theory s.t.  $\mathcal{C} \hookrightarrow \mathcal{A}$  iff  $\mathcal{C}$  model of the theory.



- ▶  $\text{Pol}(\mathcal{B})$ : clone of homomorphisms  $\mathcal{B}^k \rightarrow \mathcal{B}$ ,
- ▶ Clonoid homomorphism:  
 $f(x_{i_1}, \dots, x_{i_k}) \approx g(x_{j_1}, \dots, x_{j_l}) \Rightarrow \xi(f)(\dots) \approx \xi(g)(\dots)$ ,
- ▶  $\mathcal{A}$  finitely bounded homogeneous structure  $\Leftrightarrow$  exists finite universal theory s.t.  $\mathcal{C} \hookrightarrow \mathcal{A}$  iff  $\mathcal{C}$  model of the theory.

### Conjecture (Bodirsky-Pinsker, '11 (rephrased))

*Let  $\mathcal{B}$  be a reduct of a finitely bounded homogeneous structure. If there is no uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

- ▶  $\text{Pol}(\mathcal{B})$ : clone of homomorphisms  $\mathcal{B}^k \rightarrow \mathcal{B}$ ,
- ▶ Clonoid homomorphism:  
 $f(x_{i_1}, \dots, x_{i_k}) \approx g(x_{j_1}, \dots, x_{j_l}) \Rightarrow \xi(f)(\dots) \approx \xi(g)(\dots)$ ,
- ▶  $\mathcal{A}$  finitely bounded homogeneous structure  $\Leftrightarrow$  exists finite universal theory s.t.  $\mathcal{C} \hookrightarrow \mathcal{A}$  iff  $\mathcal{C}$  model of the theory.

### Conjecture (Bodirsky-Pinsker, '11 (rephrased))

*Let  $\mathcal{B}$  be a reduct of a finitely bounded homogeneous structure. If there is no uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in P.*

Interesting?

- ▶ statement and its consequences: ★★☆☆☆
- ▶ proofs: ★★★★★

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)


*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)

$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?

Rephrased: do  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$  and  $\text{CSP}(\mathcal{B})$  have same complexity?



A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)

*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*

### Proposition (Bodirsky, '07)

*For  $\omega$ -categorical model-complete cores, it is possible to add **constants** without changing the complexity.*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, '15)


*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*

Proposition (Bodirsky, '07)


*For  $\omega$ -categorical model-complete cores, it is possible to add constants without changing the complexity.*


Good news: we can choose the MMSNP structure  $\mathcal{B}$  so that  $(\mathcal{B}, \neq)$  is an  $\omega$ -categorical model-complete core.





- ▶ Obstructions  $\mathcal{F}$ : 
- ▶  $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :





- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :




- ▶ Obstructions  $\mathcal{F}$ : 
- ▶  $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :





- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :








- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



## Proposition

*There is a unif. cont. clonoid homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$  iff there is one  $\text{Pol}(\mathcal{B}, \bullet, \bullet) \rightarrow \mathcal{P}$ .*

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion



## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)

## Definition


$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)




Every canonical function determines a **behaviour** on colours.



Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.  
Only **finitely many** behaviours of a given arity.

Every canonical function determines a **behaviour** on colours.




We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

### Theorem (Bodirsky-M, '16)

*Let  $\mathcal{B}$  be in the class (reduct of...). If  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers operation modulo  $\overline{\text{Aut}(\mathcal{B})}$  that is canonical with respect to  $\text{Aut}(\mathcal{B})$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

### Theorem (Bodirsky-M, '16)

*Let  $\mathcal{B}$  be in the class (reduct of...). If  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers operation modulo  $\overline{\text{Aut}(\mathcal{B})}$  that is canonical with respect to  $\text{Aut}(\mathcal{B})$ , then  $\text{CSP}(\mathcal{B})$  is in P.*

$\Rightarrow$  If there is no clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in P.



Three cases:

1. No clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ : done (in P),
2. Clone homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ : done (NP-hard),
3. not case 1 or 2: ?

Three cases:

1. No clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ : done (in P),
2. Clone homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ : done (NP-hard),
3. not case 1 or 2: ?

## Proposition

*For  $\mathcal{B}$  an MMSNP structure, case 3 does not happen.*

Three cases:

1. No clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ : done (in P),
2. Clone homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ : done (NP-hard),
3. not case 1 or 2: ?

## Proposition

*For  $\mathcal{B}$  an MMSNP structure, case 3 does not happen.*

## Proof.

- Every function in  $\text{Pol}(\mathcal{B})$  has canonical friends in  $\text{Pol}(\mathcal{B})_{\text{can}}$ ,

Three cases:

1. No clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ : done (in P),
2. Clone homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ : done (NP-hard),
3. not case 1 or 2: ?

## Proposition

*For  $\mathcal{B}$  an MMSNP structure, case 3 does not happen.*

## Proof.

- ▶ Every function in  $\text{Pol}(\mathcal{B})$  has canonical friends in  $\text{Pol}(\mathcal{B})_{\text{can}}$ ,
- ▶ Some clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$  is constant on the sets of canonical friends,

Three cases:

1. No clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ : done (in P),
2. Clone homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ : done (NP-hard),
3. not case 1 or 2: ?

## Proposition

*For  $\mathcal{B}$  an MMSNP structure, case 3 does not happen.*

## Proof.

- ▶ Every function in  $\text{Pol}(\mathcal{B})$  has canonical friends in  $\text{Pol}(\mathcal{B})_{\text{can}}$ ,
- ▶ Some clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$  is constant on the sets of canonical friends,
- ▶ Define extension of  $\xi$  in natural way.



- ▶  $\sigma$ : set of colour symbols.
- ▶ A **trivial subfactor** of  $\mathcal{C}$  is a partition  $S \uplus T \subseteq \sigma$  such that  $\mathcal{C} / \sim$  is isomorphic to  $\mathcal{P}$ .

## Proposition

*Suppose there is  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ . There exists  $S, T$  trivial subfactor of  $\mathcal{C}$  and  $E$  undirected graph on  $\sigma$  s.t.:*

- ▶  *$E$  is preserved by  $\text{Pol}(\mathcal{B})$ ,*
- ▶  *$E$  contains an edge from  $S$  to  $T$  but does not contain **pseudo-loops**;*
- ▶ *there is no  $E$ -path of even length between  $S$  and  $T$ .*

$\rightsquigarrow$  clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$  that is constant on every set of canonical friends.

**Theorem (Hubička-Nešetřil, 2016)**

*Let  $\mathcal{B}$  be an MMSNP structure. Then there is a linear order  $<$  on  $B$  such that  $(\mathcal{B}, <)$  is  $\omega$ -categorical and Ramsey.*

**Theorem (Hubička-Nešetřil, 2016)**

*Let  $\mathcal{B}$  be an MMSNP structure. Then there is a linear order  $<$  on  $B$  such that  $(\mathcal{B}, <)$  is  $\omega$ -categorical and Ramsey.*

**Theorem (Bodirsky-Pinsker-Tsankov, 2010)**

*Suppose that  $\mathcal{G}$  is the automorphism group of an  $\omega$ -categorical ordered Ramsey structure. For every  $f: B^k \rightarrow B$ , there exists a function  $g \in \overline{\mathcal{G}f\mathcal{G}}$  that is canonical with respect to  $\mathcal{G}$ .*



Summing up:

## Theorem

*Let  $\mathcal{B}$  be an MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is  $NP$ -complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be an MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be an MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
- 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be an MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
  - 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
  - 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*
  - 4.  $\text{Pol}(\mathcal{B}, \bullet, \bullet)$  contains a colour-canonical pseudo-Siggers,*
- and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be an MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
- 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*
- 4.  $\text{Pol}(\mathcal{B}, \bullet, \bullet)$  contains a colour-canonical pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Items 3. and 4. are decidable.