

MMSNP: Towards a Proof of the Universal-Algebraic Dichotomy Conjecture

Manuel Bodirsky, **Antoine Mottet**

September 25, 2018

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity:

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

-
- ▶ Input: undirected graph G
 - ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: in P.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\quad)$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x})) \quad)$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

In general, for some fixed set \mathcal{F} of vertex-coloured graphs, the problem $\text{FPP}(\mathcal{F})$ is:

- ▶ Input: finite graph G ,
- ▶ Question: can one colour the vertices of G and **avoid** \mathcal{F} ?

(i.e., find G^* such that $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$)

MMSNP: formulas of the form


$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:


- ▶ M_1, \dots, M_n are **unary** predicates,
- ▶ α_i is a conjunction of **positive** literals $E(x_j, x_k)$,
- ▶ β_i is a conjunction of literals.

MMSNP and FPP are computationally equivalent.


Def: Structure is **clique-like** if its **Gaifman graph** is a clique.



Def: Structure is **clique-like** if its **Gaifman graph** is a clique.



Def: Structure is **clique-like** if its **Gaifman graph** is a clique.



For general structures: Gaifman graph is obtained by replacing tuples by cliques.

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden clique-like patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden clique-like patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden clique-like patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,
- ▶ Completely solve question posed by Lutz and Wolter (ICDT'15),

New proofs of known results:

- ▶ For every finite set \mathcal{F} of forbidden clique-like patterns, the corresponding problem is in P or NP-complete,
- ▶ Algorithm that takes \mathcal{F} and outputs complexity of $\text{FPP}(\mathcal{F})$.

New results:

- ▶ An algebraic characterisation of membership in P,
- ▶ Completely solve question posed by Lutz and Wolter (ICDT'15),
- ▶ Partial confirmation of conjecture by Bodirsky-M (LICS'16).

Introduction

MMSNP “ \subseteq ” CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Theorem and Example

Conclusion

Definition

$\mathfrak{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathfrak{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathfrak{A} ,
- ▶ **Question:** is there a homomorphism $\mathfrak{A} \rightarrow \mathfrak{B}$?

Definition

$\mathfrak{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathfrak{B})$ is the problem:

- ▶ **Input:** a **finite** graph \mathfrak{A} ,
- ▶ **Question:** is there a homomorphism $\mathfrak{A} \rightarrow \mathfrak{B}$?

Everything in this talk generalises to **relational** structures.

Definition

$\mathfrak{B} = (B; E)$ a **graph**.

$\text{CSP}(\mathfrak{B})$ is the problem:


- ▶ **Input:** a **finite** graph \mathfrak{A} ,
- ▶ **Question:** is there a homomorphism $\mathfrak{A} \rightarrow \mathfrak{B}$?


Everything in this talk generalises to **relational** structures.


In general, the forbidden patterns problem (FPP) for \mathcal{F} is **not a CSP**, but a **finite union** of CSPs.


Proposition


Every FPP reduces in polynomial-time to a finite number of FPP of connected structures.











Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$.

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$.

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

FPP of finite sets of finite **connected** structures

Theorem (Cherlin-Shelah-Shi, Adv. Appl. Math. 1999)

For every finite set \mathcal{F} of finite connected coloured graphs, there exists an ω -categorical partially coloured graph \mathfrak{B}^ such that $\mathfrak{A}^* \rightarrow \mathfrak{B}^*$ iff \mathfrak{A}^* avoids \mathcal{F} .*

Construct a new \mathfrak{B} by:

- ▶ deleting the uncoloured elements in \mathfrak{B}^* ,
- ▶ forgetting about the colours.

For this talk: we call \mathfrak{B} an **MMSNP structure**.

Proposition (Bodirsky-Dalmau, STACS'06)

$\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$.

Moreover, \mathfrak{B} belongs to the Bodirsky-Pinsker class.

FPP of finite sets of finite **connected** structures



CSPs in the BP class


Obstructions \mathcal{F} :



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :




Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Obstructions \mathcal{F} :



Structure \mathfrak{B} such that $\text{CSP}(\mathfrak{B}) = \text{FPP}(\mathcal{F})$:



Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Theorem and Example

Conclusion

For each \mathfrak{B} in the BP class,

complexity of $\text{CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B})$.

For each \mathfrak{B} in the BP class,

complexity of $\text{CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B})$.

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

For each \mathfrak{B} in the BP class,

complexity of $\text{CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B})$.

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

- ▶ Topology: put a metric on $\text{Pol}(\mathfrak{B})$ by putting two functions close together if they agree on a large part of B ,

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

- ▶ Topology: put a metric on $\text{Pol}(\mathfrak{B})$ by putting two functions close together if they agree on a large part of B ,
- ▶ Algebra: a **height 1 equation** is an equation of the form

$$f(\text{variables}) = g(\text{variables}).$$

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

- ▶ Topology: put a metric on $\text{Pol}(\mathfrak{B})$ by putting two functions close together if they agree on a large part of B ,
- ▶ Algebra: a **height 1 equation** is an equation of the form

$$f(\text{variables}) = g(\text{variables}).$$

Examples:

- ▶ Commutativity is height 1: $f(x, y) = f(y, x)$,

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

- ▶ Topology: put a metric on $\text{Pol}(\mathfrak{B})$ by putting two functions close together if they agree on a large part of B ,
- ▶ Algebra: a **height 1 equation** is an equation of the form

$$f(\text{variables}) = g(\text{variables}).$$

Examples:

- ▶ Commutativity is height 1: $f(x, y) = f(y, x)$,
- ▶ Cyclicity is height 1: $f(x_1, x_2, \dots, x_n) = f(x_2, \dots, x_n, x_1)$,

For each \mathfrak{B} in the BP class,

$$\text{complexity of CSP}(\mathfrak{B}) \leftrightarrow \text{Pol}(\mathfrak{B}).$$

Known: $\text{Pol}(\mathfrak{B})$ “small/trivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ NP-hard.

Conjectured: $\text{Pol}(\mathfrak{B})$ “big/nontrivial” $\Rightarrow \text{CSP}(\mathfrak{B})$ in P.

Triviality is a topological-algebraic property:

- ▶ Topology: put a metric on $\text{Pol}(\mathfrak{B})$ by putting two functions close together if they agree on a large part of B ,
- ▶ Algebra: a **height 1 equation** is an equation of the form

$$f(\text{variables}) = g(\text{variables}).$$

Examples:

- ▶ Commutativity is height 1: $f(x, y) = f(y, x)$,
- ▶ Cyclicity is height 1: $f(x_1, x_2, \dots, x_n) = f(x_2, \dots, x_n, x_1)$,
- ▶ Associativity is **not** height 1: $f(x, f(y, z)) = f(f(x, y), z)$.

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** $:\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0,1\}$.
By definition trivial ($\xi = \text{identity}$).

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0,1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0, 1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

$$f(x, y) = f(y, x)$$

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0, 1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

$$\begin{aligned} f(x, y) &= f(y, x) \\ \Rightarrow \xi(f)(x, y) &= \xi(f)(y, x) \end{aligned}$$

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0, 1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

$$\begin{aligned} f(x, y) &= f(y, x) \\ \Rightarrow \xi(f)(x, y) &= \xi(f)(y, x) \\ \Rightarrow pr_i(x, y) &= pr_i(y, x), \text{ for some } i \in \{1, 2\} \end{aligned}$$

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0, 1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

$$\begin{aligned} f(x, y) &= f(y, x) \\ \Rightarrow \xi(f)(x, y) &= \xi(f)(y, x) \\ \Rightarrow pr_i(x, y) &= pr_i(y, x), \text{ for some } i \in \{1, 2\} \\ \Rightarrow x &= y \end{aligned}$$

Definition (projective height 1 homomorphisms)

$\text{Pol}(\mathfrak{B})$ is **trivial** : $\Leftrightarrow \exists$ uniformly continuous map $\xi: \text{Pol}(\mathfrak{B}) \rightarrow \mathcal{P}$ that preserves height 1 equations.

- ▶ \mathcal{P} : set of all projection operations on $\{0,1\}$.
By definition trivial ($\xi = \text{identity}$).
- ▶ $\text{Pol}(\mathfrak{B})$ contains a commutative binary operation \Rightarrow nontrivial.

$$\begin{aligned} f(x, y) &= f(y, x) \\ \Rightarrow \xi(f)(x, y) &= \xi(f)(y, x) \\ \Rightarrow pr_i(x, y) &= pr_i(y, x), \text{ for some } i \in \{1, 2\} \\ \Rightarrow x &= y \end{aligned}$$

Theorem (Barto-Opršal-Pinsker, Israel Jour. Math. 2015)

If \mathfrak{B} is ω -categorical and $\text{Pol}(\mathfrak{B})$ is trivial, $\text{CSP}(\mathfrak{B})$ is NP-hard.

Conjecture (Bodirsky-Pinsker, 2011)

*Let \mathfrak{B} be a reduct of a finitely bounded homogeneous structure.
If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .*

Conjecture (Bodirsky-Pinsker, 2011)

*Let \mathfrak{B} be a reduct of a finitely bounded homogeneous structure.
If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .*

Relativise to MMSNP:

Conjecture

*Let \mathfrak{B} be ω -categorical such that $\text{CSP}(\mathfrak{B})$ is in MMSNP.
If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .*

Conjecture (Bodirsky-Pinsker, 2011)

Let \mathfrak{B} be a reduct of a finitely bounded homogeneous structure. If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .

Relativise to MMSNP:

Conjecture

Let \mathfrak{B} be ω -categorical such that $\text{CSP}(\mathfrak{B})$ is in MMSNP. If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .

Theorem (Bodirsky-M, 2017)

*Let \mathfrak{B} be ω -categorical such that $\text{CSP}(\mathfrak{B})$ is in MMSNP **with clique-like obstructions**. If $\text{Pol}(\mathfrak{B})$ is not trivial, then $\text{CSP}(\mathfrak{B})$ is in P .*

- ▶ Precolouring, i.e., allow partially coloured instances,

- ▶ Precolouring, i.e., allow partially coloured instances,
- ▶ $\text{Pol}(\mathfrak{B})$ not trivial implies existence of some particular function in $\text{Pol}(\mathfrak{B})$, called **pseudo-Siggers** (negative \Rightarrow positive),

- ▶ Precolouring, i.e., allow partially coloured instances,
- ▶ $\text{Pol}(\mathfrak{B})$ not trivial implies existence of some particular function in $\text{Pol}(\mathfrak{B})$, called **pseudo-Siggers** (negative \Rightarrow positive),
- ▶ make the pseudo-Siggers **canonical** (positive \Rightarrow positive++),

- ▶ Precolouring, i.e., allow partially coloured instances,
- ▶ $\text{Pol}(\mathfrak{B})$ not trivial implies existence of some particular function in $\text{Pol}(\mathfrak{B})$, called **pseudo-Siggers** (negative \Rightarrow positive),
- ▶ make the pseudo-Siggers **canonical** (positive \Rightarrow positive++),
- ▶ reduction to finite-domain CSPs (Bodirsky-M, LICS'16),

- ▶ Precolouring, i.e., allow partially coloured instances,
- ▶ $\text{Pol}(\mathfrak{B})$ not trivial implies existence of some particular function in $\text{Pol}(\mathfrak{B})$, called **pseudo-Siggers** (negative \Rightarrow positive),
- ▶ make the pseudo-Siggers **canonical** (positive \Rightarrow positive++),
- ▶ reduction to finite-domain CSPs (Bodirsky-M, LICS'16),
- ▶ finite-domain CSPs satisfy the conjecture (Bulatov-Zhuk, FOCS'17).

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Theorem and Example

Conclusion

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)


\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Rephrased: do $\text{CSP}(\mathfrak{B}, \bullet, \bullet)$ and $\text{CSP}(\mathfrak{B})$ have same complexity?



A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Proposition (Bodirsky, LMCS 2007)

*For some structures (ω -categorical model-complete cores), it is possible to add **constants** without changing the complexity.*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

Question (Lutz-Wolter, ICDT'15)

\mathcal{F} finite family of finite connected coloured graphs. Do $\text{FPP}(\mathcal{F})$ and its precoloured version have the same complexity?

Proposition (Bodirsky, LMCS 2007)

*For some structures (ω -categorical model-complete cores), it is possible to add **constants** without changing the complexity.*

Good news: we can choose the MMSNP structure \mathfrak{B} so that (\mathfrak{B}, \neq) is an ω -categorical model-complete core.

Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :




Suppose the vertex x is precoloured in the input:



Obstructions \mathcal{F} :



Suppose the vertex x is precoloured in the input:



Proposition

The input precoloured graph is colourable iff the graph obtained by adding the gadgets is colourable.

Theorem

Every FPP and its precoloured version have the same complexity.

Theorem

Every FPP and its precoloured version have the same complexity.

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{CSP}(\mathfrak{B})$ is in P

Theorem

Every FPP and its precoloured version have the same complexity.

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet) \text{ nontrivial} \quad (\text{precolouring})$

$\Rightarrow \text{CSP}(\mathfrak{B}) \text{ is in P}$

A 6-ary function $s: B^6 \rightarrow B$ is **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{B})}$** if $\exists e_1, e_2 \in \overline{\text{Aut}(\mathfrak{B})}$ s.t.

A 6-ary function $s: B^6 \rightarrow B$ is **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{B})}$** if $\exists e_1, e_2 \in \overline{\text{Aut}(\mathfrak{B})}$ s.t.

$$e_1(s(x, y, x, z, y, z)) = e_2(s(y, x, z, x, z, y))$$

for all $x, y, z \in B$.

A 6-ary function $s: B^6 \rightarrow B$ is **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{B})}$** if $\exists e_1, e_2 \in \overline{\text{Aut}(\mathfrak{B})}$ s.t.

$$e_1(s(x, y, x, z, y, z)) = e_2(s(y, x, z, x, z, y))$$

for all $x, y, z \in B$.

Theorem (Barto et al., LICS'16 + LICS'17)

*If \mathfrak{A} is a **model-complete core** in the BP class and $\text{Pol}(\mathfrak{A})$ is nontrivial, then $\text{Pol}(\mathfrak{A})$ contains a **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{A})}$** .*

A 6-ary function $s: B^6 \rightarrow B$ is **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{B})}$** if $\exists e_1, e_2 \in \overline{\text{Aut}(\mathfrak{B})}$ s.t.

$$e_1(s(x, y, x, z, y, z)) = e_2(s(y, x, z, x, z, y))$$

for all $x, y, z \in B$.

Theorem (Barto et al., LICS'16 + LICS'17)

If \mathfrak{A} is a **model-complete core** in the BP class and $\text{Pol}(\mathfrak{A})$ is nontrivial, then $\text{Pol}(\mathfrak{A})$ contains a **pseudo-Siggers modulo $\overline{\text{Aut}(\mathfrak{A})}$** .

To prove: if $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers, then $\text{CSP}(\mathfrak{B})$ is in P.

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Theorem and Example

Conclusion

Definition

$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

Definition

$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)

Definition


$f: B^k \rightarrow B$, a group \mathcal{G} acting on B . f is **canonical** (wrt \mathcal{G}) if for every finite subset $S \subseteq B$ of B and $\alpha_1, \dots, \alpha_k \in \mathcal{G}$, there exists $\beta \in \mathcal{G}$ such that $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$.

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)




Every canonical function determines a **behaviour** on colours.



Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.
Only **finitely many** behaviours of a given arity.

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

Theorem (Bodirsky-M, LICS'16)

*Let \mathfrak{B} be in the BP class. If $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers operation *modulo* $\overline{\text{Aut}(\mathfrak{A})}$ that is canonical with respect to \mathfrak{A} , then $\text{CSP}(\mathfrak{B})$ is in P.*

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ nontrivial

(precolouring)

$\Rightarrow \text{CSP}(\mathfrak{B})$ is in P

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ nontrivial

(precolouring)

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a pseudo-Siggers

(Barto et al.)

$\Rightarrow \text{CSP}(\mathfrak{B})$ is in P

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ nontrivial (precolouring)

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a pseudo-Siggers (Barto et al.)

$\text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a canonical pseudo-Siggers

$\Rightarrow \text{CSP}(\mathfrak{B})$ is in P (Bodirsky-M.)

$\text{Pol}(\mathfrak{B})$ nontrivial

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ nontrivial (precolouring)

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a pseudo-Siggers (Barto et al.)

$\Rightarrow \text{Pol}(\mathfrak{B}, \bullet, \bullet)$ contains a canonical pseudo-Siggers (?!?)

$\Rightarrow \text{CSP}(\mathfrak{B})$ is in P (Bodirsky-M.)

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is an expansion \mathfrak{C} of \mathfrak{B} that is ω -categorical, ordered, and Ramsey.

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is an expansion \mathfrak{C} of \mathfrak{B} that is ω -categorical, ordered, and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is an expansion \mathfrak{C} of \mathfrak{B} that is ω -categorical, ordered, and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .

Problems:

- ▶ g is not unique,
- ▶ if f is pseudo-Siggers, we might get a g that is not pseudo-Siggers.

Theorem (Hubička-Nešetřil, 2016)

Let \mathfrak{B} be an MMSNP structure. Then there is an expansion \mathfrak{C} of \mathfrak{B} that is ω -categorical, ordered, and Ramsey.

Theorem (Bodirsky-Pinsker-Tsankov, 2010)

Suppose that \mathcal{G} is the automorphism group of an ω -categorical ordered Ramsey structure. For every $f: B^k \rightarrow B$, there exists a function $g \in \overline{\mathcal{G}f\mathcal{G}}$ that is canonical with respect to \mathcal{G} .


Problems:


- ▶ g is not unique,
- ▶ if f is pseudo-Siggers, we might get a g that is not pseudo-Siggers.


(Partial) solution: Mashups!


$\text{Pol}(\mathfrak{B})$


A large, empty circle is centered on the right side of the slide. The label $\text{Pol}(\mathfrak{B})$ is positioned to the upper left of the circle's top edge.











Theorem (Bodirsky-M, LICS'16)

*If $\text{Pol}(\mathfrak{B})$ has the **mashup property**,*

$\text{Pol}(\mathfrak{B})$ has pseudo-Siggers $\Leftrightarrow \text{Pol}(\mathfrak{B})$ has a colour-canonical Siggers.

Theorem (Bodirsky-M, LICS'16)

*If $\text{Pol}(\mathfrak{B})$ has the **mashup property**,*

$\text{Pol}(\mathfrak{B})$ has pseudo-Siggers $\Leftrightarrow \text{Pol}(\mathfrak{B})$ has a colour-canonical Siggers.

What we have now: if Φ is an MMSNP sentence with clique-like obstructions, the corresponding $\text{Pol}(\mathfrak{B})$ has the mashup property.

Theorem (Bodirsky-M, LICS'16)

If $\text{Pol}(\mathfrak{B})$ has the *mashup property*,

$\text{Pol}(\mathfrak{B})$ has pseudo-Siggers $\Leftrightarrow \text{Pol}(\mathfrak{B})$ has a colour-canonical Siggers.

What we have now: if Φ is an MMSNP sentence with clique-like obstructions, the corresponding $\text{Pol}(\mathfrak{B})$ has the mashup property.

Conjecture

For *every* MMSNP structure, $\text{Pol}(\mathfrak{B})$ has the mashup property.

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Theorem and Example

Conclusion

Summing up:

Theorem

*Let \mathfrak{B} be an MMSNP structure with clique-like obstructions.
Then either the following equivalent statements hold:*

1. $\text{Pol}(\mathfrak{B})$ is nontrivial,

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

*Let \mathfrak{B} be an MMSNP structure with clique-like obstructions.
Then either the following equivalent statements hold:*

- 1. $\text{Pol}(\mathfrak{B})$ is nontrivial,*
- 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

*Let \mathfrak{B} be an MMSNP structure with clique-like obstructions.
Then either the following equivalent statements hold:*

- 1. $\text{Pol}(\mathfrak{B})$ is nontrivial,*
- 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*
- 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,*

and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.

Summing up:

Theorem

*Let \mathfrak{B} be an MMSN structure with clique-like obstructions.
Then either the following equivalent statements hold:*

- 1. $\text{Pol}(\mathfrak{B})$ is nontrivial,*
 - 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,*
 - 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,*
 - 4. $\text{Pol}(\mathfrak{B}, \textcolor{red}{\bullet}, \textcolor{teal}{\bullet})$ contains a colour-canonical pseudo-Siggers,*
- and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.*

Summing up:

Theorem

*Let \mathfrak{B} be an MMSNP structure with clique-like obstructions.
Then either the following equivalent statements hold:*

1. $\text{Pol}(\mathfrak{B})$ is nontrivial,
 2. $\text{Pol}(\mathfrak{B})$ contains a pseudo-Siggers,
 3. $\text{Pol}(\mathfrak{B})$ contains a canonical pseudo-Siggers,
 4. $\text{Pol}(\mathfrak{B}, \bullet, \circ)$ contains a colour-canonical pseudo-Siggers,
- and $\text{CSP}(\mathfrak{B})$ is in P , or $\text{CSP}(\mathfrak{B})$ is NP-complete.*

Items 3. and 4. can be checked effectively.

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: ?

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: ?

Theorem \Rightarrow it suffices to check for colour-canonical Siggers.

- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: ?

Theorem \Rightarrow it suffices to check for colour-canonical Siggers.

Two colours \Rightarrow it suffices to check for ternary cyclic operation.


- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: ?

Theorem \Rightarrow it suffices to check for colour-canonical Siggers.

Two colours \Rightarrow it suffices to check for ternary cyclic operation.




- ▶ Input: undirected graph G ,
- ▶ Question: can one colour the vertices of G in a way to avoid the following patterns:



- ▶ Complexity: ?

Theorem \Rightarrow it suffices to check for colour-canonical Siggers.

Two colours \Rightarrow it suffices to check for ternary cyclic operation.



No Siggers \Rightarrow the problem is NP-complete.

Introduction

MMSNP " \subseteq " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP


Canonical Functions

Theorem and Example

Conclusion

- ▶ Solve the general MMSNP case (natural test-case for the Bodirsky-Pinsker conjecture).
- ▶ When is $\text{CSP}(\mathfrak{B})$ in Datalog? Is it decidable? (Rewritability of MMSNP into Datalog programs, Lutz et al.)
- ▶ MMSNP_2 : instead of colouring vertices, we colour edges. It is **more expressive** than MMSNP, but it is open whether it has a complexity dichotomy (Lutz et al.).

Example: is it possible to colour the edges of an input graph and avoid:



Since last year:

- ▶ Result from last year “Reasoning with Discrete Time” (i.e., quantitative temporal reasoning) submitted to Journal of the ACM and in second round of reviewing,
- ▶ LICS’16 result (reduction infinite-domain CSPs to finite-domain + mashup technique) strengthened and submitted to LMCS (reports received yesterday).