

# MMSNP: An algebraic proof of the dichotomy

Manuel Bodirsky, Florent Madelaine, **Antoine Mottet**

September 25, 2018

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity:

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.

- ▶ Input: undirected graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity: NP-complete.
- 

- ▶ Input: undirected graph  $G$
- ▶ Question: can one colour the vertices of  $G$  in a way to avoid the following patterns:



- ▶ Complexity: in P.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSN: formulas of the form

$$\exists M_1 \cdots \exists M_n$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg($$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x})) \quad )$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ ,



In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ ,
- ▶  $\beta_i$  is a conjunction of literals.

In general, for some fixed set  $\mathcal{F}$  of vertex-coloured graphs, the problem  $\text{FPP}(\mathcal{F})$  is:

- ▶ Input: finite graph  $G$ ,
- ▶ Question: can one colour the vertices of  $G$  and **avoid**  $\mathcal{F}$ ?

(i.e., find  $G^*$  such that  $\forall F^* \in \mathcal{F}, F^* \not\rightarrow G^*$ )

MMSNP: formulas of the form

$$\exists M_1 \cdots \exists M_n \forall \bar{x} \bigwedge_i \neg(\alpha_i(\bar{x}) \wedge \beta_i(\bar{x}, M_1, \dots, M_n))$$

where:

- ▶  $M_1, \dots, M_n$  are **unary** predicates,
- ▶  $\alpha_i$  is a conjunction of **positive** literals  $E(x_j, x_k)$ ,
- ▶  $\beta_i$  is a conjunction of literals.

MMSNP and FPP are computationally equivalent.

Last monday:

Theorem (Bulatov, Zhuk '17)

*Finite-domain CSPs have a complexity dichotomy.*

Last monday:

Theorem (Bulatov, Zhuk '17)

*Finite-domain CSPs have a complexity dichotomy.*

A couple Mondays ago:

Theorem (Feder-Vardi '93, Kun '13)

*MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.*

Last monday:

Theorem (Bulatov, Zhuk '17)

*Finite-domain CSPs have a complexity dichotomy.*

A couple Mondays ago:

Theorem (Feder-Vardi '93, Kun '13)

*MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.*

The End.

Last monday:

Theorem (Bulatov, Zhuk '17)

*Finite-domain CSPs have a complexity dichotomy.*

A couple Mondays ago:

Theorem (Feder-Vardi '93, Kun '13)

*MMSNP has a complexity dichotomy if and only if finite-domain CSPs have a complexity dichotomy.*

# The End?

- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.

- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.

## Theorem

*Let  $\mathcal{A}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{A})$  is in MMSNP. Then one of the following holds:*

- ▶ *there is a uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{A}) \rightarrow \mathcal{P}$ , and  $\text{CSP}(\mathcal{A})$  is NP-complete,*
- ▶  *$\mathcal{A}$  has a pseudo-Siggers polymorphism and  $\text{CSP}(\mathcal{A})$  is in P.*



- ▶ Finite-domain dichotomy is algebraic, but Kun's reduction is not.
- ▶ MMSNP can describe some infinite-domain CSPs, and the conjecture for infinite domains is still open.

## Theorem

*Let  $\mathcal{A}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{A})$  is in MMSNP. Then one of the following holds:*

- ▶ *there is a uniformly continuous clonoid homomorphism  $\text{Pol}(\mathcal{A}) \rightarrow \mathcal{P}$ , and  $\text{CSP}(\mathcal{A})$  is NP-complete,*
- ▶  *$\mathcal{A}$  has a pseudo-Siggers polymorphism and  $\text{CSP}(\mathcal{A})$  is in P.*

In particular, this confirms the infinite-domain conjecture for CSPs in MMSNP.

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

## Definition

$\mathcal{B} = (B; E)$  a **graph**.

$\text{CSP}(\mathcal{B})$  is the problem:

- ▶ **Input:** a **finite** graph  $\mathcal{A}$ ,
- ▶ **Question:** is there a homomorphism  $\mathcal{A} \rightarrow \mathcal{B}$ ?

## Definition

$\mathcal{B} = (B; E)$  a **graph**.

$\text{CSP}(\mathcal{B})$  is the problem:

- ▶ **Input:** a **finite** graph  $\mathcal{A}$ ,
- ▶ **Question:** is there a homomorphism  $\mathcal{A} \rightarrow \mathcal{B}$ ?

In general, the forbidden patterns problem (FPP) for  $\mathcal{F}$  is **not a CSP**, but a **finite union** of CSPs.

## Proposition

*Every FPP reduces in polynomial-time to a finite number of FPP of connected structures.*

**Theorem (Cherlin-Shelah-Shi, '99)**

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

**Theorem (Cherlin-Shelah-Shi, '99)**

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

Construct a new  $\mathcal{B}$  by:

- ▶ deleting the uncoloured elements in  $\mathcal{B}^*$ ,
- ▶ forgetting about the colours.

For this talk: we call  $\mathcal{B}$  an **MMSNP structure**.

### Theorem (Cherlin-Shelah-Shi, '99)

*For every finite set  $\mathcal{F}$  of finite connected coloured graphs, there exists an  $\omega$ -categorical partially coloured graph  $\mathcal{B}^*$  such that  $\mathcal{A}^* \rightarrow \mathcal{B}^*$  iff  $\mathcal{A}^*$  avoids  $\mathcal{F}$ .*

Construct a new  $\mathcal{B}$  by:

- ▶ deleting the uncoloured elements in  $\mathcal{B}^*$ ,
- ▶ forgetting about the colours.

For this talk: we call  $\mathcal{B}$  an **MMSNP structure**.

### Proposition (Bodirsky-Dalmau, '06)

$$\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F}).$$

Moreover,  $\mathcal{B}$  belongs to the class of **reducts of finitely bounded homogeneous structures**.


Obstructions  $\mathcal{F}$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :



Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :






Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :




Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Obstructions  $\mathcal{F}$ :



Structure  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \text{FPP}(\mathcal{F})$ :



Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

**Conjecture (Bodirsky-Pinsker, '11 (rephrased))**

*Let  $\mathcal{B}$  be a reduct of a finitely bounded homogeneous structure.  
If there is no uniformly continuous clonoid homomorphism  
 $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

**Conjecture (Bodirsky-Pinsker, '11 (rephrased))**

*Let  $\mathcal{B}$  be a reduct of a finitely bounded homogeneous structure.  
If there is no uniformly continuous clonoid homomorphism  
 $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

**Theorem (Bodirsky-Madelaine-M, '18)**

*Let  $\mathcal{B}$  be  $\omega$ -categorical such that  $\text{CSP}(\mathcal{B})$  is in MMSNP.  
If there is no uniformly continuous clonoid homomorphism  
 $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.



A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)


*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)

$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?

Rephrased: do  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$  and  $\text{CSP}(\mathcal{B})$  have same complexity?



A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.

### Question (Lutz-Wolter, '15)

*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*

### Proposition (Bodirsky, '07)

*For  $\omega$ -categorical model-complete cores, it is possible to add **constants** without changing the complexity.*

A **precoloured** forbidden patterns problem is an FPP where the input can be partially coloured.


### Question (Lutz-Wolter, '15)

*$\mathcal{F}$  finite family of finite connected coloured graphs. Do  $\text{FPP}(\mathcal{F})$  and its precoloured version have the same complexity?*


### Proposition (Bodirsky, '07)


*For  $\omega$ -categorical model-complete cores, it is possible to add **constants** without changing the complexity.*


Good news: we can choose the MMSNP structure  $\mathcal{B}$  so that  $(\mathcal{B}, \neq)$  is an  $\omega$ -categorical model-complete core.


- ▶ Obstructions  $\mathcal{F}$ : 
- ▶  $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :





- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :




- ▶ Obstructions  $\mathcal{F}$ : 
- ▶  $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :




- ▶ Obstructions  $\mathcal{F}$ : 
- ▶  $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :











- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



- Obstructions  $\mathcal{F}$ : 
- $x$  precoloured in the instance of  $\text{CSP}(\mathcal{B}, \bullet, \bullet)$ :



## Proposition

*The input precoloured graph is colourable iff the graph obtained by adding the gadgets is colourable.*

Introduction

MMSNP " $\subseteq$ " CSP

A Dichotomy Conjecture for Infinite-Domain CSPs

Precoloured MMSNP

Canonical Functions

Conclusion

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)

## Definition


$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:

“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)




Every canonical function determines a **behaviour** on colours.






Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.  
Only **finitely many** behaviours of a given arity.

Every canonical function determines a **behaviour** on colours.




We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

### Theorem (Bodirsky-M, '16)

*Let  $\mathcal{B}$  be in the class (reduct of...). If  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers operation modulo  $\overline{\text{Aut}(\mathcal{B})}$  that is canonical with respect to  $\text{Aut}(\mathcal{B})$ , then  $\text{CSP}(\mathcal{B})$  is in  $P$ .*

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a finite set.

Only **finitely many** behaviours of a given arity.

### Theorem (Bodirsky-M, '16)

*Let  $\mathcal{B}$  be in the class (reduct of...). If  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers operation modulo  $\overline{\text{Aut}(\mathcal{B})}$  that is canonical with respect to  $\text{Aut}(\mathcal{B})$ , then  $\text{CSP}(\mathcal{B})$  is in P.*

$\Rightarrow$  If there is no clone homomorphism  $\text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ , then  $\text{CSP}(\mathcal{B})$  is in P.

1. We suppose that there is a clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ .

1. We suppose that there is a clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ .
2.  $\text{Pol}(\mathcal{B})_{\text{can}}$  is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of  $\text{Pol}(\mathcal{B})_{\text{can}}$  that is isomorphic to  $\mathcal{P}$ .

1. We suppose that there is a clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ .
2.  $\text{Pol}(\mathcal{B})_{\text{can}}$  is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of  $\text{Pol}(\mathcal{B})_{\text{can}}$  that is isomorphic to  $\mathcal{P}$ .
3. Using this subfactor and trickery, we show that the polymorphisms of  $\mathcal{B}$  **canonize in a unique way**,

1. We suppose that there is a clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ .
2.  $\text{Pol}(\mathcal{B})_{\text{can}}$  is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of  $\text{Pol}(\mathcal{B})_{\text{can}}$  that is isomorphic to  $\mathcal{P}$ .
3. Using this subfactor and trickery, we show that the polymorphisms of  $\mathcal{B}$  **canonize in a unique way**<sup>†</sup>,



1. We suppose that there is a clone homomorphism  $\xi: \text{Pol}(\mathcal{B})_{\text{can}} \rightarrow \mathcal{P}$ .
2.  $\text{Pol}(\mathcal{B})_{\text{can}}$  is an idempotent clone by the precolouring. This implies the existence of a **subfactor** of  $\text{Pol}(\mathcal{B})_{\text{can}}$  that is isomorphic to  $\mathcal{P}$ .
3. Using this subfactor and trickery, we show that the polymorphisms of  $\mathcal{B}$  **canonize in a unique way**<sup>‡</sup>,
4. Define a clonoid homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$  by **canonizing** and composing with  $\xi$ .

- ▶  $\sigma$ : set of colour symbols.
- ▶ A **trivial subfactor** of  $\mathcal{C}$  is a partition  $S \uplus T \subseteq \sigma$  such that  $\mathcal{C} / \sim$  is isomorphic to  $\mathcal{P}$ .

## Proposition

$S, T$  trivial subfactor of  $\mathcal{C}$ .  $\exists (B, E)$  undirected graph s.t.:

- ▶  $(B, E)$  contains an edge from  $S$  to  $T$  but does not contain *pseudo-loops*;
- ▶ the connected components of  $N$  are included in  $S$ , included in  $T$ , or bipartite with the bipartition induced by  $S$  and  $T$ ;
- ▶  $E$  is preserved by  $\text{Pol}(B)$ .

**Theorem (Hubička-Nešetřil, 2016)**

*Let  $\mathcal{B}$  be an MMSNP structure. Then there is a linear order  $<$  on  $B$  such that  $(\mathcal{B}, <)$  is  $\omega$ -categorical and Ramsey.*

**Theorem (Hubička-Nešetřil, 2016)**

*Let  $\mathcal{B}$  be an MMSNP structure. Then there is a linear order  $<$  on  $B$  such that  $(\mathcal{B}, <)$  is  $\omega$ -categorical and Ramsey.*

**Theorem (Bodirsky-Pinsker-Tsankov, 2010)**

*Suppose that  $\mathcal{G}$  is the automorphism group of an  $\omega$ -categorical ordered Ramsey structure. For every  $f: B^k \rightarrow B$ , there exists a function  $g \in \overline{\mathcal{G}f\mathcal{G}}$  that is canonical with respect to  $\mathcal{G}$ .*

### Theorem (Hubička-Nešetřil, 2016)

*Let  $\mathcal{B}$  be an MMSNP structure. Then there is a linear order  $<$  on  $B$  such that  $(\mathcal{B}, <)$  is  $\omega$ -categorical and Ramsey.*

### Theorem (Bodirsky-Pinsker-Tsankov, 2010)

*Suppose that  $\mathcal{G}$  is the automorphism group of an  $\omega$ -categorical ordered Ramsey structure. For every  $f: B^k \rightarrow B$ , there exists a function  $g \in \overline{\mathcal{G}f\mathcal{G}}$  that is canonical with respect to  $\mathcal{G}$ .*

Given  $f \in \text{Pol}(\mathcal{B})$ , and  $\xi: \mathcal{C} \rightarrow \mathcal{P}$  given by subfactor  $\{S, T\}$ , define  $\phi(f) := \xi(g)$  where  $g$  is an arbitrary function in  $\mathcal{C} \cap \overline{\text{Aut}(\mathcal{B}, <)f\text{Aut}(\mathcal{B}, <)}$ .

### Proposition

$\phi: \text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$  is a well-defined uniformly continuous height 1 homomorphism.

Summing up:

## Theorem

*Let  $\mathcal{B}$  be a MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be a MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be a MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
- 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*



Summing up:

## Theorem

*Let  $\mathcal{B}$  be a MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
  - 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
  - 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*
  - 4.  $\text{Pol}(\mathcal{B}, \bullet, \bullet)$  contains a colour-canonical pseudo-Siggers,*
- and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Summing up:

## Theorem

*Let  $\mathcal{B}$  be a MMSNP structure.*

*Then either the following equivalent statements hold:*

- 1. there is no uniformly continuous height 1 homomorphism  $\text{Pol}(\mathcal{B}) \rightarrow \mathcal{P}$ ,*
- 2.  $\text{Pol}(\mathcal{B})$  contains a pseudo-Siggers,*
- 3.  $\text{Pol}(\mathcal{B})$  contains a canonical pseudo-Siggers,*
- 4.  $\text{Pol}(\mathcal{B}, \bullet, \bullet)$  contains a colour-canonical pseudo-Siggers,*

*and  $\text{CSP}(\mathcal{B})$  is in  $P$ , or  $\text{CSP}(\mathcal{B})$  is NP-complete.*

Items 3. and 4. can be checked effectively.