

Constraint Satisfaction Problems over the Integers with Successor

Manuel Bodirsky, Barnaby Martin, **Antoine Mottet**

ICALP 2015

Let σ be a relational signature.

A σ -sentence ϕ is primitive positive if it is of the form

$$\exists x_1, \dots, x_n \bigwedge_{i=1}^m R_i(x_{i_1}, \dots, x_{i_k}), \quad R_i \in (\sigma \cup \{=\}).$$

Let σ be a relational signature.

A σ -sentence ϕ is primitive positive if it is of the form

$$\exists x_1, \dots, x_n \bigwedge_{i=1}^m R_i(x_{i_1}, \dots, x_{i_k}), \quad R_i \in (\sigma \cup \{=\}).$$

What is the complexity of deciding satisfiability of pp-formulas over \mathbb{Z} , depending on σ ?

Let σ be a relational signature.

A σ -sentence ϕ is primitive positive if it is of the form

$$\exists x_1, \dots, x_n \bigwedge_{i=1}^m R_i(x_{i_1}, \dots, x_{i_k}), \quad R_i \in (\sigma \cup \{=\}).$$

What is the complexity of deciding satisfiability of pp-formulas over \mathbb{Z} , depending on σ ?

Examples:

- ▶ $\{<, +\}$: in **P** (consequence of LP tractability and **scalability**)

Let σ be a relational signature.

A σ -sentence ϕ is primitive positive if it is of the form

$$\exists x_1, \dots, x_n \bigwedge_{i=1}^m R_i(x_{i_1}, \dots, x_{i_k}), \quad R_i \in (\sigma \cup \{=\}).$$

What is the complexity of deciding satisfiability of pp-formulas over \mathbb{Z} , depending on σ ?

Examples:

- ▶ $\{<, +\}$: in **P** (consequence of LP tractability and **scalability**)
- ▶ $\{<, +, 1\}$: **NP-complete** (integer linear programming)

Let σ be a relational signature.

A σ -sentence ϕ is primitive positive if it is of the form

$$\exists x_1, \dots, x_n \bigwedge_{i=1}^m R_i(x_{i_1}, \dots, x_{i_k}), \quad R_i \in (\sigma \cup \{=\}).$$

What is the complexity of deciding satisfiability of pp-formulas over \mathbb{Z} , depending on σ ?

Examples:

- ▶ $\{<, +\}$: in **P** (consequence of LP tractability and **scalability**)
- ▶ $\{<, +, 1\}$: **NP-complete** (integer linear programming)
- ▶ $\{+, \times\}$: **undecidable** (Hilbert's Tenth Problem)

Definition ($\text{CSP}(\Gamma)$)

Let Γ be a relational structure with a finite signature. The constraint satisfaction problem of Γ is the following decision problem:

INPUT: a primitive positive sentence ϕ in the language of Γ ,

QUESTION: is ϕ true in Γ ?

The structure Γ is called the template of $\text{CSP}(\Gamma)$.

The CSP of $(\mathbb{Z}; \text{succ})$:

The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .


The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .
- ▶ The input is true in $(\mathbb{Z}, \text{succ})$ iff there is a graph homomorphism from G to $(\mathbb{Z}; \text{succ})$.

The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .
- ▶ The input is true in $(\mathbb{Z}, \text{succ})$ iff there is a graph homomorphism from G to $(\mathbb{Z}; \text{succ})$.

$$\exists x_1, \dots, x_6 (x_2 = \text{succ}(x_1) \wedge x_4 = \text{succ}(x_2) \wedge \dots) \quad \cong$$




The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .
- ▶ The input is true in $(\mathbb{Z}, \text{succ})$ iff there is a graph homomorphism from G to $(\mathbb{Z}; \text{succ})$.

$$\exists x_1, \dots, x_6 (x_2 = \text{succ}(x_1) \wedge x_4 = \text{succ}(x_2) \wedge \dots) \cong$$

satisfying assignment




\cong homomorphism to $(\mathbb{Z}; \text{succ})$

The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .
- ▶ The input is true in $(\mathbb{Z}, \text{succ})$ iff there is a graph homomorphism from G to $(\mathbb{Z}; \text{succ})$.
- ▶ Complexity:

$$\exists x_1, \dots, x_6 (x_2 = \text{succ}(x_1) \wedge x_4 = \text{succ}(x_2) \wedge \dots) \cong$$

satisfying assignment




\cong homomorphism to $(\mathbb{Z}; \text{succ})$

The CSP of $(\mathbb{Z}; \text{succ})$:

- ▶ We view the input formula as a directed graph G .
- ▶ The input is true in $(\mathbb{Z}, \text{succ})$ iff there is a graph homomorphism from G to $(\mathbb{Z}; \text{succ})$.
- ▶ Complexity: in **P**.

$$\exists x_1, \dots, x_6 (x_2 = \text{succ}(x_1) \wedge x_4 = \text{succ}(x_2) \wedge \dots) \cong$$



satisfying assignment

\cong homomorphism to $(\mathbb{Z}; \text{succ})$

Definition

Let Γ be a relational structure and $R \subseteq \Gamma^n$ be a relation.

Definition

Let Γ be a relational structure and $R \subseteq \Gamma^n$ be a relation. We say that R is first-order definable in Γ if there exists a first-order formula $\phi(x_1, \dots, x_n)$ in the language of Γ such that

$$(a_1, \dots, a_n) \in R \Leftrightarrow \Gamma \models \phi(a_1, \dots, a_n).$$

Definition

Let Γ be a relational structure and $R \subseteq \Gamma^n$ be a relation. We say that R is first-order definable in Γ if there exists a first-order formula $\phi(x_1, \dots, x_n)$ in the language of Γ such that

$$(a_1, \dots, a_n) \in R \Leftrightarrow \Gamma \models \phi(a_1, \dots, a_n).$$

Examples:

- ▶ The unary $R = \{x \in \mathbb{Z} : x \text{ is even}\}$ is definable in $(\mathbb{Z}; +)$,

Definition

Let Γ be a relational structure and $R \subseteq \Gamma^n$ be a relation. We say that R is first-order definable in Γ if there exists a first-order formula $\phi(x_1, \dots, x_n)$ in the language of Γ such that

$$(a_1, \dots, a_n) \in R \Leftrightarrow \Gamma \models \phi(a_1, \dots, a_n).$$

Examples:

- ▶ The unary $R = \{x \in \mathbb{Z} : x \text{ is even}\}$ is definable in $(\mathbb{Z}; +)$,
- ▶ $\{(x, y) \in \mathbb{Z}^2 : x \text{ divides } y\}$ is definable in $(\mathbb{Z}; \times)$.

Definition

Let Γ be a relational structure and $R \subseteq \Gamma^n$ be a relation. We say that R is first-order definable in Γ if there exists a first-order formula $\phi(x_1, \dots, x_n)$ in the language of Γ such that

$$(a_1, \dots, a_n) \in R \Leftrightarrow \Gamma \models \phi(a_1, \dots, a_n).$$

Examples:

- ▶ The unary $R = \{x \in \mathbb{Z} : x \text{ is even}\}$ is definable in $(\mathbb{Z}; +)$,
- ▶ $\{(x, y) \in \mathbb{Z}^2 : x \text{ divides } y\}$ is definable in $(\mathbb{Z}; \times)$.

Definition

Let Γ, Δ be structures over the same domain. We say that Γ is a reduct of Δ when all the relations of Γ are (fo-)definable in Δ .

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in P,

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in P,
- ▶ Let R be the ternary relation

$$\{(a + 1, a, a), (a, a + 1, a), (a + 1, a + 1, a) \mid a \in \mathbb{Z}\}.$$

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in P,
- ▶ Let R be the ternary relation

$$\{(a + 1, a, a), (a, a + 1, a), (a + 1, a + 1, a) \mid a \in \mathbb{Z}\}.$$

$\text{CSP}(\mathbb{Z}; R)$:

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in \mathbf{P} ,
- ▶ Let R be the ternary relation

$$\{(a+1, a, a), (a, a+1, a), (a+1, a+1, a) \mid a \in \mathbb{Z}\}.$$

$\text{CSP}(\mathbb{Z}; R)$: in \mathbf{P} ,

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in P ,
- ▶ Let R be the ternary relation

$$\{(a + 1, a, a), (a, a + 1, a), (a + 1, a + 1, a) \mid a \in \mathbb{Z}\}.$$

$\text{CSP}(\mathbb{Z}; R)$: in P ,

- ▶ $\text{CSP}(\mathbb{Z}; |x - y| = 1, |x - y| = 5)$: **NP-complete**,

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in P,
- ▶ Let R be the ternary relation

$$\{(a + 1, a, a), (a, a + 1, a), (a + 1, a + 1, a) \mid a \in \mathbb{Z}\}.$$

$\text{CSP}(\mathbb{Z}; R)$: in P,

- ▶ $\text{CSP}(\mathbb{Z}; |x - y| = 1, |x - y| = 5)$: NP-complete,
- ▶ $\text{CSP}(\mathbb{Z}; \neq, y - x \in \{1, 2\})$: NP-complete.

Definition

A distance CSP is a CSP whose template is a reduct of $(\mathbb{Z}; \text{succ})$.

Examples:

- ▶ $\text{CSP}(\mathbb{Z}; \text{succ})$: in \mathbf{P} ,
- ▶ Let R be the ternary relation

$$\{(a + 1, a, a), (a, a + 1, a), (a + 1, a + 1, a) \mid a \in \mathbb{Z}\}.$$

$\text{CSP}(\mathbb{Z}; R)$: in \mathbf{P} ,

- ▶ $\text{CSP}(\mathbb{Z}; |x - y| = 1, |x - y| = 5)$: **NP-complete**,
- ▶ $\text{CSP}(\mathbb{Z}; \neq, y - x \in \{1, 2\})$: **NP-complete**.

Problem (Complexity classification project for $(\mathbb{Z}; \text{succ})$)

Give a complete classification of the complexity of distance CSPs.


A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

- ▶ $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.


A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

► $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.



A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .


- ▶ $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.



- ▶ $(\mathbb{Z}; x = y \vee u = v, \text{succ})$ is not.

A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

- ▶ $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.




- ▶ $(\mathbb{Z}; x = y \vee u = v, \text{succ})$ is not.

Theorem (Bodirsky, Dalmau, Martin, Pinsker '10)

Let Γ be a locally finite reduct of $(\mathbb{Z}; \text{succ})$. Then $\text{CSP}(\Gamma)$ is in P or NP -complete, or $\text{CSP}(\Gamma)$ is the CSP of a finite structure.

A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

- ▶ $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.



- ▶ $(\mathbb{Z}; x = y \vee u = v, \text{succ})$ is not.

Theorem (Bodirsky, Dalmau, Martin, Pinsker '10)


Let Γ be a locally finite reduct of $(\mathbb{Z}; \text{succ})$. Then $\text{CSP}(\Gamma)$ is in P or NP -complete, or $\text{CSP}(\Gamma)$ is the CSP of a finite structure.

Our result:

- ▶ **Complete** classification of the complexity of distance CSPs.

A reduct Γ of $(\mathbb{Z}; \text{succ})$ is locally finite if every $x \in \mathbb{Z}$ is contained in finitely many tuples of relations of Γ .

- ▶ $(\mathbb{Z}; |y - x| \leq 3)$ is locally finite.



- ▶ $(\mathbb{Z}; x = y \vee u = v, \text{succ})$ is not.

Theorem (Bodirsky, Dalmau, Martin, Pinsker '10)

Let Γ be a locally finite reduct of $(\mathbb{Z}; \text{succ})$. Then $\text{CSP}(\Gamma)$ is in P or NP -complete, or $\text{CSP}(\Gamma)$ is the CSP of a finite structure.

Our result:

- ▶ **Complete** classification of the complexity of distance CSPs.
- ▶ **Systematic approach** using **universal algebraic methods**.

Fact

Let Γ be a relational structure, and let R be a relation that has a **primitive positive definition** in Γ . Then $\text{CSP}(\Gamma)$ and $\text{CSP}(\Gamma, R)$ are polynomial-time equivalent.

Fact

Let Γ be a relational structure, and let R be a relation that has a **primitive positive definition** in Γ . Then $\text{CSP}(\Gamma)$ and $\text{CSP}(\Gamma, R)$ are polynomial-time equivalent.

How to determine if R is pp-definable in Γ ?

Fact

Let Γ be a relational structure, and let R be a relation that has a **primitive positive definition** in Γ . Then $\text{CSP}(\Gamma)$ and $\text{CSP}(\Gamma, R)$ are polynomial-time equivalent.

How to determine if R is pp-definable in Γ ?

Definition

Let $f: D^k \rightarrow D$ and let $R \subseteq D^n$ be a relation. We say that f preserves R iff

Fact

Let Γ be a relational structure, and let R be a relation that has a **primitive positive definition** in Γ . Then $\text{CSP}(\Gamma)$ and $\text{CSP}(\Gamma, R)$ are polynomial-time equivalent.

How to determine if R is pp-definable in Γ ?

Definition

Let $f: D^k \rightarrow D$ and let $R \subseteq D^n$ be a relation. We say that f preserves R iff

$$\begin{pmatrix} a_1^1 & a_2^1 & \cdots & a_n^1 \\ a_1^2 & a_2^2 & \cdots & a_n^2 \\ & \vdots & & \\ a_1^k & a_2^k & \cdots & a_n^k \end{pmatrix} \in R$$

Fact

Let Γ be a relational structure, and let R be a relation that has a **primitive positive definition** in Γ . Then $\text{CSP}(\Gamma)$ and $\text{CSP}(\Gamma, R)$ are polynomial-time equivalent.

How to determine if R is pp-definable in Γ ?

Definition

Let $f: D^k \rightarrow D$ and let $R \subseteq D^n$ be a relation. We say that f preserves R iff

$$\begin{array}{ccccccc} \left(\begin{array}{cccc} a_1^1 & a_2^1 & \cdots & a_n^1 \\ a_1^2 & a_2^2 & \cdots & a_n^2 \\ & \vdots & & \\ a_1^k & a_2^k & \cdots & a_n^k \end{array} \right) & \in R \\ \downarrow & \downarrow & & \downarrow & \\ (b_1 & b_2 & \cdots & b_n) & \in R \end{array}$$

A polymorphism of Γ is a function that preserves all the relations of Γ .

Lemma

*If Γ is a finite structure, a relation R is pp-definable in Γ **if and only if** R is preserved by all the polymorphisms of Γ .*

A polymorphism of Γ is a function that preserves all the relations of Γ .

Lemma

If Γ is a finite structure, a relation R is pp-definable in Γ if and only if R is preserved by all the polymorphisms of Γ .

- ▶ The previous lemma generalizes to infinite structures which have many automorphisms.

A polymorphism of Γ is a function that preserves all the relations of Γ .

Lemma

If Γ is a finite structure, a relation R is pp-definable in Γ if and only if R is preserved by all the polymorphisms of Γ .

- ▶ The previous lemma generalizes to infinite structures which have many **automorphisms**.
- ▶ In general, a reduct of $(\mathbb{Z}; \text{succ})$ does not satisfy this condition.

A polymorphism of Γ is a function that preserves all the relations of Γ .

Lemma

*If Γ is a finite structure, a relation R is pp-definable in Γ **if and only if** R is preserved by all the polymorphisms of Γ .*


- ▶ The previous lemma generalizes to infinite structures which have many **automorphisms**.
- ▶ In general, a reduct of $(\mathbb{Z}; \text{succ})$ does not satisfy this condition.
- ▶ Solution: we can recover a part of the connection if Γ has enough **elements**.

Definition

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. There exists a unique countable model of $\text{Th}(\Gamma)$ which contains every other countable model. We call this structure the ω -saturated model of Γ , denoted by $\omega.\Gamma$.

Definition

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. There exists a unique countable model of $\text{Th}(\Gamma)$ which contains every other countable model. We call this structure the ω -saturated model of Γ , denoted by $\omega.\Gamma$.



Definition

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. There exists a unique countable model of $\text{Th}(\Gamma)$ which contains every other countable model. We call this structure the ω -saturated model of Γ , denoted by $\omega.\Gamma$.

Γ and $\omega.\Gamma$ have the **same CSP**.

Definition

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. There exists a unique countable model of $\text{Th}(\Gamma)$ which contains every other countable model. We call this structure the ω -saturated model of Γ , denoted by $\omega.\Gamma$.

Γ and $\omega.\Gamma$ have the **same CSP**.

Lemma

*Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. Let R be a relation fo-definable in $(\omega.\mathbb{Z}; \text{succ})$ that consists of n orbits under $\text{Aut}(\omega.\Gamma)$. Then R is pp-definable in $\omega.\Gamma$ **if and only if** R is preserved by all the polymorphisms of arity n of $\omega.\Gamma$.*

Definition

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. There exists a unique countable model of $\text{Th}(\Gamma)$ which contains every other countable model. We call this structure the ω -saturated model of Γ , denoted by $\omega.\Gamma$.

Γ and $\omega.\Gamma$ have the **same CSP**.

Lemma

*Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$. Let R be a relation fo-definable in $(\omega.\mathbb{Z}; \text{succ})$ that consists of n orbits under $\text{Aut}(\omega.\Gamma)$. Then R is pp-definable in $\omega.\Gamma$ **if and only if** R is preserved by all the polymorphisms of arity n of $\omega.\Gamma$.*

Example: succ consists of 1 orbit under $\text{Aut}(\omega.\Gamma)$. By the lemma, succ is pp-definable in $\omega.\Gamma$ iff it is preserved by all the endomorphisms of $\omega.\Gamma$.

Theorem (Bodirsky, Martin, AM '15)

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$ with a finite signature. There exists a structure Δ with $\text{CSP}(\Gamma) = \text{CSP}(\Delta)$ and such that one of the following cases applies.

Theorem (Bodirsky, Martin, AM '15)

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$ with a finite signature. There exists a structure Δ with $\text{CSP}(\Gamma) = \text{CSP}(\Delta)$ and such that one of the following cases applies.

1. Δ is a finite structure. In this case, $\text{CSP}(\Gamma)$ is conjectured to be in P or NP -complete.

Theorem (Bodirsky, Martin, AM '15)

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$ with a finite signature. There exists a structure Δ with $\text{CSP}(\Gamma) = \text{CSP}(\Delta)$ and such that one of the following cases applies.

- 1. Δ is a finite structure. In this case, $\text{CSP}(\Gamma)$ is conjectured to be in P or NP -complete.*
- 2. Δ is a reduct of $(\mathbb{Z}; =)$. In this case, $\text{CSP}(\Gamma)$ is either in P or NP -complete (Bodirsky, Kára '08).*

Theorem (Bodirsky, Martin, AM '15)

Let Γ be a reduct of $(\mathbb{Z}; \text{succ})$ with a finite signature. There exists a structure Δ with $\text{CSP}(\Gamma) = \text{CSP}(\Delta)$ and such that one of the following cases applies.

- 1. Δ is a finite structure. In this case, $\text{CSP}(\Gamma)$ is conjectured to be in P or NP -complete.*
- 2. Δ is a reduct of $(\mathbb{Z}; =)$. In this case, $\text{CSP}(\Gamma)$ is either in P or NP -complete (Bodirsky, Kára '08).*
- 3. Δ is a reduct of $(\mathbb{Z}; \text{succ})$ whose endomorphisms are all isometries. In this case, $\text{CSP}(\Gamma)$ is in P or NP -complete. Moreover, the tractability of $\text{CSP}(\Gamma)$ is characterized by the existence of certain polymorphisms of finite arity.*

- ▶ We obtain a characterization of those reducts of $(\mathbb{Z}; \text{succ})$ that are not **homomorphically equivalent** to a finite structure or to a reduct of $(\mathbb{Z}; =)$.

- ▶ We obtain a characterization of those reducts of $(\mathbb{Z}; \text{succ})$ that are not **homomorphically equivalent** to a finite structure or to a reduct of $(\mathbb{Z}; =)$.
- ▶ This characterization tells us that the **endomorphisms** of $\omega.\Gamma$ have infinite range; and that there exists $t > 0$ such that for every endomorphism e of $\omega.\Gamma$ and all $x \in \omega.\mathbb{Z}$, we have

$$|e(x + t) - e(x)| \leq t.$$

- ▶ We obtain a characterization of those reducts of $(\mathbb{Z}; \text{succ})$ that are not **homomorphically equivalent** to a finite structure or to a reduct of $(\mathbb{Z}; =)$.
- ▶ This characterization tells us that the **endomorphisms** of $\omega.\Gamma$ have infinite range; and that there exists $t > 0$ such that for every endomorphism e of $\omega.\Gamma$ and all $x \in \omega.\mathbb{Z}$, we have

$$|e(x + t) - e(x)| \leq t.$$

- ▶ We inductively reduce t to 1 by replacing Γ , which finally gives us that all the endomorphisms of $\omega.\Gamma$ satisfy

$$|e(x + 1) - e(x)| = 1.$$

- ▶ Does the class of reducts of $(\mathbb{Z}; <)$ exhibit a P/NP-complete dichotomy?

- ▶ Does the class of reducts of $(\mathbb{Z}; <)$ exhibit a P/NP-complete dichotomy?
- ▶ More ambitious project: classify the complexity of reducts of $(\mathbb{Z}; <, +)$, i.e., reducts of **Presburger Arithmetic**.