

# Cyclic terms, CSP, MMSNP

Antoine Mottet  
(j.w. Manuel Bodirsky, Florent Madelaine)

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,
  - ▶ equations satisfied in  $\mathcal{P}$  are satisfied in every clone (aka boring equations).

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,
  - ▶ equations satisfied in  $\mathcal{P}$  are satisfied in every clone (aka boring equations).
- ▶ A clone  $\mathcal{A}$  can be:

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,
  - ▶ equations satisfied in  $\mathcal{P}$  are satisfied in every clone (aka boring equations).
- ▶ A clone  $\mathcal{A}$  can be:
  - ▶ the set of all term operations of a universal algebra **A** (always),

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,
  - ▶ equations satisfied in  $\mathcal{P}$  are satisfied in every clone (aka boring equations).
- ▶ A clone  $\mathcal{A}$  can be:
  - ▶ the set of all term operations of a universal algebra  $\mathbf{A}$  (always),
  - ▶ the set  $\text{Pol}(\mathcal{A})$  of **polymorphisms** of a relational structure  $\mathcal{A}$  (sometimes),

- ▶ A set  $\mathcal{A}$  of finitary operations on  $A$  is a **clone** if:
  - ▶ every **projection**  $\pi_i^n$  is in  $\mathcal{A}$ ,
  - ▶  $\mathcal{A}$  is closed under composition.
- ▶  $\mathcal{P}$ :
  - ▶ the smallest clone on  $\{0, 1\}$ , containing only projections,
  - ▶ equations satisfied in  $\mathcal{P}$  are satisfied in every clone (aka boring equations).
- ▶ A clone  $\mathcal{A}$  can be:
  - ▶ the set of all term operations of a universal algebra  $\mathbf{A}$  (always),
  - ▶ the set  $\text{Pol}(\mathcal{A})$  of **polymorphisms** of a relational structure  $\mathcal{A}$  (sometimes),
  - ▶ shown in a beautiful picture (never).



- $f: A^n \rightarrow A$  is **cyclic** if it satisfies

$$f(x_1, \dots, x_n) \approx f(x_2, \dots, x_n, x_1)$$

- $f: A^n \rightarrow A$  is **cyclic** if it satisfies

$$f(x_1, \dots, x_n) \approx f(x_2, \dots, x_n, x_1)$$

- **clonoid homomorphism**: a map  $\xi: \mathcal{A} \rightarrow \mathcal{B}$  such that

$$\xi(f \circ (\pi_{i_1}, \dots, \pi_{i_n})) = \xi(f) \circ (\pi_{i_1}, \dots, \pi_{i_n})$$

- $f: A^n \rightarrow A$  is **cyclic** if it satisfies

$$f(x_1, \dots, x_n) \approx f(x_2, \dots, x_n, x_1)$$

- **clonoid homomorphism**: a map  $\xi: \mathcal{A} \rightarrow \mathcal{B}$  such that

$$\xi(f \circ (\pi_{i_1}, \dots, \pi_{i_n})) = \xi(f) \circ (\pi_{i_1}, \dots, \pi_{i_n})$$

### Theorem (Barto-Kozik + Barto-Opršal-Pinsker)

Let  $\mathcal{A}$  be a **finite** relational structure. Then exactly one of the following holds:

- (+)  $\text{Pol}(\mathcal{A})$  contains a cyclic operation,
- (-) there exists a clonoid homomorphism  $\text{Pol}(\mathcal{A}) \rightarrow \mathcal{P}$ .



► Cyclic?

- ▶ Cyclic? Too strong for infinite structures:  
 $\mathcal{A}$  = clone generated by all injective operations on  $A$ .  
 $\mathcal{A}$  contains no cyclic operation and  $\mathcal{A} \not\rightarrow \mathcal{P}$ .

- Cyclic? Too strong for infinite structures:  
 $\mathcal{A}$  = clone generated by all injective operations on  $A$ .  
 $\mathcal{A}$  contains no cyclic operation and  $\mathcal{A} \not\rightarrow \mathcal{P}$ .

Pseudo-cyclic:

$$e_1 f(x_1, \dots, x_n) \approx e_2 f(x_2, \dots, x_n, x_1)$$

for unary  $e_1, e_2$ .

- ▶ Cyclic? Too strong for infinite structures:  
 $\mathcal{A}$  = clone generated by all injective operations on  $A$ .  
 $\mathcal{A}$  contains no cyclic operation and  $\mathcal{A} \nrightarrow \mathcal{P}$ .

Pseudo-cyclic:

$$e_1 f(x_1, \dots, x_n) \approx e_2 f(x_2, \dots, x_n, x_1)$$

for unary  $e_1, e_2$ .

- ▶ Clonoid homomorphisms: required to be **uniformly continuous**.



- **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  $\omega$ -categorical if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

- ▶ **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.
- ▶  $\text{Pol}(\mathbb{Q}; <, \neq)$ :

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  $\omega$ -categorical if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

- ▶ **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.
- ▶  $\text{Pol}(\mathbb{Q}; <, \neq)$ :
  - ▶ no pseudo-cyclic,

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  $\omega$ -categorical if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

- ▶ **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.
- ▶  $\text{Pol}(\mathbb{Q}; <, \neq)$ :
  - ▶ no pseudo-cyclic,
  - ▶ no uniformly continuous clonoid homomorphism to  $\mathcal{P}$  (because other non-boring equations are satisfied).

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  **$\omega$ -categorical** if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

- ▶ **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.
- ▶  $\text{Pol}(\mathbb{Q}; <, \neq)$ :
  - ▶ no pseudo-cyclic,
  - ▶ no uniformly continuous clonoid homomorphism to  $\mathcal{P}$  (because other non-boring equations are satisfied).
  - ▶ The structure is even  $\omega$ -categorical<sup>1</sup>, the next best thing after being finite.

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  $\omega$ -categorical if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

- ▶ **Goal:** find other structures  $\mathcal{A}$  for which the cyclic term theorem holds.
- ▶  $\text{Pol}(\mathbb{Q}; <, \neq)$ :
  - ▶ no pseudo-cyclic,
  - ▶ no uniformly continuous clonoid homomorphism to  $\mathcal{P}$  (because other non-boring equations are satisfied).
  - ▶ The structure is even  $\omega$ -categorical<sup>1</sup>, the next best thing after being finite.
- ▶ **Here:** structures  $\mathcal{B}$  for which the set  $\{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}$  has a nice logical description.

---

<sup>1</sup>A countable structure  $\mathcal{A}$  is  $\omega$ -categorical if  $\text{Aut}(\mathcal{A})$  acts with finitely many orbits on  $A^n$ , for all  $n \geq 1$ .

**Here:** structures  $\mathcal{B}$  for which the set  $\{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}$  has a nice logical description.

**Here:** structures  $\mathcal{B}$  for which the set  $\{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}$  has a nice logical description.

### Definition (CSP)

$\text{CSP}(\mathcal{B}) := \{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}.$

- ▶  $\text{CSP}(K_3) =$  all finite 3-colourable graphs,
- ▶  $\text{CSP}(\mathbb{Q}, <) =$  all finite directed acyclic graphs.



**Here:** structures  $\mathcal{B}$  for which the set  $\{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}$  has a nice logical description.

### Definition (CSP)

$\text{CSP}(\mathcal{B}) := \{\mathcal{A} \mid \mathcal{A} \text{ finite}, \mathcal{A} \rightarrow \mathcal{B}\}.$

- ▶  $\text{CSP}(K_3) =$  all finite 3-colourable graphs,
- ▶  $\text{CSP}(\mathbb{Q}, <) =$  all finite directed acyclic graphs.

Building structures with given CSPs:

### Theorem (Cherlin, Shelah, Shi)

*Let  $\mathfrak{F}$  be a finite set of finite connected graphs. There exists an  $\omega$ -categorical  $\mathcal{B}$  such that  $\mathcal{A} \rightarrow \mathcal{B}$  iff  $\forall \mathcal{F} \in \mathfrak{F}, \mathcal{F} \not\rightarrow \mathcal{A}$ .*

Example: there exists an  $\omega$ -categorical graph  $\mathcal{B}$  such that

$$\text{CSP}(\mathcal{B}) = \{\mathcal{A} \mid \mathcal{A} \text{ is finite and triangle-free}\}.$$

A fragment of **second-order logic** to describe CSPs:

A fragment of **second-order logic** to describe CSPs:

$$\begin{aligned} \exists \bullet, \circ \forall x, y, z & (\bullet(x) \vee \circ(x)) \wedge \neg(\bullet(x) \wedge \circ(x)) \\ & \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \bullet(x) \wedge \bullet(y) \wedge \bullet(z)) \\ & \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \circ(x) \wedge \circ(y) \wedge \circ(z)) \end{aligned}$$

A fragment of **second-order logic** to describe CSPs:

$$\begin{aligned} \exists \textcolor{red}{\bullet}, \textcolor{teal}{\bullet} \forall x, y, z (&\textcolor{red}{\bullet}(x) \vee \textcolor{teal}{\bullet}(x)) \wedge \neg(\textcolor{red}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(x)) \\ &\neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{red}{\bullet}(x) \wedge \textcolor{red}{\bullet}(y) \wedge \textcolor{red}{\bullet}(z)) \\ &\neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{teal}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(y) \wedge \textcolor{teal}{\bullet}(z)) \end{aligned}$$

- $\mathcal{A}$  satisfies  $\Phi_{\text{No-mono-tri}}$  iff there exists a **colouring**  $\mathcal{A}^*$  of  $\mathcal{A}$  such that  $\mathcal{A}^*$  contains no monochromatic triangle.

A fragment of **second-order logic** to describe CSPs:

$$\begin{aligned} \exists \textcolor{red}{\bullet}, \textcolor{teal}{\bullet} \forall x, y, z (\textcolor{red}{\bullet}(x) \vee \textcolor{teal}{\bullet}(x)) \wedge \neg(\textcolor{red}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(x)) \\ \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{red}{\bullet}(x) \wedge \textcolor{red}{\bullet}(y) \wedge \textcolor{red}{\bullet}(z)) \\ \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{teal}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(y) \wedge \textcolor{teal}{\bullet}(z)) \end{aligned}$$

- ▶  $\mathcal{A}$  satisfies  $\Phi_{\text{No-mono-tri}}$  iff there exists a **colouring**  $\mathcal{A}^*$  of  $\mathcal{A}$  such that  $\mathcal{A}^*$  contains no monochromatic triangle.
- ▶ Cherlin-Shelah-Shi: There exists an  $\omega$ -categorical graph  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \{\mathcal{A} \mid \mathcal{A} \models \Phi_{\text{No-mono-tri}}\}$ .

A fragment of **second-order logic** to describe CSPs:

$$\begin{aligned} \exists \textcolor{red}{\bullet}, \textcolor{teal}{\bullet} \forall x, y, z (\textcolor{red}{\bullet}(x) \vee \textcolor{teal}{\bullet}(x)) \wedge \neg(\textcolor{red}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(x)) \\ \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{red}{\bullet}(x) \wedge \textcolor{red}{\bullet}(y) \wedge \textcolor{red}{\bullet}(z)) \\ \neg(E(x, y) \wedge E(y, z) \wedge E(z, x) \wedge \textcolor{teal}{\bullet}(x) \wedge \textcolor{teal}{\bullet}(y) \wedge \textcolor{teal}{\bullet}(z)) \end{aligned}$$

- ▶  $\mathcal{A}$  satisfies  $\Phi_{\text{No-mono-tri}}$  iff there exists a **colouring**  $\mathcal{A}^*$  of  $\mathcal{A}$  such that  $\mathcal{A}^*$  contains no monochromatic triangle.
- ▶ Cherlin-Shelah-Shi: There exists an  $\omega$ -categorical graph  $\mathcal{B}$  such that  $\text{CSP}(\mathcal{B}) = \{\mathcal{A} \mid \mathcal{A} \models \Phi_{\text{No-mono-tri}}\}$ .
- ▶ More generally, we consider formulas with:
  - ▶ existential unary second-order quantifiers,
  - ▶ universal first-order quantifiers,
  - ▶ a conjunction of forbidden patterns.

## Theorem

Let  $\mathcal{B}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{B}) \in \text{MMSNP}$ . Then:

- (-) There is a *uniformly continuous* clonoid homomorphism from  $\text{Pol}(\mathcal{B})$  to  $\mathcal{P}$ , or
- (+)  $\text{Pol}(\mathcal{B})$  contains a *pseudo-cyclic* operation.

## Theorem

Let  $\mathcal{B}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{B}) \in \text{MMSNP}$ . Then:

- (-) There is a *uniformly continuous* clonoid homomorphism from  $\text{Pol}(\mathcal{B})$  to  $\mathcal{P}$ , or
- (+)  $\text{Pol}(\mathcal{B})$  contains a *pseudo-cyclic* operation.

A by-product of the proof gives the following corollary:

## Corollary

The logic MMSNP has a P/NP-complete dichotomy.



## Theorem

Let  $\mathcal{B}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{B}) \in \text{MMSNP}$ . Then:

- (-) There is a *uniformly continuous* clonoid homomorphism from  $\text{Pol}(\mathcal{B})$  to  $\mathcal{P}$ , or
- (+)  $\text{Pol}(\mathcal{B})$  contains a *pseudo-cyclic* operation.

A by-product of the proof gives the following corollary:

## Corollary

The logic MMSNP has a P/NP-complete dichotomy.

- Only focus on *particular* structures  $\mathcal{C}_\Phi$ , for *particular* MMSNP sentences  $\Phi$ .

## Theorem

Let  $\mathcal{B}$  be  $\omega$ -categorical and such that  $\text{CSP}(\mathcal{B}) \in \text{MMSNP}$ . Then:

- (-) There is a *uniformly continuous* clonoid homomorphism from  $\text{Pol}(\mathcal{B})$  to  $\mathcal{P}$ , or
- (+)  $\text{Pol}(\mathcal{B})$  contains a *pseudo-cyclic* operation.

A by-product of the proof gives the following corollary:

## Corollary

The logic MMSNP has a P/NP-complete dichotomy.

- ▶ Only focus on *particular* structures  $\mathcal{C}_\Phi$ , for *particular* MMSNP sentences  $\Phi$ .
- ▶ Make a bet: if  $\text{Pol}(\mathcal{C}_\Phi)$  has a cyclic operation, it has a very regular one.

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)

## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:

“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)




## Definition

$f: B^k \rightarrow B$ , a group  $\mathcal{G}$  acting on  $B$ .  $f$  is **canonical** (wrt  $\mathcal{G}$ ) if for every finite subset  $S \subseteq B$  of  $B$  and  $\alpha_1, \dots, \alpha_k \in \mathcal{G}$ , there exists  $\beta \in \mathcal{G}$  such that  $\beta \circ f|_S = f \circ (\alpha_1, \dots, \alpha_k)|_S$ .

In our case, we only care about the following consequence:


“the colour of the output only depends on the colours of the inputs” (**colour-canonical**)



Every canonical function determines a **behaviour** on colours.




Every canonical function determines a **behaviour** on colours.






Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a **finite** set.

Every canonical function determines a **behaviour** on colours.



We view this behaviour as an operation on a **finite** set.

### Theorem (Bodirsky-Pinsker-Pongrácz)

*Let  $\mathcal{C}$  be a clone consisting of canonical functions with respect to a homogeneous structure. Then:*

- (+)  $\mathcal{C}$  contains a cyclic operation, or*
- (-) there exists a **clone** homomorphism  $\mathcal{C} \rightarrow \mathcal{P}$ .*

**Theorem (Hubička-Nešetřil, 2016)**

*There is a linear order  $<$  on  $\mathcal{C}_\Phi$  such that  $\text{Aut}(\mathcal{C}_\Phi, <)$  is oligomorphic and extremely amenable.*


**Theorem (Hubička-Nešetřil, 2016)**

*There is a linear order  $<$  on  $\mathcal{C}_\Phi$  such that  $\text{Aut}(\mathcal{C}_\Phi, <)$  is oligomorphic and extremely amenable.*


**Theorem (Bodirsky-Pinsker-Tsankov, 2010)**

*Suppose that  $\mathcal{G} := \text{Aut}(\mathcal{B})$  is oligomorphic and extremely amenable. For every  $f: B^k \rightarrow B$ , there exists a function  $g \in \overline{\mathcal{G} f \mathcal{G}}$  that is canonical with respect to  $\mathcal{G}$ .*


- $\mathcal{C} \subseteq \text{Pol}(\mathcal{C}_\Phi)$ , so simply extend  $\mathcal{C} \rightarrow \mathcal{P}$  to  $\text{Pol}(\mathcal{C}_\Phi) \rightarrow \mathcal{P}$ !




- $\mathcal{C} \subseteq \text{Pol}(\mathcal{C}_\Phi)$ , so simply extend  $\mathcal{C} \rightarrow \mathcal{P}$  to  $\text{Pol}(\mathcal{C}_\Phi) \rightarrow \mathcal{P}$ !



- $\mathcal{C} \subseteq \text{Pol}(\mathcal{C}_\Phi)$ , so simply extend  $\mathcal{C} \rightarrow \mathcal{P}$  to  $\text{Pol}(\mathcal{C}_\Phi) \rightarrow \mathcal{P}$ !




- $\mathcal{C} \subseteq \text{Pol}(\mathcal{C}_\Phi)$ , so simply extend  $\mathcal{C} \rightarrow \mathcal{P}$  to  $\text{Pol}(\mathcal{C}_\Phi) \rightarrow \mathcal{P}$ !






- $\mathcal{C} \subseteq \text{Pol}(\mathcal{C}_\Phi)$ , so simply extend  $\mathcal{C} \rightarrow \mathcal{P}$  to  $\text{Pol}(\mathcal{C}_\Phi) \rightarrow \mathcal{P}$ !




►  $\Phi = \Phi_{\text{No-mono-tri}}$


►  $\phi = \phi_{\text{No-mono-tri}}$




- ▶  $\Phi = \Phi_{\text{No-mono-tri}}$
- ▶  $\mathcal{C} = \text{clone on } \{\text{red}, \text{teal}\}, \text{ no cyclic operation.}$



- ▶  $\Phi = \Phi_{\text{No-mono-tri}}$
- ▶  $\mathcal{C} = \text{clone on } \{\text{red}, \text{blue}\}$ , no cyclic operation.
- ▶ The relation  $N \subseteq (\mathcal{C}_\Phi)^2$  is invariant under  $\text{Pol}(\mathcal{C}_\Phi)$ .



- ▶  $\Phi = \Phi_{\text{No-mono-tri}}$
- ▶  $\mathcal{C} = \text{clone on } \{\text{red}, \text{blue}\}, \text{ no cyclic operation.}$
- ▶ The relation  $N \subseteq (\mathcal{C}_\Phi)^2$  is invariant under  $\text{Pol}(\mathcal{C}_\Phi)$ .
- ▶ So  $\text{Pol}(\mathcal{C}_\Phi) = \mathcal{C}$ !



- ▶ In general,  $\mathcal{C} \subsetneq \text{Pol}(\mathcal{C}_\Phi)$  and  $N$  is not invariant under  $\text{Pol}(\mathcal{C}_\Phi)$ .
- ▶ But some almost-bipartite simple graph is.
- ▶ Forces **canonizations of a single function** to have the same image under a clonoid homomorphism  $\mathcal{C} \rightarrow \mathcal{P}$ .

- ▶ In general,  $\mathcal{C} \subsetneq \text{Pol}(\mathcal{C}_\Phi)$  and  $N$  is not invariant under  $\text{Pol}(\mathcal{C}_\Phi)$ .
- ▶ But some almost-bipartite simple graph is.
- ▶ Forces **canonizations of a single function** to have the same image under a clonoid homomorphism  $\mathcal{C} \rightarrow \mathcal{P}$ .

## Theorem

$\Phi$  MMSNP formula. One of the following holds:

- (-) There is a **uniformly continuous** clonoid homomorphism from  $\text{Pol}(\mathcal{C}_\Phi)$  to  $\mathcal{P}$ , or
- (+)  $\text{Pol}(\mathcal{C}_\Phi)$  contains a **pseudo-cyclic** operation.