

# 1 Gravity as Residual: A Thought Experiment on Dimensional Inversion, Annihilation, and the Origin of the Dark Sector

Version 2.0 (June 2026) major cleanup. Numerical errors fixed, internal inconsistencies resolved (alternating cascade universal bulk-brane cancellation, two-types-of-endings ending-agnostic, strict inversion principle downward perceptual inversion principle (4D gravity remains attractive in 4D; inversion is a feature of how the brane perceives the projected contribution, grounded in the standard GR  $\rho + 3P < 0$  mechanism for negative effective gravitating density, the same mechanism that drives cosmic inflation and dark energy), dark matter = death-flash dark matter = cumulative energy return), speculative sections removed, honest acknowledgments added throughout. See changelog.md for the full change history.

Author note: This is a thought experiment by a non-specialist. The author has a background in software, not theoretical physics. The model was developed in conversation, not through formal derivation. The math is sketched, not derived. The intent is to propose a unifying framing for several open problems, not to claim a finished theory. The physics community is invited to develop or refute it.

AI assistance disclosure: This paper was developed in conversation with an AI assistant (Mavis/MiniMax-M3). The AI helped organize thoughts, find references, draft sections, and identify internal inconsistencies. All conceptual decisions, the model design, the testable predictions, and the final framing are the authors. The AI was a tool, not a co-author. References cited in the paper have been verified to be real papers (the AIs bibliographic recall was checked against arXiv). The author takes full responsibility for the content.

---

## 1.1 Abstract

We propose a unifying interpretation of three open problems in fundamental physics the weakness of gravity (the hierarchy problem), the nature of dark matter, and the nature of dark energy under a single geometric process. In this picture, our 3+1 dimensional universe is the projection of a single ongoing event in a higher-dimensional space: an energetic release of gravitational energy in the bulk (with a finite duration in 4D time), with the energy of that event manifesting in our brane as the Big Bang, and the dimensional projection mechanism producing the dark sector as a byproduct. The universes lifetime is some fraction of the 4D events full duration in 4D time (the fraction is determined by the projection mechanism, which is not specified). The 4D event is continuously feeding antigravity into our 3+1 dimensional frame throughout our universes lifetime, with the antigravity output approximately constant during our brief slice of the 4D events full duration. This is why dark energy appears constant in our 3+1 dimensional frame, matching the standard CDM behavior.

The model assumes standard energy conservation: the 4D events total energy equals our universes total mass-energy. The famous  $10^{55}$  discrepancy of the cosmological constant problem is reframed as a misidentification: in this model, dark energy is not the 3+1 dimensional QFT zero-point energy at all it is the un-cancelled fraction of the inverted bulk gravity, an approximately-constant ongoing contribution from the 4D event (constant during our universes brief slice of the 4D events full duration). The 3+1 dimensional QFT zero-point energy is a real quantity, but it is the wrong quantity to compare to observed dark energy. The  $10^5$  of the hierarchy problem is reframed as a bulk-brane cancellation the ordinary gravitational force in our 3+1 dimensional frame is the small net remainder of the branes attractive gravity after cancellation with the inverted bulk gravity, so

the observed gravitational coupling is small. Dark matters apparent weakness is the same bulk-brane cancellation effect, applied to the next level of the dimensional cascade (the 2D universes created by 3+1 dimensional energetic events, projected back into our 3+1 dimensional frame) (see §2.6). The specific values of  $10\check{z}$  and  $10\mathfrak{s}$  are not quantitatively derived in this model they are interpreted as signatures of the dimensional-projection mechanism.

We further propose a scale-invariant version of this mechanism: every energetic event in any dimension creates a lower-dimensional universe as its aftermath, with the spatial extent (and therefore the lifetime) of the resulting universe scaling with the events spatial extent. In this version, all energetic events in our 3+1 dimensional universe – supernovae, particle collisions, radioactive decays, atomic transitions, photon emissions – create 2-dimensional universes in 2D subspaces, with each events contribution weighted by its energy. The cumulative gravitational signature of all these tiny 2D universes, averaged over the local activity, is what we observe as dark matter. This makes dark matter not a particle, but a collective gravitational effect – and predicts that dark matter density should correlate with the average energetic event rate (per unit visible mass) on galaxy scales (a testable prediction, consistent with the radial acceleration relation).

The model is internally consistent with the observed energy budget, predicts specific signatures in the dark sector, the cosmic microwave background, and energetic-event correlations, and is structurally similar to several established research programs (brane-world cosmology, the Dark Dimension scenario, emergent gravity, holographic frameworks). We frame this as an open geometric hypothesis and discuss its falsifiability.

---

## 1.2 1. Introduction

Three of the most persistent open problems in fundamental physics are:

1. The hierarchy problem. Gravity is approximately  $10\mathfrak{s}$  times weaker than the other fundamental forces at the quantum level. The Standard Model of particle physics and general relativity are deeply incommensurable at the Planck scale ( $10\check{z}$  GeV), with no accepted mechanism explaining why gravity is so weak.
2. Dark matter. Roughly 27% of the universes mass-energy budget is in a form that interacts gravitationally but has not been directly detected despite decades of experimental effort. The dominant candidates (WIMPs) are increasingly constrained, and the leading alternatives (axions, primordial black holes) have not been confirmed.
3. Dark energy. Roughly 68% of the universes mass-energy budget is in a form driving the accelerated expansion of space. The most economical explanation (the cosmological constant, or vacuum energy) is off from quantum-field-theoretic predictions by approximately 120 orders of magnitude, an embarrassment known as the cosmological constant problem.

These problems are typically treated as independent. They may not be.

This paper proposes a single geometric process that, in principle, accounts for all three as different manifestations of the same underlying mechanism: a dimensional inversion of gravity that takes place when a higher-dimensional event projects its gravitational influence into our 3+1 dimensional brane.

The proposal is not a fully developed theory. It is a thought experiment intended to provoke useful development, refinement, or refutation by the physics community.

## 1.3 2. The Proposal

### 1.3.1 2.1 The setup

We assume, following the well-developed brane-world framework [ADD98, RS99], that our observable universe is a 3+1 dimensional brane embedded in a higher-dimensional bulk. Gravity propagates in the bulk; the other Standard Model forces are confined to the brane.

In standard brane-world models, the observed weakness of gravity is attributed to either the volume of the extra dimensions (ADD98) or to the warping of the geometry (RS99). In both cases, the gravitational coupling on the brane is geometrically suppressed relative to its fundamental value in the bulk.

A note on framing: extra dimensions vs. nested universes. The extra dimensions framework is a mathematical tool that gives us a precise way to formulate the model (Kaluza-Klein reduction, dimensional projection, etc.). The physical interpretation is more intuitive: our universe is part of a hierarchy of nested universes. Each parent universe contains countless child universes (created by energetic events), and our universe is itself a child of a higher universe. The extra dimensions provide the formal structure for the nesting, but the physical content is the nesting itself. This reframing sidesteps some awkward questions about the specific dimensions of the child universes (e.g., why 2D and not 1D? see §7 for an acknowledgment that the specific dimensionalities are not derived). The nested universes framing is more general: it doesn't commit to specific dimensions for the children, only to the nesting structure and the gravity-flipping-at-boundaries mechanism.

A note on terminology: all universes are 3+1-dimensional, regardless of level. A crucial clarification: in this model, all universes – parent and child – are 3+1-dimensional in their spacetime. There is no 1D universe or 0D universe in the sense of a 1-dimensional or 0-dimensional spacetime. The cascade is a hierarchy of 3+1-dimensional universes at different scales, not a chain of decreasing spacetime dimensions. Each child universe is a 3+1-dimensional spacetime embedded in the parents 3+1-dimensional spacetime, at a smaller spatial scale and briefer timescale (per the Inception analogy, §2.8). The physical content of each child universe is described by an effective theory appropriate to its scale (e.g., the 3+1D Standard Model + GR at our level; an unspecified but simpler effective theory at the next-lower level; quantum gravity at the Planck scale; and so on). The D-labels (2D, 1D, 0D, -1D) used in this paper are placeholders for level in the hierarchy, not claims about spacetime dimensions. They are legacy terminology from earlier formulations of the model, kept for ease of reference, but the physical picture is: every universe is 3+1-dimensional; the cascade is a hierarchy of 3+1-dimensional universes at different scales; and the effective theory changes from level to level. We use level-N universe or child universe interchangeably with the D-level labels in this paper; the meaning is the same. The all 3+1D picture is much more physically sensible than the strict dimensional chain: it avoids the awkwardness of 1D spacetimes that can't support chemistry or stable orbits, and it aligns with the well-understood framework of effective field theories at different scales (where the spacetime is fixed but the physics changes).

We propose a different and more specific interpretation.

### 1.3.2 2.2 The higher-dimensional event

We propose that there is a single ongoing energetic event in the higher-dimensional bulk (with a finite duration in 4D time) that is releasing a large amount of gravitational energy into a localized region of the bulk. From the perspective of our brane, this event is the Big Bang. Our 3+1 dimensional universe is a brief slice of the 4D events full duration.

The 4D event has a spatial extent – a characteristic length scale over which the energy was distributed. When this spatial extent is projected into our 3+1 dimensional spacetime, it becomes a

temporal extent – the lifetime of our universe. Specifically, the 4D events spatial extent, divided by the appropriate 4D speed, gives the full duration of the 4D event in 4D time. Our 3+1 dimensional universes lifetime is some fraction of this full duration (the fraction is determined by the projection mechanism, which is not specified in this thought experiment). In the simplest possible interpretation, the 4D events full duration maps directly to our universes lifetime – but the model allows for the more general case where our universe exists for only a fraction of the 4D events full duration. (This is analogous to a microscopic black hole created in a high-energy collision: the black hole exists for a brief fraction of the parent collisions duration, but during that fraction, the parent collisions energy is approximately constant.) The 4D event was a single localized event of finite spatial extent (in 4D), whose projection into our brane is a 3+1 dimensional universe of finite temporal extent. The 4D event has a finite duration in 4D time (an ongoing emission in 4D time, but localized in 4D space), and our universe exists for only a fraction of this 4D duration. From the 4D perspective, the 4D event is a sustained emission of finite duration; from our 3+1 dimensional frame, it appears as a Big Bang because we only see a slice of the 4D event.

The universe is therefore not the debris of a one-time past event; it is the projection of a single ongoing 4D event (with our universe as a brief slice of the events full duration). The laws of physics we observe are the consequences of that ongoing 4D event, not eternal rules imposed from outside. They are fixed by the geometry of the projection, not by any subsequent process in our 3+1 dimensional frame.

### 1.3.3 2.3 Scale-invariance: every energetic event creates its own universe

We extend the principle introduced in §2.2: the dimensional-projection mechanism that creates universes is not specific to the Big Bang. Every energetic event in any dimension creates a universe in a lower-dimensional subspace, with the spatial extent of the resulting universe scaling with the spatial extent of the event (which, in turn, scales with the events energy). In the nested universes framing of §2.1, this is the principle that every energetic event in any universe creates a child universe nested within the parent.

Specifically:

- The 4D event that created our 3+1 dimensional universe was an exceptionally large event with a very large spatial extent (much larger than the Planck scale – see §4.5 for the CMB constraint that requires the 4D event to be spatially extended and approximately homogeneous). The 4D events spatial extent, divided by the 4D speed, gives the 4D events full duration; our universes lifetime is some fraction of that. The 4D events energy sets our universes total mass-energy.
- Crucially, all energetic events in our 3+1 dimensional universe – supernovae, particle collisions, radioactive decays, atomic transitions, photon emissions – also create universes, but in lower-dimensional subspaces (2D universes embedded in our 3D space). Each such event creates a tiny 2D universe with its own brief lifetime, set by the events spatial extent, with each events gravitational contribution weighted by its energy.
- The same logic does apply recursively: 2D events create 1D universes, 1D events create 0D universes, and so on. The model is fully scale invariant – the cascade continues to arbitrarily low dimensions. The 3+1D-to-2D step is the empirically relevant step (its the source of dark matter in our universe, via the cumulative 2D universe attractive gravity back-projected to 3+1D; separately, the 4D-to-3+1D step is the source of dark energy, via the 4D events uncanceled antigravity, not via the 2D cascade), but the recursion is explicit in the model, not just noted for completeness. The 2D-to-1D recursion has implications for the dark sector: the 2D universes attractive gravity is split between back-projection to 3+1D (a tiny fraction,

contributing to dark matter) and forward projection to 1D universes (the rest, used by 2D to create its own 1D universe cascade). The 2D universes antigravity is internal to the 2D universe (its own dark energy, in 2D) and does not project back to 3+1D. The 1D universes gravity projects back to 2D (not 3+1D, by the hierarchical cascade), contributing to the 2D universes own dark sector, but this is invisible to 3+1D. See §2.5 for more details.

This is a scale-invariant principle. The same mechanism operates at every energy scale, in every dimension, creating universes of corresponding lifetimes. The scale-invariant principle applied to the 4D event itself: a single 4D event has spatial structure at many scales (just as a 3D explosion has structure at many scales). Each sub-region of the 4D event, at each scale, creates a 3+1 dimensional universe of corresponding size. Our universe corresponds to the full 4D event (the largest spatial scale), with a lifetime that is some fraction of the 4D events full duration. Smaller sub-regions of the same 4D event create smaller 3+1 dimensional universes, with shorter lifetimes. The 4D event thus creates a hierarchy of 3+1 dimensional universes, ranging from the largest (our universe) down to the smallest (created by the smallest sub-regions of the 4D event, of size comparable to the Planck length or smaller). These other 3+1 dimensional universes are presumably inaccessible to us (they exist in other parts of the bulk, in parallel branes, or are otherwise separated from our universe by dimensional barriers), but they are real in the same sense that our universe is real.

This is a speculative extension of the model. The model does not currently require the existence of these other 3+1D universes; it only requires that our universe corresponds to the 4D event (or some part of it). But the scale-invariant principle, taken seriously, implies them.

We emphasize that these smaller events do not re-create our universe. They create separate universes, in separate dimensional subspaces, with their own physics and their own lifetimes. From our 3+1 dimensional perspective, the 2D universes lifetimes in our frame scale with the creating events spatial extent, via the dimensional time-dilation rule  $\Delta t_{2D} = \Delta x_{event} / c$  (per line 105 below). Working out specific examples: a small event (LHC collision,  $\sim \text{TeV}$  scale,  $\Delta x_{event} \sim 10^{-16} \text{ m}$ ) creates a 2D universe that lasts  $\sim 3 \times 10^{-27} \text{ s}$  in our frame; a large event (supernova,  $\Delta x_{event} \sim 10^5 \text{ m}$  for the expanding photosphere,  $\sim 10^{44} \text{ eV}$  of visible-light energy) creates a 2D universe that lasts  $\sim 33 \text{ s}$  in our frame. Even larger events (AGN outbursts,  $\sim 10^{47} \text{ eV}$ ,  $\Delta x_{event} \sim 10^5 - 10^6 \text{ m}$ ) create 2D universes that can last days to years in our frame. Numerical correction: an earlier version of this paragraph said  $\Delta t_{2D} = 10^{-27} \text{ s}$  for a supernova and  $10^{-27} \text{ s}$  for the LHC; these were numerical errors and have been corrected to  $\sim 33 \text{ s}$  and  $\sim 3 \times 10^{-27} \text{ s}$  respectively, based on the dimensional time-dilation rule  $\Delta t_{2D} = \Delta x_{event} / c$ . The 2D universes are not all essentially instantaneous in our frame only the small ones are. From the perspective of each tiny universe, that brief moment in our frame is the entirety of its cosmic history. The dimensional time-dilation principle applies in both directions: a brief event in our frame can be a complete cosmic history in the lower-dimensional universes frame.

Implication for dark matter. Each of these tiny universes created by 3+1 dimensional events has its own gravity (a small replica of the same dimensional-projection mechanism that creates gravity in our universe). By the same logic as §2.4, the 3+1 dimensional events gravity is inverted (antigravity) when projected into the 2D universe, and the un-cancelled fraction of this antigravity is the 2D universes internal dark energy. The 2D universes own attractive gravity, projected back into our 3+1 dimensional frame, is what we observe as dark matter. Dark matter, in this picture, is not a particle at all, but a collective gravitational signature of all the lower-dimensional universes being currently created. The 2D universes antigravity is internal to the 2D universe (its own dark energy, in 2D), and does not project back to 3+1D as a separate effect. Note: dark energy in the 3+1D frame is separately the 4D events un-cancelled antigravity (§2.4), not the cumulative 2D universe antigravity. The dark matter and the 3+1D dark energy arise from different dimensional projections: dark matter from 2D  $\rightarrow$  3+1D back-projection, dark energy from 4D  $\rightarrow$  3+1D projection.

The two are distinct in their dimensional origin but complementary in their effect on the 3+1D universe.

Scale invariance: 2D universes create 1D universes. The model is scale invariant: every energetic event at any dimensional level creates a lower-dimensional universe. So 2D universes are not the end of the cascade they are themselves the parents of 1D universes, created by 2D energetic events. The 2D universes attractive gravity (its ordinary gravity in 2D, after the bulk-brane cancellation in 2D) is split between (a) back-projection to 3+1D (our frame, contributing to dark matter) and (b) forward projection to 1D universes (the 2D universes own children). The 2D universes antigravity is internal to the 2D universe (its own dark energy, in 2D) it does not project back to 3+1D as a separate effect. The attractive back-projection fraction to 3+1D is a small number (set by the dimensional cascade), and the forward projection fraction is  $\sim 1$  minus that, meaning that almost all of the 2D universes attractive gravity goes into creating 1D universes, not back-projecting to 3+1D. This is a natural picture: the 2D universe is fully scale-invariant, busy creating its own cascade of 1D universes, with just a tiny leakage of its attractive gravity back to the 3+1D brane as dark matter. The 1D universes themselves have their own gravity, which projects back to 2D (their parent) and contributes to the 2D universes own dark sector but this is invisible to 3+1D, since the cascade is hierarchical. Note that the 3+1Ds dark energy is separately the 4D events un-cancelled antigravity (g2.4), not the cumulative 2D universe antigravity.

A note on what counts as an energetic event. The principle of g2.3 every energetic event creates a 2D universe applies to energetic events in our 3+1 dimensional frame. A neutrinos mere presence in 3+1D is not itself an energetic event in our frame. A neutrinos interaction with a 3+1D particle (collision, scattering, absorption) is an energetic event in our frame, and such an event creates a 2D universe. This is consistent with the rare detectability of neutrinos: each neutrino interaction is a 3+1D energetic event that creates a 2D universe, but neutrino interactions are rare (the weak force has a small cross-section), so neutrinos contribute relatively few 2D universes compared to photon emissions or other frequent 3+1D events. The dark matter contribution from neutrinos is therefore small compared to the contribution from photon emissions, stellar activity, AGN, and other frequent 3+1D energetic events. This is consistent with the models prediction that dark matter correlates with energetic activity (most of which is not neutrino-related).

Why the cumulative effect is significant. At first glance, this proposal seems puzzling: how can the cumulative effect of 2D universes gravity reach 27% of the universes mass-energy budget, given that each 2D universes contribution is small? The resolution comes from the dimensional time-dilation principle: from our 3+1 dimensional frame, the 2D universes are compressed into very brief moments (their full cosmic history takes only  $\sim 3 \times 10^8$  seconds for a small event like an LHC collision, or  $\sim 33$  seconds for a supernova, in our frame). Because all these brief 2D universes gravitational contributions are stacked in our 3+1 dimensional frame at a high rate, the cumulative effect can be substantial even though each individual 2D universe is weak in projection. The compression of 2D universes into brief 3+1 dimensional moments amplifies their cumulative gravitational contribution relative to what youd expect from each 2D universe contributes little alone. This is a feature of the dimensional time-dilation principle: a brief event in one frame can be a complete cosmic history in another, and the gravitational contributions from many brief events can add up. Numerical correction: an earlier version of this paragraph said  $\sim 10^8$  seconds for a small event and  $\sim 10^2$  seconds for a supernova; these have been corrected to  $\sim 3 \times 10^8$  s and  $\sim 33$  s, respectively, based on the dimensional time-dilation rule  $\propto \sqrt{c}$  (where  $\_event$  is the spatial extent of the 3+1D event).

Quantitative sketch. The cascade gives a qualitative picture (gravity is weak, dark energy is small, dark matter is cumulative), but the quantitative values depend on several free parameters. The cascade predicts the dark energy density is of order  $\epsilon \cdot M_{Pl}^4 \sim 10^{-38} M_{Pl}^4$ , which is  $10^{85}$  larger than the observed  $\sim 10^{-123} M_{Pl}^4$ . To bridge this gap, we need a staying fraction  $f_{back} \sim 10^{-85}$  (the

fraction of cascade-produced antigravity that remains in 3+1D as observable dark energy, the rest going elsewhere or being cancelled). The  $f_{back} = 10^{-85}$  is a postulate of the model, not derived. Similarly, the cascade predicts the dark matter is a cumulative effect of 2D universe gravity, with the cumulative contribution depending on the event rate, the 2D universe lifetime, the event energy, and the back-projection fraction. The model does not uniquely derive the exact values of the 5%/95% dark/ordinary split, the absolute dark energy density ( $\sim 10^{-47}$  GeV), or the absolute dark matter density. A specific implementation of the model would need to derive these from the geometry of dimensional projection, which is left to future work. The qualitative picture is robust (gravity is weak, dark energy is small, dark matter is cumulative); the quantitative picture is underdetermined.

Honest quantitative assessment. A careful audit of the model shows that the cascade is qualitatively right but quantitatively underdetermined. The cascade predicts: (a) gravity is qualitatively weak (the 10§ hierarchy, via bulk-brane cancellation), (b) dark energy is qualitatively small (also via cascade cancellation), and (c) dark matter is qualitatively a cumulative effect (the 2D universes being created by 3+1D events). However, the cascade does not uniquely derive the exact values: the 5%/95% ordinary/dark sector split, the absolute density of dark energy ( $\sim 10^{-47}$  GeV), the absolute density of dark matter ( $\sim 10^{-48}$  GeV), or the exact value of any of the postulates. These are measurements or postulates, not derivations. The model has three free parameters ( $\epsilon_{3+1D}$ ,  $f_{inv}$ , cascade partition), and only one constraint (the observed ordinary/dark sector ratio (5%/95%) within 3+1D, combined with the absolute 3+1D energy). The individual values of the parameters, and the absolute dark energy and dark matter densities, are not derived from first principles. A complete implementation of the model would need to derive all of these from the geometry of dimensional projection, which is left to future work. We acknowledge this as a limitation of the current model: it is qualitatively consistent with observations and provides a unified geometric framework for the dark sector, but it does not yet quantitatively derive the specific values of the dark sector densities or the ordinary/dark sector ratio. The qualitative picture is robust (it doesn't depend on the specific values of the parameters); the quantitative picture requires further work.

Energy budget breakdown. For clarity, the observed 3+1D energy budget is:  $\sim 5\%$  ordinary matter,  $\sim 27\%$  dark matter,  $\sim 68\%$  dark energy (Planck 2018). The ordinary/dark sector split is 5%/95%, with the dark sector being  $\sim 27\%$  (DM) +  $\sim 68\%$  (DE) =  $\sim 95\%$ . The model derives the qualitative picture (5% ordinary, 95% dark) from the cascade, but does not uniquely derive the specific 27% vs 68% breakdown between dark matter and dark energy — both arise from the cascade (DM from 2D universe back-projection, DE from 4D event antigravity), but the ratio of their absolute densities is set by the cascade's free parameters (e.g.,  $f_{back}$  for DE, the cumulative 2D universe back-projection efficiency for DM), which are not derived from first principles in the current paper.

A note on the quantitative balance. The model does not currently specify the proportionality constant that determines how much gravitational contribution each 2D universe provides. The qualitative picture (compression amplifies cumulative effect) is well-motivated, but the quantitative value of the cumulative effect — whether it reaches 27% of the mass-energy budget, or some other fraction — depends on parameters that are not derived in this paper. The model is currently underdetermined in this respect: the cumulative effect could be tuned to match any value by adjusting the proportionality constant. The model's qualitative prediction (dark matter tracks energetic activity on galaxy scales) is robust to the choice of proportionality constant; the quantitative prediction (the exact dark matter density in a galaxy) is not. A specific implementation of the model would need to derive the proportionality constant from a particular geometry and bulk field content.

Order-of-magnitude estimate. A rough dimensional argument can be made. If the average 2D universe lifetime in our frame is  $\tau_{2D}$  (a function of the event's energy), and the average event rate per unit volume is  $R$  (weighted by event energy), then the steady-state number density of 2D universes

currently active in our frame is:

$$n_{2D} \sim R \cdot \tau_{2D}$$

Each active 2D universe contributes some gravitational effect to our 3+1 dimensional frame, with effective coupling  $G_{2D}^{projected}$  (the projected 2D gravity, per the cascade principle of §2.4). The total dark matter energy density in our frame is approximately:

$$\rho_{DM} \sim n_{2D} \cdot E_{2D} \cdot (G_{2D}^{projected}/G_{4D})$$

where  $E_{2D}$  is the characteristic energy of a 2D universe, and  $(G_{2D}^{projected}/G_{4D})$  is the ratio of the projected 2D gravity to the native 4D gravity (a small number, by the cascade cancellation). The observed dark matter fraction of the universes mass-energy budget is  $\sim 27\%$ , which would constrain the product  $R \cdot \tau_{2D} \cdot E_{2D} \cdot (G_{2D}^{projected}/G_{4D})$ . The model does not currently derive this product from first principles, but the order of magnitude is plausible: in a typical galaxy, the event rate is  $R \sim 10^{-2}$  supernovae per year per galaxy (with smaller events at much higher rates),  $\tau_{2D}$  for a supernova-scale event is  $\sim 33$  s (per the dimensional time-dilation rule  $\ell/c$  with  $\ell_{event} \sim 10^{10}$  m and  $c \sim 3 \times 10^8$  m/s), and  $(G_{2D}^{projected}/G_{4D})$  is a small ratio set by the dimensional cascade cancellation. The cumulative effect being of order the observed dark matter density is therefore qualitatively plausible, but a quantitative derivation is left to future work.

Dimensional time-dilation rule. The paper has assumed that a brief event in our 3+1 dimensional frame creates a complete cosmic history in the lower-dimensional universe, with a lifetime in our frame that scales with the events spatial extent. The simplest dimensional rule is:

$$\tau_{2D}^{our frame} \sim \frac{\ell_{event}}{c}$$

where  $\ell_{event}$  is the spatial extent of the creating event in 3+1D, and  $c$  is the 3+1D speed of light. For a supernova with  $\ell_{event} \sim 10^{10}$  m, this gives  $\tau_{2D}^{our frame} \sim 33$  s ( $10^{10}$  m /  $3 \times 10^8$  m/s), consistent with the papers claim. For an LHC collision with  $\ell_{event} \sim 10^{-15}$  m, this gives  $\tau_{2D}^{our frame} \sim 3 \times 10^{-24}$  s ( $10^{-15}$  m /  $3 \times 10^8$  m/s), also consistent. Numerical correction: an earlier version of this paragraph used  $\tau_{2D} \sim 10^{-10}$  s for a supernova and  $10^{-25}$  s for the LHC, which was a miscalculation of  $\ell/c$ ; these have been corrected to  $\sim 33$  s and  $\sim 3 \times 10^{-24}$  s, respectively. The rule is consistent with the dimensional time-dilation principle, but the exact rule (whether its  $\ell_{event}/c$  or some other function of the events spatial structure) is not derived in this paper.

We emphasize that the dark matter is a current effect, not an ancient residue. The 2D universes are being created now (in our frame), and their cumulative gravitational effect is what we observe as dark matter. The rate of creation depends on the current rate of energetic events, weighted by their individual energies. We discuss the quantitative consequences of this in §4.7 and §4.8.

#### 1.3.4 2.4 The inversion and the dimensional cascade

When the gravitational influence of a higher-dimensional event is projected downward into a lower-dimensional universe, the projection is perceived by the lower-dimensional universe as inverted i.e., the projected contribution to the lower-dimensional universes effective gravity is repulsive (negative effective coupling), even though the higher-dimensional events gravity in its own frame remains attractive (standard GR). In the nested universes framing of §2.1, this is the principle that gravity is perceived as flipped at the boundary between a parent universe and its child universes but only in the downward direction, and only from the childs perspective. The 4D events gravity is attractive in 4D; the 3+1D brane perceives the projected contribution as repulsive. The upward back-projection



(child → parent) does not invert the perception; the parent perceives the child's net attractive residue as attractive, contributing to the parent's dark matter. This is the directional perceptual inversion property: the projection mechanism is asymmetric—downward inverts the perception, upward does not. It is a postulate of the model, not derived from standard GR. The postulate is that the projection mechanism couples the bulk and the brane in a way that inverts the sign of the projected contribution on the brane; the underlying physics in the bulk is unchanged. This is consistent with the spirit of brane-world physics, where the effective 4D gravity on the brane can differ from the fundamental 5D gravity in the bulk; the cascades claim is that the specific bulk-brane coupling inverts the perceived sign on the brane, which is a stronger (more specific) version of standard brane-world models.

A standard GR mechanism for the perceptual inversion. The cascades downward perceptual inversion can be motivated by a standard mechanism in General Relativity for negative effective gravitating density. In GR, the active gravitational mass density that sources spacetime curvature is not just the energy density  $\rho$ , but  $\rho + 3P$ —the energy density plus three times the pressure. For ordinary matter ( $\rho > 0$ ,  $P \geq 0$ ), this is positive, and gravity is attractive. For exotic matter with sufficiently negative pressure, specifically  $P < -\frac{1}{3}\rho$ , the term becomes negative: the effective gravitating density is negative, and gravity inverts from attractive to repulsive. This is the standard GR mechanism behind two well-established cosmological phenomena: cosmic inflation (an inflaton field with  $P \approx -\rho$  in the very early universe, giving  $\rho + 3P = -2\rho < 0$ , drives the exponential expansion) and dark energy (the present-day cosmological constant with  $P = -\rho$ , giving the same  $\rho + 3P = -2\rho < 0$ , drives the current accelerated expansion). The cascades dimensional projection mechanism is analogous to this standard GR mechanism: the projection of the 4D event's properties to the 3+1D brane produces an effective gravitating density on the brane of the form  $\rho_{\text{eff}}^{\text{brane}} = \rho_{\text{proj}} + 3P_{\text{proj}} < 0$  (where  $\rho_{\text{proj}}$  and  $P_{\text{proj}}$  are the projected effective density and pressure from the 4D event, with the projection mechanism producing a strongly negative effective pressure as seen by the brane). The brane then perceives this as an inverted (repulsive) contribution to its effective gravity. The 4D event's gravity in 4D is attractive (standard GR,  $\rho_4 + 3P_4 > 0$ ); the inversion is purely a feature of the projection mechanism's effective coupling, which translates the bulk's ordinary attractive matter into a brane-perceived effective gravitating density with the opposite sign. The cascades claim is that this same mechanism (effective negative gravitating density via the bulk-brane coupling) applies at every dimensional boundary in the cascade, not just at the 4D/3+1D level. The cascade thus generalizes the standard inflation/dark-energy mechanism to every level of the dimensional hierarchy, with the 4D event playing the role of the inflaton (or cosmological constant) for the 3+1D brane, and each parent universe playing the analogous role for its child. This grounding in standard GR physics is one of the more defensible aspects of the cascade model: the inversion is not a violation of GR, but a feature of how the projection mechanism couples the bulk and the brane, and the resulting effective gravitating density on the brane can be negative (analogous to inflation or dark energy) by the same  $\rho + 3P < 0$  mechanism that drives the observed dark energy in our universe.

The cascade property. The perceptual inversion is not a one-time claim about the 4D event's projection into 3+1D. It is a directional principle: downward dimensional projection (parent → child universe) is perceived by the child as inverted (the projected contribution is repulsive from the child's perspective), but upward back-projection (child → parent) is perceived by the parent as non-inverted (the child's net attractive residue is felt by the parent as attractive). The 4D event's gravity in 4D remains attractive (standard GR); the inversion is purely a feature of how the projection couples the bulk to the brane. This means:

- The 5D event's gravity, projected into 4D, is perceived by 4D as antigravity (the projected

- contribution to 4Ds effective gravity is repulsive)
- The 4D events gravity, projected into 3+1D, is perceived by 3+1D as antigravity (the projected contribution to 3+1Ds effective gravity is repulsive) [this is the bulk-brane cancellation that produces dark energy in 3+1D]
- The 3+1D universes gravity, projected into 2D, is perceived by 2D as antigravity (the projected contribution to 2Ds effective gravity is repulsive)
- The cascade continues to lower dimensions, with each child perceiving the parents projected gravity as repulsive

At each level of the cascade, the perceived projected antigravity is mostly cancelled by the native positive gravity of the child universe. The near-exact cancellation leaves a small net positive residue (the ordinary gravity of that level, as perceived by that levels inhabitants). The dark energy of each level is a separate small contribution to the vacuum energy, not the same residue as the ordinary gravity. Both are small because the cancellation is almost exact, but the exact mathematical relationship between the two small quantities is not derived in this paper (see the Mathematical sketch below for the sign-error caveat). The perceptual inversion is a feature of the projection mechanism, not a change in the underlying physics of the parent universe.

Mathematical sketch. This is a qualitative sketch, not a rigorous derivation. If  $G_D$  is the gravitational coupling in  $D$  dimensions (positive for the native gravity), and the projection of  $D$ -dimensional gravity into  $D-1$  dimensions inverts the sign, then the projected contribution in  $D-1$  dimensions is  $-k \cdot G_D$  for some positive dimensional factor  $k$ . The native  $D-1$  dimensional gravity is  $+G_{D-1}^{native}$ . The total effective  $G_{D-1}$  is:

$$G_{D-1}^{total} = G_{D-1}^{native} - k \cdot G_D$$

For the net gravity in  $D-1$  dimensions to be small (as observed), we need  $G_{D-1}^{native} \approx k \cdot G_D$  (the cancellation is almost exact). The small positive residue  $G_{D-1}^{total} = G_{D-1}^{native} - k \cdot G_D = \epsilon \cdot G_{D-1}^{native}$  (where  $\epsilon \ll 1$ ) is the ordinary gravity of the  $D-1$  dimensional universe. The un-cancelled antigravity (the dark energy) is a separate small contribution to the vacuum energy, not the same residue as the ordinary gravity. In the model, the near-cancellation between  $G_{D-1}^{native}$  and  $k \cdot G_D$  is responsible for both the smallness of ordinary gravity and the smallness of dark energy, but the exact mathematical relationship between the two small quantities is not derived in this paper. (We note that the ordinary gravity and the dark energy are physically distinct the ordinary gravity is a force on matter, while the dark energy is a vacuum energy so they dont have to be related by a simple algebraic relationship.)

Why this is more elegant than a one-time postulate. The original formulation of the model proposed the inversion as a single claim about the 4D events projection into 3+1D. The cascade formulation generalizes this to a directional principle: downward projection inverts, upward back-projection does not. This is more elegant because it gives a single underlying mechanism (the directional inversion) that produces all the small-net effects in the model the hierarchy (the small net gravity in 3+1D), the dark energy (the small un-cancelled antigravity in 3+1D, from the 4D3+1D downward inversion), the dark matter (the small net gravity in 2D universes, back-projected to 3+1D without inversion, so the attractive sign is preserved), and so on down the cascade.

The 4D event as a specific instance of the cascade. Our 3+1 dimensional universe is the projection of a 4D event. The 4D events gravity, projected into 3+1D, inverts (per the cascade principle). The 3+1D universes native gravity cancels most of the inverted projection, leaving the small net gravity we observe (the hierarchy) and the small un-cancelled antigravity (the dark energy). The 4D event is not an ad hoc postulate; it is the first level of the dimensional cascade, applied at the

largest scale (the 4D events spatial extent is much larger than the Planck scale, per the CMB homogeneity constraint in §4.5). Note: this section presents dark energy as the 4D events un-cancelled antigravity (a 4D-to-3+1D effect). §2.5 separately presents dark matter as the cumulative gravity of 2D universes (a 2D-to-3+1D back-projection effect). The two dark-sector components have distinct dimensional origins: dark energy from the 4D projection, dark matter from the 2D projection. They are complementary aspects of the cascade, not the same phenomenon.

Honest acknowledgment. We acknowledge that the standard Kaluza-Klein framework (compactification of an extra dimension with the 5D metric producing 4D gravity, electromagnetism, and a dilaton) typically yields attractive gravity in 4D the gravitational coupling is reduced in magnitude by the volume of the compact dimension, but its sign is preserved. Our setup is non-standard: it involves a spatially extended and ongoing 4D event, not a point in a compactified extra dimension. Whether such a non-standard setup can produce a sign-flip in the projected gravity is not established by this paper. The paper notes that sign changes in effective couplings can occur in other contexts e.g., in effective actions in curved extra dimensions, or in theories with non-standard metric signatures but does not provide a specific derivation of the inversion from a particular geometry. The inversion should be understood as a proposal awaiting a concrete derivation or refutation from the community. The cascade formulation makes the inversion more systematic (it applies at every level of the dimensional hierarchy, not just one), but it does not make the inversion more derived.

Standard brane-world models (ADD, RS) typically describe gravity as suppressed by geometric dilution, not inverted by sign change. Our claim is stronger: the projection not only weakens gravity, the projected contribution is perceived by the brane as having the opposite sign i.e., the brane perceives the bulks attractive gravity as repulsive. The underlying gravity in the bulk remains attractive (standard GR); the inversion is a perceptual effect from the branes perspective, a feature of the specific bulk-brane coupling. This is a non-standard claim that would need to be checked against data if the model were to be developed further.

If the perceptual inversion holds, the projected bulk gravity (as perceived by the brane) is not the gravity we observe as attraction. What we observe as gravitational attraction must be the residual brane gravity the small net remainder of the branes attractive gravity after cancellation with the (perceived-inverted) bulk gravity. The branes gravity exceeds the perceived-inverted bulk gravity by a tiny amount; that tiny excess is what we measure as the gravitational force. The 4D events gravity, in 4D, remains attractive the inversion is only in the projection to the brane.

We distinguish two different un-cancelled contributions from the inversion:

- The small net brane excess (brane gravity minus inverted bulk gravity, taken as a force on matter) is the ordinary attractive gravity we observe. This is a force on matter, not a vacuum energy.
- The small un-cancelled fraction of the inverted bulk gravity is the dark energy a vacuum-energy-like contribution that drives cosmic expansion. This is a vacuum energy, not a force on matter.

Both arise from the inversion, but they play different roles: the first couples to matter (as a force), the second contributes to the vacuum energy (approximately constant during our brief slice of the 4D events full duration). The model does not currently derive the absolute magnitudes of either contribution; we note only that the qualitative structure (a small un-cancelled fraction of the inverted bulk gravity, approximately constant in our 3+1 dimensional frame because our universe is a brief slice of the 4D events full duration, acting as a cosmological-constant-like vacuum energy) is what the model proposes. The famous 10<sup>123</sup> discrepancy of the cosmological constant problem does not arise in this model from a ratio of these two un-cancelled fractions it arises

from the misidentification discussed in §2.6 (we were comparing observed dark energy to the 3+1 dimensional QFT vacuum energy, which is the wrong quantity to compare to).

The antigravity nature of the inverted bulk gravity is determined by the sign of the gravitational coupling, which is fixed by the inversion mechanism (a geometric postulate). The sign of the dark energy contribution is therefore fixed (repulsive, driving cosmic expansion). The magnitude of the dark energy contribution is approximately constant in our 3+1 dimensional frame because our universes lifetime is a brief slice of the 4D events full duration, during which the 4D events antigravity output is approximately constant. The dark energy density is therefore approximately constant in our frame (matching standard CDM behavior), and the total dark energy grows as the universe expands (because the universes volume grows while the density stays constant). The antigravity claim is about the sign, not the magnitude, and remains valid even as the dark energy total grows.

The dark energy is approximately constant in our 3+1 dimensional frame (because the 4D events antigravity output is approximately constant during our brief slice), and is not a current rate-dependent effect. (This is distinct from dark matter, which is the current rate-dependent effect of 2D universes being created by 3+1 dimensional energetic events see §2.6 and §2.7.) Dark energy is a 4D-event-driven geometric contribution; dark matter is the current activity in our 3+1 dimensional universe.

### 1.3.5 2.5 The two dark-sector products

The model has two distinct observable effects in the dark sector, which arise from different aspects of the dimensional projection mechanism:

- Dark energy arises from the bulk-brane gravity interaction (§2.4). The bulk gravity is inverted (repulsive) when projected into our 3+1 dimensional brane, and the un-cancelled fraction of this inverted bulk gravity is the dark energy. This is a 4D-event-driven geometric contribution a repulsive contribution that is approximately constant in our 3+1 dimensional frame (because our universe is a brief slice of the 4D events full duration, during which the 4D events antigravity output is approximately constant).
- Dark matter arises from the scale-invariant principle (§2.3) and the energy-conserving return of 2D universe energy to 3+1D as 2D universes lives end. Every energetic event in our 3+1 dimensional universe creates a 2D universe. Each 2D universe lives for some time (per  $\tau_{2D} = \ell/c$ ), evolves (its own Big Bang, expansion, physics), and eventually ends. The form of the ending depends on the 2D universes internal dynamics a 2D universe with high matter density (relative to its own dark energy) may undergo a Big Crunch (gravitational collapse, brief intense death-flash); a 2D universe where its own dark energy dominates may reach heat death (slow, diffuse dispersal of matter); other endings (cyclic, Big Rip, etc.) are also possible in principle. The model does not currently specify which ending is typical for 2D universes of different sizes; however, since every 2D universe has its own dark energy (per the universal bulk-brane cancellation, §2.6) and that dark energy is repulsive, the natural ending for small 2D universes (where matter density is low) is heat death the 2D universes own dark energy slowly pulls its matter apart, just as our own dark energy is pulling our universe apart. The energy of the 2D universe has to return to 3+1D eventually (energy conservation), and this cumulative return whether as brief death-flashes (Big Crunch) or slow diffuse leakage (heat death) is the dark matter. The total dark matter is set by the cumulative energy of all 2D universe endings; the form depends on the mix of endings. The observable dark matter we see is consistent with a smooth, distributed background (matching heat-death-dominant

returns), with potentially a small impulsive component (from occasional Big Crunch returns) at SN/BH/AGN sites that is currently below detection thresholds. The model is intentionally ending-agnostic at the 2D universe level, just as it is at the 3+1D level (§2.8) – the specific ending affects the form of the dark matter, not the total.

These two effects are distinct in origin: dark energy comes from the bulk-brane interaction (the one 4D event), while dark matter comes from the scale-invariant principle applied to many 3+1 dimensional events. Both are aspects of dimensional projection, but they are not both products of the bulk-brane cancellation – only dark energy is. Dark matter is a separate effect of the same underlying dimensional principle.

Lensing and the inversion principle. A natural question is: where does the inversion happen, and where does normal gravity apply? The model has a specific answer: the downward dimensional projection (parent → child universe) inverts the sign of gravity, but the upward back-projection (child → parent) does not. This gives a clean and consistent picture:

- Ordinary matter lensing (3+1D mass): attractive (normal lensing, light bends toward mass). No inversion within 3+1D; ordinary matter is just normal attractive gravity.
- Dark matter lensing (2D universe back-projection): attractive (normal lensing). The 2D universes net gravity is attractive (per the universal bulk-brane cancellation, below), and the back-projection from 2D to 3+1D does not invert the sign – so 2Ds attractive gravity projects back to 3+1D as attractive = dark matter. (Note: the downward projection from 3+1D to 2D does invert 3+1Ds net attractive gravity into 2Ds bulk antigravity, which is then mostly cancelled by 2Ds native attractive gravity – leaving 2Ds small net attractive residue. It is this residue that back-projects up to 3+1D without further inversion.)
- Dark energy lensing (4D event projection): repulsive (anti-lensing in principle, but dark energy is approximately uniform in 3+1D, so it doesn't cause local lensing – its anti-gravity is observed on cosmological scales as cosmic expansion, not local anti-lensing). The 4D events gravity is attractive in 4D; the downward projection to 3+1D inverts it, giving repulsive gravity in 3+1D. The un-cancelled fraction of this 4D→3+1D antigravity is the dark energy.
- 2D universe internal physics: similar to 3+1D, with attractive net gravity and its own dark energy (which is repulsive). The universal bulk-brane cancellation mechanism (per §2.6) gives attractive gravity on each brane, regardless of which level. The 2D universes own gravity is attractive (not repulsive), because the bulk-brane cancellation in 2D gives the same kind of weak attractive gravity that we observe in 3+1D. The 2D universe also has its own dark energy (per the cascade), which is repulsive. The competition between attractive gravity and repulsive dark energy determines the 2D universes ending: if matter density is high (large events, e.g., AGN or BH-scale), attractive gravity wins and the 2D universe undergoes a Big Crunch (gravitational collapse); if matter density is low (small events, e.g., LHC or small SN), dark energy wins and the 2D universe slowly disperses in heat death. Both endings return the 2D universes energy to 3+1D as dark matter – the Big Crunch case as a brief, intense, localized death-flash; the heat death case as a slow, diffuse, distributed leakage. The mix of these depends on the 2D universes size, which is set by the original 3+1D event energy. The model is ending-agnostic at the 2D level (just as at the 3+1D level, §2.8), and acknowledges that the form of dark matter depends on this mix.

The inversion principle: downward only. The inversion is a property of the downward dimensional projection (parent → child universe). When a parents gravity is projected into a child universe, the sign inverts – so the child experiences the parent as antigravity. The childs own native gravity,

plus its bulk-brane cancellation with the inverted parent projection, leaves a small net attractive residue (the child's ordinary gravity). The upward back-projection (child  $\rightarrow$  parent) does not invert the child's small net attractive residue; it is felt in the parent as attractive, contributing to the parent's dark matter. This asymmetry is what makes the cascade consistent with both the dark matter observations (attractive, normal lensing) and the dark energy observations (repulsive, cosmic expansion).

The universal bulk-brane cancellation. A cleaner way to think about the cascade is that every level has the same basic structure as 3+1D: a bulk (the level above) and a brane (the level itself), with the bulk-brane interaction giving a weak attractive gravity on each brane. The downward perceptual inversion principle (downward projection is perceived as inverted by the child, upward back-projection is not perceived as inverted by the parent) is still a postulate (not a derivation), but its consequence is that every level is similar to 3+1D in structure:

- All universes have 3+1D spacetime (per §2.1)
- All universes have bulk-brane cancellation (per §2.6)
- All universes have attractive net gravity
- All universes can have stable structures (in principle)
- All universes can gravitationally collapse (Big Crunch)
- All universes end (Big Crunch, heat death, or other), and the energy return to the parent is the parent's dark matter contribution (the form depends on the ending)
- Death-flashes project back to the parent level as dark matter

So the cascade is a fractal hierarchy of similar universes, each with: - Attractive net gravity (similar to our universe) - Stable structures (in principle, at appropriate scales) - An ending (Big Crunch, heat death, or other) that returns the universe's energy to its parent as dark matter contribution (the form depends on the ending: Big Crunch gives a brief, intense death-flash; heat death gives a slow, diffuse return)

The depth of the cascade just means smaller and briefer levels, not fundamentally different physics. Each level is similar to 3+1D in structure, with bulk-brane cancellation giving a weak attractive gravity, its own dark energy (repulsive), and its own ending (Big Crunch, heat death, or other). The 2D universe is a miniature 3+1D universe with its own gravity, its own dark energy, its own ending, and its own energy return to 3+1D as dark matter. The 1D universe is a miniature 2D universe with its own gravity, its own dark energy, its own ending, and its own energy return to 2D. And so on.

This is similar to the standard brane-world picture (RS99), but applied to every level of the cascade. Each level has a bulk (above) and a brane (itself); gravity inverts at the bulk-brane boundary; the cancellation gives a weak attractive gravity on each brane. The cascade extends this picture to multiple levels, with each level being a miniature brane-world in its own right.

The inversion principle is a postulate, not a derivation. The universal bulk-brane cancellation picture (every level similar to 3+1D) depends on the downward perceptual inversion principle that downward dimensional projection (parent  $\rightarrow$  child universe) is perceived by the child as having the opposite sign, with the bulk-brane interaction giving weak attractive gravity on each brane. The upward back-projection (child  $\rightarrow$  parent) does not invert the perception; the parent perceives the child's net attractive gravity as attractive, contributing to the parent's dark matter. This is a postulate of the model, not a derivation. We do not currently know why the downward projection is perceived as inverted but the upward back-projection is not; the cascade simply assumes this asymmetry of the projection mechanism. Importantly, the underlying gravity in each parent universe remains attractive (standard GR); the inversion is a perceptual effect from the child's perspective, a feature

of the specific bulk-brane coupling. The postulate is motivated by the standard GR mechanism for negative effective gravitating density: a quantum field with  $P < -\frac{1}{3}\rho$  has effective gravitating density  $\rho + 3P < 0$ , which sources repulsive gravity in standard GR (the same mechanism that drives cosmic inflation and dark energy in our universe). The cascades bulk-brane coupling is postulated to produce a similar effect: the projected effective density on the brane is  $\rho_{\text{eff}} = \rho_{\text{proj}} + 3P_{\text{proj}} < 0$ , sourcing repulsive gravity on the brane. The asymmetry (down inverts, up doesn't) is a specific feature of the bulk-brane coupling that the model does not derive; it is the least constrained postulate in the cascade. The user (in private communication) has noted that this is a strong claim, and that the consequence of the claim is that every level has the same basic structure as 3+1D (bulk above, brane itself, weak attractive gravity, dark energy, an ending that returns energy to the parent as dark matter). If the downward perceptual inversion principle is wrong (i.e., the brane does not perceive the projected bulk gravity as inverted), the entire framework changes – the dark matter and dark energy mechanisms would need to be re-derived, and the universal bulk-brane cancellation would not hold. The model is committed to the downward perceptual inversion principle as a fundamental postulate, but acknowledges that this is a strong claim that requires experimental or theoretical justification. The consistency of the principle with the dark matter / dark energy observations (e.g., dark matter lensing normally, dark energy being repulsive, dark energy equation of state  $w \approx -1$  matching a cosmological-constant-like effective pressure) is suggestive but not conclusive evidence. A specific implementation of the model would need to derive the perceptual asymmetry (down inverts, up doesn't) and the specific bulk-brane coupling that produces the negative effective gravitating density, from a deeper theory (e.g., the geometry of the bulk-brane coupling), which is left to future work.

The cascades inversion principle is therefore directional: the downward projection inverts, the upward back-projection does not. The consequence of this directional inversion is the universal bulk-brane cancellation – every level has the same structure as 3+1D, with attractive net gravity, dark energy (repulsive), an ending (Big Crunch, heat death, or other), and energy return to the parent as dark matter. The cascade does not alternate between collapsing and expanding levels based on parity; it applies the same bulk-brane physics at every level, just at different scales. This universal interpretation is what allows the model to produce both dark energy (anti-gravity from the 4D3+1D downward inversion, no local lensing) and dark matter (attractive, from the 2D3+1D upward back-projection of the 2D universes net attractive gravity, lensing normally) from the same dimensional-projection mechanism, without contradiction, at every level of the cascade. The model is ending-agnostic at every level (§2.8) – the specific ending (Big Crunch vs heat death vs other) affects the form of the dark matter, not the total.

The specific microphysical mechanism by which the dimensional projection produces these effects is not determined by this proposal alone. The geometry of the extra dimensions, the field content of the bulk, and the coupling structure would together determine the details. We discuss the observable consequences of these products in §2.6 and §2.7.

Summary of the cascade framework. The cascades core claims, distilled from §2.1–§2.9, are:

1. Dimensional projection mechanism: A single ongoing 4D event projects into our 3+1D universe, with the 4D events antigravity inverting on projection, giving a weak attractive gravity on the 3+1D brane (per the bulk-brane cancellation, §2.6).
2. Scale invariance: Every energetic event in 3+1D creates a child universe at the next level (per §2.3). The child universe is embedded in the parents spacetime, with 3+1D spacetime at smaller scales (per §2.1).
3. Universal bulk-brane cancellation: Every level has the same basic structure as 3+1D (bulk above, brane itself, weak attractive gravity, dark energy which is repulsive, an ending that

returns energy to the parent). The depth of the cascade means smaller and briefer levels, not fundamentally different physics. The form of the energy return (Big Crunch brief death-flash; heat death slow diffuse return) depends on the specific ending at each level.

4. Two dark-sector products: Dark energy is the un-cancelled fraction of the 4D events anti-gravity, projected down to 3+1D (one downward inversion, repulsive). Dark matter is the back-projection of 2D (child) universes from 3+1D energetic events (one downward inversion at 3+1D2D that sets up 2Ds bulk antigravity, then bulk-brane cancellation leaves 2Ds small net attractive residue, which projects up to 3+1D without further inversion = attractive).
5. The downward perceptual inversion principle (postulate): Downward dimensional projection (parent child universe) is perceived by the child as having the opposite sign i.e., the projected contribution to the childs effective gravity is repulsive, even though the parents gravity in its own frame remains attractive (standard GR). Upward back-projection (child parent) does not invert the perception; the parent perceives the childs net attractive residue as attractive. This asymmetry is a postulate, not a derivation, and the model is committed to it as a fundamental claim. The underlying physics in each parent universe is unchanged; the inversion is a perceptual feature of the projection mechanism, grounded in the standard GR  $\rho + 3P < 0$  mechanism for negative effective gravitating density (the same mechanism that drives cosmic inflation and dark energy in our universe). It is what makes the dark matter attractive (back-projection from 2D to 3+1D preserves the attractive sign) and the dark energy repulsive (downward projection from 4D is perceived as inverted by 3+1D).
6. Energy return from ending universes dark matter: Each child universes life ends (Big Crunch, heat death, or other), and its energy returns to the parent. The form of the return depends on the ending: Big Crunch gives a brief, intense, localized death-flash; heat death gives a slow, diffuse, distributed return. The total dark matter is set by the cumulative energy of all 2D universe endings. The model is ending-agnostic at the 2D level, just as it is at the 3+1D level.
7. Quantum mechanics as 2D-level physics projection: 3+1D quantum mechanics is the projection of the 2D-levels Standard Model (per §2.8). The cascade transitions between regimes with different effective theories at different scales.
8. Cascade is infinite in principle, finite in practice: The cascade has no a priori depth limit (labels can extend to -1D, -2D, etc.), but in practice the depth is finite (energy decreases at each level, eventually too small to create a new universe).
9. Five possible universe endings: fixed-time boundary, cyclic, diminishing cyclic, Big Rip, Big Freeze / heat death all empirically distinguishable by future observatories (Euclid, Roman, LSST, SKA).
10. D-labels are placeholders: 2D universe, 1D universe, 0D universe, -1D universe are legacy labels for levels in the hierarchy of nested universes, not claims about spacetime dimensions. All universes are 3+1D (per §2.1).

These claims are the core of the model. §4 extends the model with speculative applications (sub-mm gravity, CMB, dark matter / activity correlation, black holes, constants, weak/strong forces, etc.).

### 1.3.6 2.6 The energy budget, the cosmological constant, and the bulk-brane cascade

Energy conservation is standard. The model does not propose a new conservation law. Energy is conserved in the usual sense: the 4D event is an ongoing energetic process with some total energy budget  $E_{4D}$  (integrated over its full duration in 4D time), and our 3+1 dimensional universes total mass-energy is the portion of that total energy that has been delivered to the brane during



our universes lifetime (a brief slice of the 4D events full duration). The 4D events full duration and the 4D-side power output are not specified by the model; the model only specifies that the standard conservation law applies at the level of total energy. There is no missing energy, no projection factor multiplying or attenuating the energy, and no infinite regress through lower dimensions. The standard conservation law suffices.

Symmetries and conservation laws. The dimensional-cascade framework takes the standard conservation laws and symmetries of physics as given: - Energy conservation (per §2.6 above): the model does not propose a new conservation law. - Momentum and angular momentum conservation: the model assumes standard conservation of 3-momentum and angular momentum in 3+1D. The 4D events internal momentum structure is unspecified; the model only requires that the projected 3-momentum and angular momentum in 3+1D are conserved. - CPT symmetry: the model assumes CPT is conserved (per standard physics). The cascade does not propose a new CPT-violating mechanism. - Lorentz invariance: the model assumes standard Lorentz invariance in 3+1D. The 4D events internal Lorentz structure is unspecified; the model only requires that the projected 3+1D physics respects Lorentz invariance. - Equivalence principle: the model assumes the equivalence principle (gravitational and inertial mass are equal) in 3+1D, per general relativity. The cascade does not propose violations of the equivalence principle. - Locality: the model assumes standard locality in 3+1D. The 2D universe back-projection (dark matter) is non-local in the sense that the 2D universes whole gravitational effect is felt at the 2D universes 3+1D location, but this is consistent with locality in 3+1D (the gravitational effect propagates at the speed of light). - Thermodynamics (2nd law): the model assumes standard thermodynamics. The cascade does not propose violations of the 2nd law.

These are assumptions of the model, not derivations. The cascade is consistent with standard physics in 3+1D; it extends the framework to include dimensional projection (4D  $\rightarrow$  3+1D  $\rightarrow$  2D), but the projected 3+1D physics is standard.

Honest acknowledgment: what the model does and does not address. The dimensional-cascade framework is a thought experiment that reinterprets the dark sector and the weakness of gravity through a single geometric process. It is not a complete theory of everything, and it explicitly does not derive several features of standard cosmology that are well-established experimentally: - The origin of the primordial perturbations (the seed fluctuations for cosmic structure) the cascade takes these as given, noting that the 4D event must be spatially extended and approximately homogeneous to match the observed near-scale-invariant CMB power spectrum. - Cosmic inflation the very early accelerated expansion that solves the horizon, flatness, and monopole problems. The cascade is compatible with inflation (the 4D event could in principle have an inflationary phase) but does not derive it. - Baryogenesis the matter-antimatter asymmetry. The cascade is compatible with baryogenesis (the 4D event could in principle generate the asymmetry via C and CP violation in the projection) but does not derive it. - Big Bang nucleosynthesis the light element abundances. The cascade takes BBN as given, noting that the 4D event scenario must be consistent with the observed D,  $^3\text{He}$ ,  $^4\text{He}$ , Li abundances. - The Standard Model particle spectrum the masses and couplings of all known particles. The cascade takes these as given; the 4D events internal physics is unspecified in the model. - Neutrino masses and oscillations the cascade takes these as given (this is why §4.6 and §4.12 were removed in v1.58).

The model is a framework for the dark sector and gravity, not a complete cosmological theory. The 22 references and §4 extensions are speculative applications of the model, not derivations.

The cosmological constant problem reframed. A natural objection is the cosmological constant problem: the natural vacuum energy from 3+1 dimensional quantum field theory is far larger than the observed dark energy density the discrepancy is often quoted as approximately  $10^{120}$  orders of magnitude, though the precise value depends on the assumed cutoff, field content, and regularization

scheme.

In our model, the actual dark energy density in our universe is not the 3+1 dimensional QFT vacuum energy. It is the un-cancelled fraction of the inverted bulk gravity (ğ2.4) a contribution from the 4D event that is approximately constant in our 3+1 dimensional frame (because the 4D events antigravity output is approximately constant during our brief slice of the 4D events full duration), and unrelated to 3+1 dimensional zero-point energy calculations. The 3+1 dimensional QFT calculation gives the right answer for what it computes (the 3+1 dimensional zero-point energy of Standard Model fields), but this is not the right physical quantity to compare to the dark energy in our universe, because dark energy in our universe is bulk gravity residue, not 3+1 dimensional QFT vacuum energy. The famous 10<sup>123</sup> is the disagreement between these two ways of computing the same conceptual quantity (vacuum energy in our universe), one of which is computing the wrong thing.

The cosmological constant problem, in this framing, becomes a problem of identification: we were computing the wrong quantity. The correct computation is the residue of the inverted bulk gravity, not the 3+1 dimensional QFT zero-point energy. (We acknowledge that the model does not yet quantitatively derive the un-cancelled fraction from the geometry of the 4D event the specific value of 10<sup>123</sup> is not predicted by the model but the qualitative point that the bulk-gravity residue is not the 3+1 dimensional QFT zero-point energy is well-motivated by the dimensional-projection mechanism.)

Dimensional analysis of the cascade. The cascade principle of ğ2.4 says that at every dimensional level, the projected (inverted) gravity and the native gravity nearly cancel. The small un-cancelled residue is the effective gravity at that level. We can parameterize the cancellation by a small dimensionless parameter  $\epsilon$  at each level:

$$G_{D-1}^{eff} = \epsilon \cdot G_{D-1}^{native}$$

where  $\epsilon \ll 1$ . The un-cancelled antigravity (the cascades gravitational contribution) is also of order  $\epsilon \cdot G_{D-1}^{native}$  in magnitude that is, the un-cancelled antigravity is of the same order as the ordinary gravity (both are small because of the near-cancellation). Crucially, the cascades un-cancelled antigravity is a gravitational coupling (units of  $G$ ), not a vacuum energy density (units of energy<sup>4</sup>). Converting the cascades gravitational coupling to a vacuum energy density requires a mass scale or length scale: for example, if we associate the antigravity with a Planck-scale vacuum energy, then  $\rho_{DE,model} \sim \epsilon \cdot M_{Pl}^4 \sim 10^{-38}$  in natural Planck units (where  $M_{Pl}^4 \sim 1$ ), but the observed dark energy is  $\sim 10^{-123}$  in natural Planck units. So the cascades un-cancelled antigravity as vacuum energy is 10<sup>85</sup> larger than the observed dark energy. The cascade therefore explains why the dark energy is qualitatively small (its a near-cancellation effect, suppressed by  $\epsilon$ ), and it gives a quantitative prediction of  $\sim 10^{-38}$  in natural units for the total antigravity produced by the cascade. The observed dark energy in 3+1D is  $\sim 10^{-123}$ , which is 10<sup>-85</sup> times smaller than the cascades prediction. The cascade produces this antigravity at the 4D 3+1D level, and only a small fraction  $f_{back} \sim 10^{-85}$  of it remains in 3+1D as observable dark energy; the rest ( $1 - f_{back} \approx 1$ ) is absorbed into the 3+1D vacuum structure in a way that does not contribute to the observable dark energy density (it could be the 4D bulk dark matter, or a non-projected component). The cascades prediction  $\times$  fraction staying in 3+1D = observed dark energy:  $10^{-38} \times 10^{-85} = 10^{-123}$  (matches observation). The fraction  $f_{back} = 10^{-85}$  is a postulate of the model, and represents the effective fraction of cascade-produced antigravity that survives as observable dark energy. (Note: this is distinct from the cumulative 2D universe antigravity, which is internal to the 2D universes and does not project back to 3+1D as dark energy. The 3+1Ds dark energy comes only from the 4D event, not from cumulative 2D universe antigravity. See ğ2.5 and ğ2.7.) The observed 10<sup>123</sup>

(hierarchy) is then  $\epsilon$  at the 3+1D level: the native 3+1D gravitational coupling is larger than the effective coupling by a factor of  $\sim 10^{38}$ . Equivalently,  $\epsilon_{3+1D} \sim 10^{-38}$ . The cascade explains the 10 $\S$  hierarchy quantitatively (as a near-cancellation of  $G_{native}$  and  $G_{proj}$ ), and it gives a qualitative explanation for the smallness of the dark energy (as another near-cancellation effect), but it does not give the quantitative value of the dark energy. The model reframes the 10 $\S$  cosmological constant problem as a misidentification: the 3+1D QFT vacuum energy is the wrong quantity to compare to the observed dark energy. The dark energy in the model is the un-cancelled antigravity residue of the cascade, modulated by the staying fraction  $f_{back} \sim 10^{-85}$ . The model does not claim to quantitatively derive the 10 $\S$  hierarchy, the absolute value of the dark energy density, or the 10 $\S$  discrepancy from the cascade alone. A specific implementation of the model would need to derive the exact un-cancelled fraction from the geometry of the dimensional projection, which would in turn predict the absolute value of the dark energy density and the quantitative value of the 10 $\S$  ratio.

We emphasize that this dimensional analysis is qualitative and not a derivation. The quantitative values of  $\epsilon$  at each level of the cascade are not derived in this model. A specific implementation would need to compute  $\epsilon_{3+1D}$  and  $\epsilon_{2D}$  from the geometry of the dimensional projection, which is left to future work.

Connection to the hierarchy problem. The hierarchy problem is the question: why is gravity  $\sim 10^{\S}$  times weaker than the other fundamental forces in 3+1 dimensional physics? In our model, ordinary gravity (the gravitational force between visible masses) is the small net remainder of the branes attractive gravity after cancellation with the inverted (repulsive) bulk gravity (§2.4). The branes gravity exceeds the inverted bulk gravity by a tiny amount; that tiny excess is what we measure as ordinary gravity. The fundamental gravitational coupling in our 3+1 dimensional frame is therefore small because the bulk-brane cancellation is almost exact the residual brane excess is small, leaving only a tiny fraction of the bulks gravity to be observed as attractive force. This is the bulk-brane cancellation interpretation of the hierarchy: gravitys weakness is a consequence of the dimensional-projection mechanism nearly cancelling the branes gravity with the bulks inverted gravity.

The 2D universes collective gravity is also a consequence of bulk-brane cancellation, applied at the next level of the dimensional cascade. In the 2D universe, the 3+1 dimensional events gravity is inverted (antigravity) and mostly cancelled with the 2D universes own attractive gravity. The 2D universes attractive gravity, projected back into our 3+1 dimensional frame, is the small net remainder of this cancellation exactly the same mechanism that makes ordinary 3+1 dimensional gravity weak. The cumulative effect of all 2D universes gravity, projected back into 3+1D, is the dark matter a weak, diffuse, gravitationally-interacting background. Dark matters effective coupling is not intrinsically weaker than ordinary gravitys coupling in 2D; it is weaker in the 3+1 dimensional projection because of the bulk-brane cancellation applied to the 2D universe.

This is a unification of two effects that might otherwise seem separate: the hierarchy problem (10 $\S$ ) and dark matters apparent weakness are both consequences of the same bulk-brane cancellation mechanism, applied at different levels of the dimensional cascade. The hierarchy is the 3+1 dimensional projection; the dark matter is the cumulative effect of 2D projections. The mechanism is the same; the scale of the dimensional projection differs.

A natural question: does the 2D universes antigravity (its internal dark energy) help cancel the 4D antigravity, contributing to gravitys weakness? In the current model, the 2D universes effect is separate from the bulk-brane cancellation the 2D universes attractive gravity is dark matter (additional to ordinary gravity), and their antigravity is internal to the 2D universe (not directly projecting back into 3+1D). But in a more developed model, the 2D universes could play a back-reaction role, contributing to the cancellation of the 4D antigravity and reducing the net effective

gravity in 3+1D. This is an interesting extension of the model, but is not currently derived.

Asymmetry between dark energy and dark matter math. The dark energy and dark matter are both products of the dimensional-cascade framework, but their quantitative derivations are asymmetric. The dark energy math works cleanly once we include the back-reaction fraction (§2.6 above): cascade prediction  $\mathbb{E} f_{\text{back}} = \text{observed}$ . The back-reaction fraction is a single postulate that fixes the dark energy to the right value. The dark matter math, by contrast, is underdetermined: it depends on the event rate  $R$ , the 2D universe lifetime  $\tau_{\text{2D}}$ , the event energy  $E_{\text{2D}}$ , and the projection fraction  $G_{\text{2D}}^{\text{projected}}/G_{\text{4D}}$  four free parameters (rather than one) that must be specified to make a quantitative prediction. The current model gives a qualitatively plausible picture (the cumulative effect of 2D universes is consistent with the observed dark matter density for plausible parameter values) but does not give a unique numerical prediction. The asymmetry reflects the fact that dark energy is a single global quantity (the vacuum energy), while dark matter is a cumulative quantity (the integral over many 2D universes). Dark energy is fixed by geometry (the projection rule), while dark matter is fixed by event statistics (the rate and energy distribution of 3+1D energetic events). We acknowledge this asymmetry as a limitation of the current model.

A note on the 2D universes antigravity. The 2D universes antigravity is internal to the 2D universe it is the 2D universes own dark energy (analogous to the 3+1D dark energy, but at the 2D level). The antigravity does not project back to 3+1D, because the cascade is hierarchical each dimensional levels gravity and antigravity are local to that level, with the attractive component projecting forward to child dimensions and the antigravity remaining internal to the universe itself. The 3+1Ds dark energy is a separate contribution from the 4D event, not the cumulative 2D universe antigravity. The two dark-sector components have distinct dimensional origins: dark energy from 4D 3+1D, dark matter from 2D 3+1D. They are not parallel back-projections from the same source.

The 10 $\eta$  (hierarchy) and 10 $\zeta$  (cosmological constant) are not the same principle in this model. 10 $\eta$  is a bulk-brane cancellation effect (the residual brane excess after cancellation with the inverted bulk gravity is small). 10 $\zeta$  is a misidentification effect (we were comparing observed dark energy to the wrong theoretical quantity). The two are independent and unrelated. We note that the quantitative relationship between these two numbers is not derived in this model they emerge from different aspects of the dimensional projection. (Dark matters apparent weakness is also a bulk-brane cancellation effect, applied to the next level of the dimensional cascade. The 10 $\eta$  hierarchy and the dark matters effective coupling are related by the dimensional cascade structure, though the specific quantitative relationship is not derived in this model.)

The universes lifetime. The total lifetime of our universe is some fraction of the 4D events full duration in 4D time. The 4D events spatial extent, divided by the 4D speed, gives the 4D events full duration; our universes lifetime is determined by the projection mechanism (which is not specified in this thought experiment). The universes energy density of matter, dark matter, and dark energy evolves as the universe expands, with the 4D event providing a continuous flux of antigravity that is approximately constant during our universes brief slice. The universe does not run down or fade out in this sense (no thermodynamic depletion, no second-law-of-thermodynamics-driven heat death). The universe ends either at a fixed-time boundary (the current papers interpretation: the boundary of the 4D events spacetime) or via a Big Crunch that re-nucleates a new 4D event (the cyclic interpretation, consistent with the cascades scale invariance see §2.8 below for the philosophical distinction). Both are consistent with the cascade framework; this paper adopts the fixed-time boundary framing for simplicity.

The nature of the boundary: what happens at the edge? The fixed-time boundary framing is under-specified in the model, and the user (in private communication) has pointed out that the boundary has to be strong (otherwise the perceptual inversion wouldnt be maintained). We do not

currently specify the exact nature of the boundary, but the perceptual-inversion constraint limits the possibilities:

- The perceptual inversion is maintained throughout the universes lifetime (the cascades postulate, §2.4: the 4D events projected antigravity is constant in 3+1D frame, giving constant dark energy). This means the boundary cannot be a gradual weakening of the antigravity that would contradict the constant-dark-energy observation.
- The boundary is therefore abrupt in the antigravity sense: the 4D events projected antigravity is constant during the universes lifetime, and then suddenly stops (or changes) at the boundary. The perceptual inversion is strong because the antigravity output is constant (and strong) throughout, not because the boundary is strong in the fade-out sense.

The boundary has two aspects:

1. Antigravity boundary (abrupt): The 4D events antigravity is constant during the universes lifetime, then stops abruptly at the boundary. This is the cascade-specific feature. The strong boundary is here.
2. Matter boundary (gradual): Matter density drops as  $a^{-3}$  (cosmic expansion) and the universe becomes empty over time. This is the Big Freeze / heat death, which is gradual and observationally consistent with standard  $\Lambda$ CDM cosmology.

The universe ends when both boundaries are reached: the antigravity stops abruptly (cascade-specific), and matter is empty (Big Freeze). These two boundaries may happen at similar times (when the 4D event ends), but they're distinct in nature.

Given these constraints, the three possibilities for the boundary become:

1. Abrupt vanishing with abrupt antigravity stop: The 4D events antigravity stops abruptly at a specific time. The 3+1D universe vanishes at the same moment. The universe simply stops existing at a specific time, with no warning and no transformation. This is the simplest interpretation of the antigravity boundary, but it raises the information paradox (what happens to matter, energy, and information at the boundary?).
2. Gradual matter fade-out with abrupt antigravity stop: The 4D events antigravity stops abruptly, but the matter in the 3+1D universe gradually disperses (Big Freeze) over some finite time after the antigravity stops. The universe becomes empty gradually, but the antigravity is gone. This is essentially the Big Freeze with an abrupt antigravity boundary on top.
3. Phase transition with abrupt antigravity stop: The 4D events antigravity stops abruptly, and the 3+1D universe undergoes a phase transition to a different state (e.g., merges back into 4D bulk, becomes a 2D universe in a higher-D cascade, etc.). This avoids the information paradox by transforming the universe rather than vanishing it.

The model does not currently specify which of these is correct, but the gravity-flip constraint is now explicit: the antigravity boundary is abrupt, not gradual. The matter boundary can be either abrupt (option 1) or gradual (option 2, Big Freeze) or transformative (option 3). A specific implementation would need to specify the exact nature of the matter boundary, which is left to future work. We note that option 2 (gradual matter fade-out with abrupt antigravity stop) is the most natural and observationally consistent interpretation, and combines the cascades antigravity constraint with the standard Big Freeze picture.

### 1.3.7 2.7 The products

The energy of the original event, projected into our 3+1 dimensional brane, would manifest as whatever particles, fields, or vacuum energy the geometry and bulk field content allow. The two distinct observable effects identified in this model are:

- **Dark matter.** A non-luminous, gravitationally-interacting substance that does not couple to electromagnetism or the strong force. In the scale-invariant version of this model (§2.3), dark matter is the current collective gravitational signature of all 2D universes being currently created by 3+1 dimensional energetic events in our 3+1 dimensional world. Dark matter is not a particle; it is the gravitational effect of tiny universes being created now, weighted by event size. The model provides both a geometric role and a specific identification for dark matter though the proportionality constants and details depend on the specific geometry and event spectrum, which we have not derived.
- **Dark energy.** A uniform vacuum-like energy driving cosmic acceleration. In this model, dark energy is the un-cancelled antigravity of the 4D event, projected into our 3+1 dimensional frame as the un-cancelled fraction of the 4D events inverted gravity (per §2.4). The 3+1D sees a single contribution to its dark energy from the 4D event, with the magnitude set by the cascade + staying fraction  $f_{back} \sim 10^{-85}$  (§2.6). The dark energy is approximately constant in our 3+1 dimensional frame because our universe is a brief slice of the 4D events full duration, during which the 4D events antigravity output is approximately constant. The dark energy has a distinct dimensional origin from dark matter: dark energy comes from the 4D 3+1D projection, dark matter from the 2D 3+1D back-projection. The two are complementary aspects of the cascade, not parallel back-projections from the same source. (The 2D universes own antigravity is internal to the 2D universes, and does not project back to 3+1D see §2.5 and §2.6.)

The model does not fully address the Standard Model itself (the origin of electron mass, quark masses, gauge couplings, etc.). The Standard Model is taken as given in the core model; the dimensional projection mechanism is proposed as an explanation specifically for the dark sector (dark matter and dark energy) and the weakness of gravity. The core model (as presented in §2.1§2.7) takes neutrino properties as given. The model does not currently offer a dimensional-cascade interpretation of neutrino mass or oscillation.

We do not claim to derive the relative abundances of dark matter and dark energy. This is left to the community.

### 1.3.8 2.8 The universes lifetime

In this picture, the universes lifetime is some fraction of the 4D events full duration in 4D time. The 4D events spatial extent, divided by the 4D speed, gives the 4D events full duration in 4D time; the projection mechanism then selects a fraction of this full duration as our universes lifetime. The specific rule by which the 4D events full duration maps to our 3+1 dimensional lifetime is not specified in this thought experiment. We assume only that some such rule exists, mapping the 4D events full duration to a 3+1 dimensional lifetime that is some fraction of it.

The model makes the following claim about the universes energy:

- The 4D event has a duration in 4D time,  $t_{4D}$ . During this entire duration, the 4D event is active it is not a one-time past event, but an ongoing process in 4D time. Our 3+1 dimensional universe exists for a fraction of  $t_{4D}$  (we dont see the 4D events full duration in our frame;

we only see a slice of it). The 4D events spatial extent and duration in 4D time are much larger than the corresponding quantities in our 3+1 dimensional frame.

- The 4D events antigravity leakage is approximately constant over the brief slice that our 3+1 dimensional universe occupies in 4D time. This is why dark energy appears constant in our 3+1 dimensional frame: from the 4D perspective, our universe is a brief moment during which the 4D events gravity is roughly unchanging. (Just as a microscopic black hole created in a high-energy collision sees a brief moment of constant high energy before the parent collision ends and the black hole evaporates.)
- Throughout our universes lifetime (a slice of the 4D events full duration), the 4D event is continuously feeding antigravity into our 3+1 dimensional frame. The dark energy is not a one-time injection at the Big Bang; it is an ongoing flux from the 4D event over the entire lifetime of the universe. The dark energy density in our 3+1 dimensional frame is approximately constant during this slice (because the 4D events antigravity output is approximately constant).
- The model does not currently specify what determines our universes lifetime (the duration of the slice of the 4D event that we occupy). The end of our universe is a boundary beyond which the projection ceases to exist not because the 4D event ends, but because of some other reason (perhaps the stability of the projection, perhaps a change in the 4D events gravity over its full duration).

The universe does not run down or fade out in this model in the sense of a purely thermodynamic heat death (no entropy-driven process depletes the universe). However, the universe can fade out in the sense of a Big Freeze (expansion-driven emptiness) if the 4D events antigravity is constant and theres no fixed-time boundary see the list of possible endings below. The cascades default is a fixed-time boundary, but the Big Freeze is a natural alternative if the boundary interpretation is wrong.

This is structurally different from both standard cosmological fates:

- Standard heat death: The universe ends as a cold, dilute, near-equilibrium state after entropy has been maximized. In our model, the end is not an entropy-driven process; it is a fixed-time boundary.
- Big Crunch / cyclic cosmology: The universe ends by recontracting to a singularity, possibly triggering a new Big Bang. In the current paper, the end is not a recontraction; it is a fixed-time boundary beyond which there is no spacetime to recontract into. However, the cascades scale invariance (g2.3) is consistent with a cyclic interpretation: a collapsing 3+1D universe (Big Crunch) is a high-energy event, and by the cascade principle, such an event could itself create a new 4D event analogous to a vacuum bubble (cavitation) in water. In this cyclic version, the Big Bang Big Crunch are paired events: the previous universes collapse nucleates our universes Big Bang, and our universes collapse (if it happens) will nucleate the next. Crucially, the cyclic cascade need not be exact recurrence: each Big Crunch releases some energy to 4D bulk (analogous to surface tension in a collapsing bubble, or to BH evaporation), so each successive 4D event is smaller than the previous, and each successive 3+1D universe is less energetic. The cascade is therefore diminishing: cycles become smaller over time, not larger, and the cascade eventually dwindles to the point where it can no longer support complex structure (a 4D event too small to create a 3+1D universe with stars, galaxies, or observers). This avoids the eternal return problem (the universe doesnt infinitely recur in the same state) and gives the cascade a direction (toward smaller cycles, not toward eternal recurrence). The principle of energetic events create universes is robust across cycles even

if the specific values (masses, couplings, dark energy density) drift from cycle to cycle. The choice between fixed-time boundary (this paper's default), cyclic vacuum-bubble (with exact recurrence), and diminishing cyclic (cyclic but with energy loss to 4D bulk) is philosophical (where does the 4D event come from? does the cascade end at a boundary or dwindle to nothing?) rather than quantitative (the dark sector and hierarchy derivations are the same in all three). We acknowledge this as a philosophical distinction in the model, not a predictive one. A specific implementation of the model would need to specify which interpretation is correct (and would need to compute the energy loss per cycle, which is left to future work). The most defensible position is the diminishing cyclic one: it preserves the cascades scale invariance, gives the cascade a direction, avoids the eternal-return problem, and naturally ends without an ad hoc boundary.

- **Big Rip / phantom energy:** The universe ends when dark energy increases without bound, eventually overcoming all gravitational binding (galaxies, solar system, atoms, etc.) and tearing spacetime apart at a finite time in the future. In standard cosmology, this requires dark energy with equation-of-state  $w < -1$  (phantom energy); current observations suggest  $w \approx -1$ , but small deviations are possible. In our model, dark energy is repulsive (per §2.4, the 4D events antigravity projected to 3+1D inverts once, giving repulsive gravity in 3+1D). The model's default is that the 4D events antigravity is constant in 3+1D frame (because our universe is a brief slice of the 4D events full duration, §2.4), giving  $w \approx -1$  and no Big Rip. However, if the 4D events antigravity is not exactly constant—if it increases over time, or if the dimensional projection gives a time-varying dark energy in 3+1D—then a Big Rip is a natural possibility. The model is agnostic on this question: it does not currently predict whether dark energy is exactly constant or slowly increasing. Testable predictions: Euclid, Roman Space Telescope, Vera Rubin Observatory / LSST, and SKA will measure  $w$  to high precision in the coming decades. If  $w$  is found to be significantly less than  $-1$  (or  $w$  is increasing over time), the model would predict a Big Rip. If  $w$  remains  $\approx -1$  to high precision, the model still allows for fixed-time boundary or cyclic endings, but rules out Big Rip. The Big Rip interpretation is therefore empirically distinguishable from the fixed-time boundary and cyclic interpretations, but the model is not currently committed to any of them. (Note: in the universal bulk-brane cancellation picture, the Big Rip applies at every level—each level could Big Rip if its dark energy is not constant.)
- **Big Freeze / heat death (expansion-driven emptiness):** The universe expands exponentially forever, driven by constant dark energy. Matter density drops as  $a^{-3}$  (where  $a$  is the scale factor), so the universe becomes infinitely dilute over time. Eventually, matter is so sparse that no structures can form or be maintained: galaxies disperse, stars burn out, planets disintegrate, atoms themselves may eventually decay (if proton decay or similar processes occur). The universe ends in a state of infinite emptiness—no matter, no structures, no observers, no events in any meaningful sense. This is sometimes called the Big Freeze (emphasizing the expansion) or heat death (emphasizing the thermodynamic equilibrium). In our model, this is the natural fate of the universe if (1) the 4D events antigravity is constant in 3+1D frame (the model's default per §2.4), (2) the universe has no fixed-time boundary (i.e., the boundary interpretation is wrong), and (3) dark energy is not time-varying (i.e.,  $w \approx -1$  exactly, ruling out Big Rip). The Big Freeze is not an ad hoc postulation in our model—it follows directly from the combination of constant dark energy, no fixed-time boundary, and ordinary thermodynamics. It is the default outcome of standard cosmology with  $\Lambda$ CDM and a cosmological constant; the cascades contribution is to explain why dark energy is constant (because the 4D events antigravity is approximately constant in 3+1D frame, §2.4) and to frame the Big Freeze as one of several possible endings (alongside fixed-time boundary, cyclic, and Big Rip).



Endings at every level. Because the cascade is universal (every level has bulk-brane cancellation, attractive net gravity, similar to 3+1D), the same set of possible endings applies at every level:

- Fixed-time boundary: each level ends at the boundary of its 4D-time slice
- Cyclic: each level Big Crunches, creates a new 4D event
- Diminishing cyclic: each level Big Crunches, creates a smaller 4D event
- Big Rip: dark energy increases, each level tears apart
- Big Freeze / heat death: each level expands forever, becomes empty

The cascade does not predict that some levels heat-death and others Big-Crunch based on parity; it predicts that all levels have similar dynamics, with the specific ending depending on the competition between attractive gravity and repulsive dark energy at that level. Our 3+1D universe is in the late stages of heat death (consistent with standard  $\Lambda$ CDM cosmology with constant dark energy, and with our dark energy winning the competition against matter on cosmological scales). The 2D universes within our 3+1D universe have endings that depend on their size: large 2D universes (from large 3+1D events like AGN or BH-scale) have high matter density that wins against dark energy, leading to Big Crunch and a brief death-flash back-projected to 3+1D; small 2D universes (from small 3+1D events like LHC or small SN) have low matter density where dark energy wins, leading to heat death and a slow diffuse return to 3+1D. Both contribute to the observed dark matter – the Big Crunch case as localized impulsive contributions (currently below detection thresholds), the heat death case as a smooth distributed background. The model is ending-agnostic about which dominates, but predicts that heat death should be common for small 2D universes (where dark energy dominates), which is consistent with the observed smooth dark matter background.

The model is intentionally ending-agnostic. The dimensional-cascade framework does not currently predict which of the five possible endings is the actual fate of our universe. This is a feature of the model, not a gap: the cascade is a framework for the dark sector and gravity's weakness, not a complete cosmological theory that derives the universes endpoint. The specific ending depends on factors that the cascade does not currently specify (e.g., the 4D events full duration in 4D time, the dimensional projections stability, whether dark energy is exactly constant or slightly time-varying, the balance between matter density and dark energy over the 4D events full duration). A specific implementation of the model would need to derive these factors from the cascade geometry to make a definite prediction about the universes end; this is left to future work. The five possible endings are empirically distinguishable by future observatories (Euclid, Roman, LSST, SKA), so the question of which is correct is testable in principle, even if the model itself is currently agnostic. We note that this agnosticism is consistent with the spirit of the cascade as a thought experiment: the model reinterprets the dark sector through dimensional projection, but does not commit to a specific cosmological endpoint. The endpoint is an empirical question that the model frames in a new way (as a question about the 4D events nature), rather than answering it directly.

A further speculation: the recursive cascade and the vacuum bubble analogy. The cascade is not a single-step process from 3+1D to 2D. It is recursive: each 2D universe created by a 3+1D event itself undergoes a Big Bang – expansion – Big Crunch cycle, and the Big Crunch is a high-energy event in 3+1D (not in 2D) – a 3+1D energetic event that the original 3+1D universe experiences as an additional energetic event at the same location. This 3+1D energetic event (the 2D universes Big Crunch, as seen from 3+1D) creates a new 2D universe (a smaller one, since the Big Crunch is smaller than the original 3+1D event). The new 2D universe also undergoes its own Big Bang – Big Crunch, which (as seen from 3+1D) is another 3+1D energetic event at the same location, creating another 2D universe, and so on.

The cascade is fractal: each levels Big Crunch (as observed in 3+1D) is the next levels Big Bang (as experienced in 3+1D). All levels of the cascade happen at the same location in 3+1D, in quick succession each levels timescale is much shorter than the previous (per  $\tau_{2D} = \ell/c$ , the 2D universes lifetime is much shorter than the 3+1D events timescale; the next-level 2D universes lifetime is much shorter than that; and so on). From 3+1D, the entire recursive cascade appears as a single event (or a very brief sequence) at one location, even though within each 2D universes own frame, the Big Bang Big Crunch cycle is a full expansion-contraction.

The vacuum bubble analogy. This is exactly the behavior of a cavitation bubble in water (e.g., from a ships propeller or a focused ultrasound pulse): (1) a 3+1D energetic event nucleates a 2D bubble; (2) the 2D bubble expands, reaches maximum size, and contracts under surface tension; (3) the 2D bubble collapses, producing a sonoluminescence flash (a high-energy event in 3+1D the flash is observed in 3+1D water, not in 2D bubble-world); (4) the flash can nucleate a smaller 2D bubble at the same location; (5) the smaller bubble also expands, contracts, collapses, producing another flash in 3+1D water; (6) and so on, with each level smaller than the previous, until the energy is too small to create a new bubble. All of this happens in microseconds in 3+1D frame, but within the bubbles own 2D world, each level is a full expansion-contraction cycle. The cavitation flash in water is literally the cascade: each levels Big Crunch (observed in 3+1D water) is the next levels Big Bang (observed in 3+1D water). The cascade happens at one location in 3+1D, in quick succession.

Implications for the cascade model. The 3+1D dark matter is the cumulative gravitational back-projection of 2D universes from all 3+1D energetic events (per §2.5, §4.7, §4.10). But each of these 2D universes itself cascades into more 2D universes (smaller, at the same 3+1D location, in quick succession), each of which also has a gravitational back-projection to 3+1D. The total 3+1D dark matter is the sum across all levels of the recursive cascade, weighted by each levels energy and back-projection efficiency. In the models current framing, the first-level 2D universe is the primary contribution to 3+1D dark matter; higher-level (smaller, faster) 2D universe contributions are secondary and likely small in comparison (since each levels energy is a fraction of the previous). The cascade has practical finite depth (even though it is conceptually infinite in principle, per §2.8): it effectively terminates when the energy per level becomes too small to create a new universe (analogous to a cavitation cascade terminating when the bubble is too small to collapse with sufficient energy). The labels (1D-level, 0D-level, -1D-level, ) can in principle extend arbitrarily deep, but in practice the cascade is dominated by the first few levels, and deeper levels are quantitatively irrelevant. The termination is a physical one (energy threshold) rather than a mathematical one (no a priori depth limit).

The 3+1D dark matter picture (refined). The ~27% of the universes mass-energy that is dark matter is the total cumulative back-projection of all cascading 2D universes (across all levels of the recursive cascade) from all 3+1D energetic events. The first-level 2D universes from current stars, supernovae, AGN, and other 3+1D activity dominate this contribution. Higher-level (smaller, faster) 2D universe back-projections are secondary. Big Crunches, if they happen, are also first-level 2D universe creators (since they are 3+1D energetic events), but the diminishing cyclic picture suggests they are a small fraction of total 3+1D activity. The 27% / 5% / 68% split remains the current cycles signature, with the recursive cascade within each 3+1D event contributing to the 27% via the sum of all levels.

Honest caveats: the recursive cascade is a speculative extension of the core model; the papers quantitative claims (§2.5, §4.7, §4.10) are based on first-level 2D universe creation only, and higher-level contributions are not currently derived. The cavitation analogy is qualitative, not quantitative the cascades depth, energy distribution, and back-projection efficiency across levels are not specified. A specific implementation would need to compute the sum across all levels, which is left to future

work.

Observational consequences of the fixed-time boundary are subtle. If the boundary occurs at a specific time in the future, it would not necessarily be visible from within the universe (just as the Big Bang is not visible from within the universe in the conventional sense). Predictions about the boundary are therefore not in the form of we will see X happen but rather in the form of we should not see any signs of a long slow decline as the end approaches. The model is consistent with current observations.

What we do and do not know about the total lifetime. The universes total lifetime  $T_{\text{total}}$  is some fraction of the 4D events full duration in 4D time.  $T_{\text{total}}$  is finite but currently unknown to us. The model is consistent with  $T_{\text{total}}$  being any finite value; the only constraint is that the universe is not yet at its boundary, because we are observing it. The model does not predict when the boundary will occur; it only predicts that there is one.

### 1.3.9 2.9 Why gravity is so constant

A natural prediction of the model is that the gravitational constant  $G$  should be approximately constant over cosmic time. This is not because of any conservation law, but because the bulk-brane cancellation is approximately constant during our universes brief slice of the 4D events full duration. The bulk-brane interaction is a continuous 4D-side process; we are seeing a brief moment of it, during which the near-cancellation between brane gravity and inverted bulk gravity is approximately constant. (Note: this is distinct from the dark energy, which is also approximately constant in our frame, but for a different reason – the dark energy is the 4D events un-cancelled antigravity, which is approximately constant in our brief slice.  $G$  and dark energy are both approximately constant in our frame, but the underlying reason is the same: our universe is a brief slice of the 4D events full duration, during which the 4D-side physics is approximately steady-state.)

This is consistent with observations:  $G$  has been constant to within  $\sim 10\%$  over the age of the universe, and to within  $\sim 10\%$  per year over the last 50 years. The model is consistent with this constancy and would be strained by any detection of significant  $G$  variation.

---

## 1.4 3. Relation to existing work

This proposal builds on several established research programs.

### 1.4.1 3.1 Brane-world cosmology

The brane-world framework [ADD98, RS99, Gregory00] provides the geometric foundation. Our contribution is the specific interpretation of the bulk-brane gravity cancellation as a productive annihilation following a single ongoing energetic event in the bulk (with a finite duration in 4D time), rather than a quiet continuous suppression. From the 3+1 dimensional frame, this ongoing 4D-side process appears as a constant dark energy, because our universe is a brief slice of the 4D events full duration.

### 1.4.2 3.2 Bulk-brane energy exchange

Several authors have studied energy exchange between brane and bulk [Tetradis04, Yousef13]. Our proposal is a specific mechanism for such exchange: a single ongoing energetic event in the bulk (with a finite duration in 4D time), whose antigravity is continuously transferred to the brane

during our universes brief slice of the 4D events full duration. From the 3+1 dimensional frame, this continuous 4D-side transfer appears as a constant dark energy (because our slice is brief). The dark sector (dark matter, dark energy) is the result of this continuous transfer, with the dark matter being a rate-dependent effect (current 2D universe gravity) and the dark energy being a 4D-event-driven geometric contribution.

#### 1.4.3 3.3 Emergent gravity

Verlindes emergent gravity [Verlinde16] proposes that gravity is not fundamental but emerges from quantum information, and that dark matter is the elastic response of this emergent gravity to ordinary matter. Our proposal shares the general spirit of geometric/emergent explanations of dark matter, but is more specific in its dimensional mechanism: we propose a concrete inversion postulate triggered by a single ongoing 4D event. We discuss the relationship between our model and Verlindes framework in more detail in §3.7.

#### 1.4.4 3.4 The Dark Dimension scenario

The recent Dark Dimension proposal [Obied23] argues for a single small extra dimension of size 1-10 m, with a tower of massive spin-2 Kaluza-Klein graviton excitations. The lightest gravitons in this tower can serve as dark matter candidates, decaying over cosmological timescales. This scenario simultaneously addresses the hierarchy problem, the nature of dark matter, and certain cosmological tensions (notably the small-scale structure of dark matter halos). A 2024 follow-up study [LawSmith24] found that astrophysical constraints (CMB distortions from graviton decay) are consistent with the natural parameter range of the scenario. A very recent 2025 paper [Borah25] extends the scenario to allow the dark matter mass to vary as dark energy decreases (Evolving Dark Sector).

Our proposal is structurally similar to the Dark Dimension scenario both invoke a small extra dimension that affects gravities apparent strength and provides dark matter candidates but differs in the specific mechanism: (1) our model uses a single ongoing 4D event (not a fixed small dimension), (2) our dark matter is the collective gravity of 2D universes being currently created (not graviton modes of a fixed dimension), and (3) our dark energy is the un-cancelled bulk gravity (not a separate cosmological constant). The Evolving Dark Sector 2025 idea is closer in spirit to our model than to the original Dark Dimension scenario, in that it suggests dark matter is not a static relic. We discuss this in more detail in §3.7.

#### 1.4.5 3.5 Holographic and AdS/CFT frameworks

The holographic principle and the AdS/CFT correspondence [Maldacena97] describe a 5-dimensional gravitational bulk (4 space + 1 time) as mathematically equivalent to a 4-dimensional quantum field theory on the boundary (in conventional AdS/CFT notation: AdS bulk CFT boundary). In this picture, the bulk gravity is cancelled from the boundary perspective, with only certain residual effects propagating to the boundary. Our proposal is a phenomenological interpretation of this cancellation that emphasizes the productive nature of the bulk-to-brane energy transfer following a single ongoing 4D event (with our universe as a brief slice of the events full duration). The relationship between the conventional AdS/CFT framework and our 4D event proposal is structural rather than direct: both involve a higher-dimensional bulk whose dynamics are not fully accessible from the lower-dimensional brane, with the brane observing only certain projected aspects of the bulk physics. We do not claim that the AdS/CFT correspondence is the correct framework for our

model we claim that the philosophy of the holographic principle (a higher-D bulk cancels from the lower-D perspective) is consistent with our inversion postulate.

#### 1.4.6 3.6 Departure from standard brane-world

We note explicitly that our models inversion claim is stronger than standard brane-world physics. The original ADD and RS frameworks describe gravity as suppressed by geometric dilution (the gravitational coupling on the brane is reduced by the volume of the extra dimensions, or by the warping of the geometry). They do not typically claim that the projection inverts the sign of the gravitational coupling.

Our proposal is therefore not a straightforward extension of ADD/RS. It is a more aggressive geometric claim: the projection not only weakens gravity, it changes its sign, leading to cancellation with brane gravity. The un-cancelled residue of the inverted bulk gravity is what we identify as dark energy (a repulsive effect, consistent with observation).

This stronger claim is more tightly constrained by data. A specific implementation of the inversion mechanism must be checked against sub-millimeter gravity experiments, gravitational wave propagation, and cosmological observations.

#### 1.4.7 3.7 Positioning relative to recent related work

The model in this paper is one of several recent proposals that attempt to unify dark matter, dark energy, and gravity through geometric or extra-dimensional mechanisms. We position this model relative to the most relevant recent work.

Note on framing. The formal framework of this model uses the language of extra dimensions, Kaluza-Klein reduction, and brane-world physics (per §2.1). The physical interpretation is one of nested universes (also per §2.1): our universe is part of a hierarchy of nested universes, with each universe containing child universes and being contained within a parent universe. The extra dimensions provide the mathematical structure; the nesting is the physical content. We use both framings throughout the paper, and we hope this sections positioning will be useful to readers who are familiar with either framing.

Verlindes emergent gravity (2016, ongoing). Erik Verlinde's emergent gravity proposes that gravity is not fundamental but emerges from quantum entanglement, and that dark matter is the elastic response of this emergent gravity to baryonic matter. Our model is structurally similar but makes a specific geometric claim (dimensional inversion following a 4D event) that Verlinde's framework does not. The physics community has been largely skeptical of Verlinde's specific predictions; a 2024 *Ars Technica* article describes emergent gravity as a dead idea, but not a bad one. We acknowledge that our model shares with Verlinde's framework the general spirit of geometric/emergent explanations of dark matter, but differs in: (1) the explicit dimensional-inversion postulate, (2) the scale-invariant principle (every energetic event creates 2D universes), and (3) the current rate-dependent framing of dark matter (dark matter is not a relic, but a current effect). A specific implementation of our model would need to derive testable predictions that distinguish it from Verlinde's framework.

The Dark Dimension scenario (Obied, Dvorkin, Gonzalo, Vafa 2023; Law-Smith, Obied, Prabhu, Vafa 2024; further work 2025). The Dark Dimension scenario proposes a single extra dimension of size  $\sim 1\text{-}10\text{ m}$ , with massive spin-2 Kaluza-Klein gravitons as dark matter candidates, decaying over cosmological timescales. The 2024 follow-up (arXiv:2307.11048) found that astrophysical constraints (CMB distortions from graviton decay) are consistent with the natural parameter range of the scenario. A very recent 2025 paper (arXiv:2507.03090) proposes that the dark matter mass varies as dark energy decreases (Evolving Dark Sector). Our model is structurally similar to the Dark

Dimension scenario both invoke extra dimensions affecting gravity's apparent strength and providing dark matter candidates but differs in the specific mechanism: (1) our model uses a single ongoing 4D event (not a fixed small dimension), (2) our dark matter is the collective gravity of 2D universes being currently created (not graviton modes of a fixed dimension), and (3) our dark energy is the un-cancelled bulk gravity (not a separate cosmological constant). The Evolving Dark Sector 2025 idea is closer in spirit to our model than to the original Dark Dimension scenario, in that it suggests dark matter is not a static relic.

MOND and modified gravity [Desmond25]. Modified Newtonian Dynamics (MOND) modifies the dynamics of visible matter to explain galaxy rotation curves without dark matter. A comprehensive 2025 review [Desmond25] finds that MOND has significant observational successes (especially the RAR) but fundamental failures (CMB power spectrum, galaxy clusters, the Bullet Cluster). The pattern of MONDs success and failure is a cautionary tale for any modified-gravity or geometric dark matter proposal: an individual observational success (like the RAR) is not enough; the model must also explain the broader cosmological context. Our model is consistent with the RAR (more visible mass more activity more dark matter) but has not yet been checked against the CMB power spectrum, galaxy cluster dark matter content, or the Bullet Cluster in detail. A specific implementation of our model would need to address these tests.

CDM with baryonic feedback [Kravtsov24] and others. Standard CDM-based galaxy formation models, with proper treatment of baryonic feedback, can reproduce the RAR and the dark matter content of ultra-diffuse galaxies including DF2 and DF4. This means that the individual observational anomalies our model addresses can also be explained by conventional physics with carefully-tuned baryonic feedback. The model's unique contribution is the geometric unification of dark matter, dark energy, and gravity not the explanation of any individual observation. A specific implementation of the model would need to demonstrate that the geometric unification predicts the baryonic feedback parameters independently, rather than just fitting them.

Other 2024-2025 work. Recent related proposals include various geometric approaches to dark energy (e.g., volume-conservation-based derivations of the cosmological constant) and various holographic/AdS/CFT based explanations of dark energy and dark matter. Our model shares with these proposals the general spirit of geometric explanations but differs in the specific dimensional-inversion mechanism. We do not attempt a comprehensive comparison here.

The competitive landscape. The current theoretical landscape for dark matter/dark energy unification is active but competitive. The most successful framework is still standard CDM with baryonic feedback; modified-gravity proposals have individual successes but face collective challenges; geometric/extra-dimensional proposals (Verlinde, Dark Dimension, this model) are interesting but not yet established. Our model contributes to the geometric-proposal class with a specific dimensional-inversion mechanism and testable predictions (DF2/DF4 correlation with stellar density, the RAR scatter-activity correlation, no direct detection). Whether the model is correct is a question for the community; whether the model is interesting is a matter of taste.

Other 2025-2026 archive submissions. A survey of the open-access archives (ai.viXra.org, rx-iverse.org, and viXra.org) reveals several recent papers exploring conceptually similar ideas, including: a Paired Universe Theory proposing a companion universe whose resistance to stretching generates gravity and dark matter (Godwin, ai.viXra:2606.0008); various dark matter as Weyl curvature proposals; and universe creation in higher dimensions frameworks. These are not direct precursors to the present model (the specific dimensional-cascade-with-sign-flipping mechanism appears to be original), but they illustrate that the general spirit of geometric dark-sector explanations is being explored in multiple directions. We welcome the community to point out any prior work we have missed.

---

## 1.5 4. Predictions and distinguishing features

If the model is correct, several observable consequences follow.

### 1.5.1 4.1 The radial acceleration relation

The radial acceleration relation (RAR) is a tight empirical correlation between the visible (baryonic) mass distribution in galaxies and the total (visible + dark) mass distribution inferred from rotation curves [McGaugh16]. This correlation is not trivially reproduced by simple particle dark matter models with no baryonic feedback (which would predict more variation between galaxies), but can be naturally produced by models in which dark matter is a response to visible matter (such as modified gravity, emergent gravity, or our scale-invariant model). The RAR is also reproduced by standard CDM-based galaxy formation models with proper treatment of baryonic feedback [Kravtsov24], as we discuss further in §3.7.

In the scale-invariant version of this model, dark matter is the collective gravitational signature of all the 2D universes created by energetic events in our 3+1 dimensional world. The current energetic activity of a galaxy – its current star formation rate, current supernova rate, current AGN activity – is strongly correlated with its visible mass: more mass – more stars – more stellar collisions, more supernovae, more AGN activity – more energetic processes – more 2D universes – more dark matter. This strong correlation naturally produces the observed tight correlation between visible mass and total mass in galaxies (the RAR), because the average energetic activity of a galaxy is strongly tied to its visible mass.

Recent RAR results (2024/2025). The RAR has been confirmed and extended by several recent studies. MIGHTEE-HI [Vrteanu25] confirmed the RAR with a large new sample using resolved stellar mass measurements. [Mistele24] combined kinematic and weak-lensing data to extend the RAR over a large dynamic range, with consistent results. However, recent studies have also identified deviations from a single universal RAR:

- The EDGE collaboration [Júlio25] found that low-mass dwarf galaxies ( $M_{\text{bar}} \sim 10 M_{\odot}$ ) lie systematically above the low-mass extrapolation of the RAR – meaning the RAR is not a single universal function at low masses.
- [Mercado24] found that the RAR has subtle hooks and bends in its shape, not a single smooth function.
- [Tian24] found that Brightest Cluster Galaxies (BCGs) follow a different RAR from typical spirals.

These results show that the RAR is approximately tight, but not perfectly universal across all galaxy types and mass ranges. The RARs tightness at intermediate masses ( $10^8 - 10^{12} M_{\odot}$ ) is the most robust feature; deviations at low masses and in BCGs are now well-established.

Implications for our model. Our model is qualitatively consistent with the RARs tightness at intermediate masses: more visible mass – more activity on average – more dark matter. The model's additional prediction is that the small scatter in the RAR at fixed visible mass should correlate with current activity, which is testable but not yet definitively tested. The recent deviations from a single universal RAR (dwarfs above the extrapolation, BCGs on a different relation) are not directly predicted by the model in its current form – but the model could potentially accommodate them by allowing the proportionality between activity and dark matter to vary with galaxy type or mass. A

specific implementation of the model would need to derive the RARs exact shape and the source of its deviations to be a quantitative match to the data.

We also note that the RAR is not uniquely a signature of modified gravity or of our model. Standard CDM-based galaxy formation models with proper treatment of baryonic physics (e.g., [Kravtsov24]) can reproduce the RAR. The RAR is therefore a necessary feature of any successful model, not a sufficient test of our model specifically.

At fixed visible mass, the model predicts that the small scatter in the RAR should correlate with the current activity of each galaxy: galaxies with more current activity (relative to their mass) should have more dark matter (and therefore higher total mass at fixed visible mass). This is a sharp, testable prediction that distinguishes our model from standard particle dark matter, which would predict that the RAR scatter is determined by halo formation history, not by current activity. The model is consistent with the observed tightness of the RAR ( $\sim 0.1$  dex scatter) at intermediate masses, because the strong correlation between visible mass and average activity dominates over the smaller variation in current activity at fixed visible mass. The models additional prediction is that this small scatter, in our model, should correlate with current activity a subtle effect that could be tested with high-precision data.

Why dark matter is only observable on galaxy scales. The RARs existence reinforces why dark matter is not directly detectable in stellar-scale or sub-stellar-scale environments. In the model, dark matter is the cumulative gravitational effect of 2D universes being created throughout a region of space. For a galaxy-sized region, this cumulative effect is substantial (it produces the observed rotation curves). For a stellar-sized or planetary-sized region, the cumulative effect is too small to detect because the local activity (e.g., solar fusion, geothermal activity) is dwarfed by the cumulative activity of the surrounding galaxy. The dark matter density at the Suns location, in our model, is set by the galaxy's rate of large-event creation (supernovae, AGN), not the Suns rate of small-event creation. The Suns own activity adds a perturbation that is far below any detectable level. This is why direct-detection experiments (looking for dark matter particles) have all returned null results: the dark matter is smeared out by the cumulative activity of the entire galaxy, with no locally-detectable signature at any specific location. The RAR is the only scale on which dark matter becomes measurable, because galaxy scales are where the cumulative effect is large enough to be observable.

#### 1.5.2 4.2 Dark matter as current collective gravity, not a relic

Standard WIMP dark matter models predict that dark matter is a relic of the early universe, with density fixed by freeze-out and subsequently diluted only by cosmic expansion. In our model, dark matter is not a static relic it is the current collective gravitational signature of 2D universes being created right now by 3+1 dimensional energetic events. The local dark matter density at any point in our 3+1 dimensional frame is determined by the current rate of 2D universe creation in that region, weighted by the energy of each event.

The dark matter at a point is the cumulative gravitational effect of all 2D universes being created now in that region, projected into our 3+1 dimensional frame. The dark matter density at that point is therefore proportional to the current event rate in that region, weighted by event energy. The dark matter density is not a constant; it varies with the local event rate, and (in the same way as ordinary matter density) it is also diluted by cosmic expansion.

This means dark matter is not an ancient residue that has been accumulating since the Big Bang. It is a current effect, with the current rate of 2D universe creation setting the local dark matter density. The model predicts that dark matter density should track the current event rate, including any cosmic evolution of stellar activity. This is a testable prediction: if dark matter density has



decreased over cosmic time in step with the decline in star formation rate (after accounting for the standard dilution by cosmic expansion), that would be evidence for the model. (Note: this is a subtle effect, since the dark matter density in halos is also affected by cosmic expansion and dynamical evolution, which would have to be disentangled from the model prediction.)

The stability of dark matter density on short timescales (within a galaxy's lifetime) is a consequence of the local event rate being approximately constant on those timescales. On cosmological timescales, the model predicts two effects on dark matter density: (a) standard dilution by cosmic expansion (decreasing density as the universe expands, just like ordinary matter), and (b) a weak evolution of dark matter density correlated with cosmic star formation history. The two effects are different in nature: the first is a geometric dilution, the second is a rate-dependent effect specific to this model.

**Total amount of dark matter.** The model implies that the total amount of dark matter in a comoving volume (a region of the universe expanding along with cosmic expansion) is set by the current event rate  $\propto$  the average lifetime of 2D universes in our frame, not by a time-integrated history. Each 2D universe contributes to dark matter for its brief lifetime in our frame, then its contribution ends (it has completed its cosmic history in 2D). The total dark matter in a comoving volume is therefore steady-state in equilibrium: as old 2D universes end, new ones are being created at a comparable rate. On cosmological timescales, the total amount of dark matter in a comoving volume is approximately constant if the average event rate per galaxy is approximately constant, since the comoving volume contains a roughly constant number of galaxies. This is similar to standard cosmology, where the total amount of dark matter in a comoving volume is also approximately conserved.

The dark matter density in a comoving volume, however, does change in this model in two ways: (a) standard dilution by cosmic expansion (decreasing as  $1/V$  as the universe expands), and (b) a weak evolution correlated with cosmic star formation history (the average event rate per galaxy has declined over cosmic time as star formation has decreased). The two effects work in the same direction – both decrease the dark matter density over cosmic time. The model predicts that the dark matter density at high redshift ( $z > 2$ ) was somewhat higher than the standard  $1/V$  dilution would predict, because the average event rate per galaxy was higher then. (Note: the total dark matter in a comoving volume is approximately conserved, but the density decreases because the volume increases – standard cosmic dilution.)

This is a subtle but testable prediction: comparing dark matter densities in galaxies at different redshifts (after accounting for standard cosmic dilution) should reveal a residual correlation with the cosmic star formation history. The effect is small (perhaps a factor of a few) and would require careful measurements to detect.

### 1.5.3 4.3 Dark energy equation of state

In standard cosmology, dark energy is treated as a true cosmological constant – a fixed vacuum energy whose equation of state  $w = p/\rho$  is exactly  $-1$ . Current observations are consistent with this to high precision (the dark energy equation of state parameter  $w$  is consistent with  $-1$  to within a few percent).

In our model, dark energy is the un-cancelled fraction of the inverted bulk gravity ( $\bar{g}_{2.4}$ ) – a contribution from the 4D event, approximately constant in our 3+1 dimensional frame because our universes' lifetime is a brief slice of the 4D events' full duration, during which the 4D events' antigravity output is approximately constant. The dark energy density is therefore approximately constant in our frame (matching standard CDM behavior), and the total dark energy grows as the universe expands (because the universes' volume grows while the density stays constant).

This is similar to standard CDM in its observable consequences: dark energy density is approximately constant ( $w = -1$ ) over cosmic time. The model does not currently predict a detectable deviation from standard CDM in dark energy observations. The distinction between our model and CDM is in the interpretation of why dark energy is constant (the 4D event is in a brief steady state during our slice), not in the observable dark energy behavior.

The model does not currently specify how the dark energy density would evolve over time, because the model does not specify how the 4D events antigravity output evolves over its full duration. The 4D events antigravity output could be increasing over its full duration (the 4D event intensifies in 4D), decreasing (fades in 4D), or constant – the model does not specify. A specific implementation of the model would need to derive the temporal profile of the 4D events antigravity output. The key observation is that in our 3+1 dimensional frame, the dark energy density appears approximately constant regardless of the 4D events long-term behavior – because our universe’s lifetime is a brief slice of the 4D events full duration, during which any 4D-side variation is too slow to detect. If the 4D events output is exactly constant over its full duration, the dark energy density in our frame is exactly constant (matching CDM). If the 4D events output varies slowly over its full duration, the dark energy density would vary slowly (correspondingly, in our frame), but the effect would be much smaller than what we can detect during our brief slice.

Why we do not derive the absolute dark energy density. A natural question: given the cascade, can we derive the absolute value of the dark energy density ( $\sim 10^{-27}$  GeV)? The honest answer is no – at least not without further input. The cascade gives a qualitative explanation of why the dark energy is small (its a near-cancellation residue), and it gives a quantitative prediction modulo the staying fraction  $f_{back}$  (§2.6):  $\rho_{DE} \sim f_{back} \cdot \epsilon \cdot M_{Pl}^4$ , where  $\epsilon \sim 10^{-38}$  is the bulk-brane cancellation factor and  $f_{back} \sim 10^{-85}$  is the staying fraction. The product matches observation:  $f_{back} \cdot \epsilon \cdot M_{Pl}^4 \sim 10^{-85} \cdot 10^{-38} \cdot M_{Pl}^4 \sim 10^{-123} M_{Pl}^4 \sim 10^{-47}$  GeV. The individual values of  $\epsilon$  and  $f_{back}$  are postulates of the model, not derivations. A complete implementation of the model would derive  $\epsilon$  and  $f_{back}$  from the geometry of the dimensional projection, which would in turn predict the absolute dark energy density from first principles. We do not claim to have done this derivation in the present paper. We note that the dark energy density is consistent with the cascade-plus-staying-fraction picture for the specific values  $\epsilon \sim 10^{-38}$  and  $f_{back} \sim 10^{-85}$ , but these values are not predicted by the model. An earlier version of this section (in v1.57 and earlier, removed in v1.58) attempted to derive the dark energy density from a dimensional transition threshold  $\lambda_{th}$ , but this attempt failed for internal-numerical reasons (the threshold value that matches dark energy,  $\lambda_{th} \sim 10^{-4}$  m, was inconsistent with the Sun-neutrino constraint that defined the threshold range). The threshold mechanism has been removed in the present version, and the dark energy density is not derived. We acknowledge this as a limitation of the current model: it is qualitatively consistent with observations and provides a unified geometric framework for the dark sector, but it does not yet quantitatively derive the absolute value of the dark energy density. The qualitative picture is robust; the quantitative value is set by the cascade + staying fraction postulate.

A note on the Hubble tension. The Hubble tension is the statistically significant disagreement (currently  $\sim 5\%$ ) between the Hubble constant  $H_0$  measured locally ( $H_0 \approx 73$  km/s/Mpc, from Cepheids and supernovae) and the value inferred from the cosmic microwave background using CDM ( $H_0 \approx 67$  km/s/Mpc, from Planck). The local measurement is higher than the early-universe extrapolation, even after accounting for the known accelerating expansion (which is built into CDM via dark energy with  $w = -1$ ). This tension is one of the most active puzzles in modern cosmology. The dimensional-cascade framework offers a potential connection: if the 4D events antigravity output varies over its full duration (per the acknowledgment above), then the early-universe antigravity and the late-universe antigravity could be slightly different. The dimensional time-dilation principle (§2.3) says the projection from 4D to 3+1D is not a simple linear time translation, so

the effective  $H_0$  at different cosmic times could differ from the CDM-extrapolated  $H_0$  in a way that reduces the tension. Specifically, if the 4D events antigravity output was slightly higher in the early universe than now, the CMB-inferred  $H_0$  would shift upward, reducing the gap with the local measurement. This is a speculative extension of the model – §4.3 already acknowledges that the antigravity output could vary, but the specific temporal profile (and whether it would explain the magnitude of the Hubble tension,  $\sim 6$  km/s/Mpc) is not derived. A specific implementation of the model would need to (a) derive the temporal profile of the 4D events antigravity, and (b) check that the resulting shift in  $H_0$  matches the observed tension. The dimensional-cascade framework is therefore qualitatively compatible with a Hubble tension resolution via time-varying antigravity, but the quantitative details are left to future work. We note that this is a natural connection that could distinguish the model from standard CDM: CDM predicts a strictly constant dark energy (no time variation), while the dimensional-cascade model allows (and may require) slight time variation over the 4D events full duration, which would be a qualitatively different prediction.

#### 1.5.4 4.4 Sub-millimeter gravity tests

If the geometric suppression of gravity depends on the size and shape of the extra dimensions, then gravity should deviate from  $1/r^2$  at length scales comparable to the size of the extra dimensions. Standard ADD-style predictions have been constrained by sub-millimeter gravity experiments (no deviation from  $1/r^2$  down to  $\sim 10$   $\mu$ m has been observed). The dimensional inversion in our model does not by itself predict a specific size for the extra dimensions, so the sub-millimeter constraint does not directly apply to the inversion part of the model – but if the model includes ADD-style geometric suppression as part of its mechanism, then the extra dimensions would need to be smaller than  $\sim 10$   $\mu$ m to be consistent with experimental constraints. The model is consistent with extra dimensions smaller than the experimental reach (e.g., at the Planck scale), but would be in tension with extra dimensions larger than  $\sim 10$   $\mu$ m unless the geometric-suppression aspect of the model is modified. Further theoretical work is needed to extract the model's specific prediction for short-range gravity.

#### 1.5.5 4.5 The Big Bang as a 4D event CMB constraints

If the Big Bang is the projection of a 4D event into our 3+1 dimensional brane, the energy spectrum and spatial structure of the early universe would be set by the projection of that event's spectrum and structure. The cosmic microwave background (CMB) power spectrum, the abundances of light elements from Big Bang nucleosynthesis, and the early-universe particle production all depend on the initial conditions. The CMB has been measured to extraordinary precision by the Planck satellite and is consistent with a nearly scale-invariant primordial power spectrum, a radiation-dominated early universe, and  $N_{\text{eff}} \approx 3.035$  relativistic species at recombination.

A specific implementation of the 4D event scenario must reproduce these observations. The model in its current form does not derive the spectral shape or the spatial structure from first principles – we have not specified the bulk field content, the event's energy, the projection rule, or the spatial structure of the 4D event. We note that:

- The early-universe energy spectrum in our 3+1 dimensional frame is set by the projection of the 4D event's energy into our 3+1 dimensional brane. The actual spectrum would be set by the higher-dimensional physics of the original event and the geometry of the dimensional projection.
- The spatial structure of the 4D event determines the initial conditions for structure formation in our 3+1 dimensional universe. The observed near-scale-invariance of the CMB power spectrum requires the 4D event to have nearly-homogeneous energy density on the largest

scales (with small fluctuations that project as the seed perturbations for cosmic structure). This is a strong constraint on the 4D event: it must be spatially extended and approximately homogeneous (in 4D), not a localized point-like event. This is consistent with the model: the 4D event is described as having a spatial extent, and that spatial extent could be very large compared to the Planck scale, with the energy density approximately uniform across that extent.

- A localized 4D event with highly non-uniform energy density would project to a highly non-uniform early universe, in conflict with the observed CMB. The model therefore requires the 4D event to be spatially extended and approximately homogeneous. This is an additional constraint on the 4D event that was not previously emphasized in this paper.
- The standard  $\Lambda$ CDM cosmology (with inflation) is in excellent agreement with current CMB data. The 4D event scenario must do at least as well, with any deviations being a target for observational test. The model does not currently explain the origin of the primordial perturbations (the inflationary quantum-fluctuation picture is one possibility; another is that the 4D event had its own small-scale structure that projects as the seed perturbations).

Inflation, matter-antimatter asymmetry, and other open issues. The dimensional-cascade framework does not currently derive: - Cosmic inflation the near-exponential expansion in the very early universe ( $\sim 10^{-36}$  to  $10^{-32}$  seconds after the Big Bang) that solves the horizon, flatness, and monopole problems. In the cascade, the 4D events projection could in principle provide an inflation-like phase (if the 4D event had a brief, intense early phase that projected to 3+1D as accelerated expansion), but this is not derived in the current model. A specific implementation would need to derive the inflationary phase from the 4D events temporal/spatial structure, and check that the resulting primordial perturbation spectrum matches observations (nearly scale-invariant,  $n_s \approx 0.965$ , with no detectable tensor modes at current sensitivity). - Matter-antimatter asymmetry the observed fact that our universe has more matter than antimatter (baryon-to-photon ratio  $\eta \sim 6 \times 10^{-10}$ ). The cascade does not currently explain this asymmetry. In standard cosmology, the asymmetry is generated by baryogenesis (Sakharov conditions: baryon number violation, C and CP violation, out-of-equilibrium processes). In the cascade, the 4D event could in principle generate the asymmetry (if the 4D events projection preferentially created matter over antimatter, or if the dimensional projection inherently violates C and CP), but this is not derived. A specific implementation would need to address why the projected 3+1D universe is matter-dominated. - Big Bang nucleosynthesis (BBN) the observed light element abundances (D,  $^3\text{He}$ , He, Li) at  $\sim 10^{-2}$  to  $10^3$  seconds after the Big Bang, which constrain the baryon-to-photon ratio and the number of relativistic species. The cascade does not currently derive the BBN predictions from the 4D event; the model takes the standard BBN picture as given and notes that the 4D event scenario must be consistent with the observed light element abundances. - Primordial black holes, topological defects, cosmic strings other features of standard cosmology that are not currently addressed by the cascade.

These are honest gaps in the current model. The cascade is a framework that addresses the dark sector (dark matter, dark energy, gravity's weakness) but does not yet derive the full set of standard cosmological predictions. A complete implementation of the cascade would need to address all of these issues, but the current paper focuses on the core dimensional-cascade model and the dark sector, leaving the broader cosmological implications for future work.

This is a target for theoretical development.

1.5.6 4.6 (Section removed: neutrino mass is a Standard Model physics question, not addressed by this model.)

1.5.7 4.7 Dark matter density should correlate with energetic event rates on galaxy scales

If dark matter is the collective gravitational signature of 2D universes being currently created by 3+1 dimensional energetic events, then the local dark matter density in a region of space should be proportional to the current rate of energetic events in that region, weighted by the energy of each event. This correlation is expected to manifest on galaxy scales (or larger), not on stellar or sub-stellar scales.

Why event size matters, not just event rate. The relevant quantity for dark matter production is not just the count of events but also their energy. Each 2D universe created by a 3+1 dimensional event has a gravitational contribution proportional to the events energy. Many small events (e.g., solar fusion reactions at  $\sim$ MeV each) contribute little to dark matter per event, even at high rates. A few large events (e.g., supernovae at  $\sim 10^9$  eV each) contribute much more per event.

The Sun, for example, hosts  $\sim 10^8$  nuclear fusion reactions per second an enormous event rate in absolute terms. Each event releases only  $\sim$ MeV, so the current power output from solar fusion is  $\sim 3.8 \times 10^8$  W. By contrast, a single supernova releases a total of  $\sim 10^{44}$ - $10^{45}$  ergs of energy ( $10^8$ - $10^9$  eV in kinetic energy plus neutrinos;  $\sim 10^{44}$  ergs  $10^9$  eV (since  $10^9$  eV =  $1.6 \times 10^8$  erg) in visible light, which is what an external observer primarily sees) in a single brief event. The supernova energy depends on the type: Type Ia releases  $\sim 10^{44}$  ergs of kinetic energy, while Type II releases  $\sim 10^8$  ergs total (mostly neutrinos). Using the visible-light energy of  $\sim 10^9$  eV as the energetic event energy (since most of the kinetic and neutrino energy does not directly create 2D universes via 3+1D electromagnetic interactions), the supernovas event energy is  $\sim 10^9$  eV. Numerical correction: an earlier draft of this section said  $10^9$  eV ( $10^8$  ergs) the  $10^9$  eV is correct (visible-light energy), but  $10^9$  eV =  $1.6 \times 10^8$  ergs, NOT  $10^8$  ergs. The discrepancy is because the total supernova energy is  $\sim 10^8$  ergs, but most of that is kinetic and neutrino energy, not visible light. The visible-light energy of  $\sim 10^9$  eV is what primarily drives the 2D universe creation in our 3+1D frame (since neutrinos and bulk kinetic energy do not directly create 2D universes via 3+1D events). For comparison, this is  $\sim 0.1\%$  of the Suns total output over its entire lifetime ( $\sim 1.2 \times 10^{44}$  J =  $1.2 \times 10^{45}$  ergs), not  $\sim 1\%$  as a previous draft claimed. The Milky Ways current supernova rate is  $\sim$ few per century, but each event contributes much more dark matter per event than solar fusion. The galaxy's current energetic activity (per unit volume) is therefore dominated by its large events (supernovae, AGN), not by stellar fusion.

(Note: the model does not depend on the cumulative energy released over cosmic history. The dark matter at any point is set by the current event rate, not the historical rate. The historical comparison here is for intuition about the relative importance of small vs. large events in the current activity budget.)

Why the Suns local dark matter isnt enhanced. In the model, each 2D universes gravitational contribution is proportional to its creating events energy. The Suns many small fusion events create 2D universes with small gravitational contributions. The galaxy's rare large events create 2D universes with large gravitational contributions. The dark matter density at the Suns location is set by the galaxy's rate of large-event creation, not the Suns rate of small-event creation. The Sun sits within the galaxy's dark matter halo, but the Suns own fusion adds a perturbation that is far below any detectable level. This is consistent with observational constraints from direct-detection experiments and solar dark matter capture arguments.

The galaxy-scale prediction. Two galaxies of the same total stellar mass but different stellar densities should have different current energetic activities, and therefore different dark matter

content:

- A dense galaxy has a higher rate of supernovae (per unit stellar mass), more black hole formation, more AGN activity. Its current energetic activity is high.
- A diffuse galaxy has a lower rate of supernovae (per unit stellar mass), less AGN activity, less energetic processing. Its current energetic activity is low.
- The model predicts: more current activity = more dark matter.

This is the testable galaxy-scale prediction: holding total stellar mass fixed, galaxies with higher stellar density (and therefore higher current activity) should have more dark matter.

What this rules out. The model does not predict: - Enhanced dark matter density in the Sun's interior due to solar fusion - Enhanced dark matter density near nuclear reactors - Enhanced dark matter density near particle accelerators - Enhanced dark matter density in the Earth's core due to geothermal activity

The Sun's cumulative activity is small compared to the galaxies, and stellar-scale activity in general does not produce a measurable local enhancement. The proportionality constant is too small for these stellar-scale and sub-stellar-scale effects to be detectable.

On the every event question. A natural concern: does the model require that every energetic event including every photon emission, every atomic transition create a 2D universe? The simplest reading is that all energetic events contribute, with each event's contribution weighted by its energy: small events (atomic transitions, photon emissions) create 2D universes with small gravitational contributions, while large events (supernovae, AGN outbursts) create 2D universes with large gravitational contributions. The cumulative effect is the sum over all events, weighted by energy. This is consistent with the radial acceleration relation (§4.1): the RAR is tight (0.1 dex scatter) because the average activity per unit visible mass is approximately the same across galaxies of similar visible mass, even though the specific event distributions differ. Dark matter correlates with the average activity, not with the specific event history. This is why the RAR works in this model: more visible mass more activity on average more dark matter. The §4.8 discussion of diffuse galaxies uses this framing: the rate of both small and large energetic events is proportional to the particle number density, so diffuse galaxies (low density) have low average activity, and therefore low dark matter.

On experimental feasibility. Dark matter has been measured gravitationally in many ways galaxy rotation curves, gravitational lensing, CMB power spectrum, large-scale structure, the Bullet Cluster but it has not yet been directly detected as a particle. Direct-detection experiments (XENON, LUX, LZ, PandaX) have not observed dark matter particles, and indirect-detection experiments (Fermi, IceCube) have not confirmed dark matter annihilation or decay signals.

The galaxy-scale correlation prediction is testable with existing data from galaxy surveys. A study comparing the dark matter content of mass-matched galaxy pairs with different stellar densities (e.g., compact dwarf spheroidals vs. ultra-diffuse galaxies of the same mass) would be a direct test. The cumulative energetic activity can be estimated from the galaxy's current star formation rate, current supernova rate, and current AGN activity (or, as a proxy, from its stellar density holding mass fixed, since denser galaxies have higher current event rates per unit mass). The model predicts a positive correlation between current activity and dark matter content.

The upcoming Vera Rubin Observatory's Legacy Survey of Space and Time (LSST) will provide unprecedented data on galaxy properties, including stellar densities, star formation histories, and dark matter content inferred from lensing and kinematics. The model predicts that, in a sample of mass-matched galaxy pairs, the more dense galaxy should have more dark matter.

We acknowledge that the proportionality constant for the galaxy-scale correlation is not yet specified. Computing this constant requires a specific geometry and event spectrum, which we have

not derived. The current observational data is qualitatively consistent with the prediction (diffuse galaxies do seem to have less dark matter, and active galaxies tend to have more), but a quantitative test has not been performed.

#### 1.5.8 4.8 Diffuse galaxies and the dark matter / activity correlation

A particularly clean application of the model is the observed low dark matter content in some diffuse galaxies. The most famous cases are NGC 1052-DF2 and NGC 1052-DF4, ultra-diffuse galaxies (UDGs) discovered in 2018 and 2019 that appear to have very little (or no) dark matter halo, in apparent contrast to standard CDM predictions that every galaxy should host a substantial dark matter halo.

Current status of the data. Multiple high-resolution spectroscopic studies have confirmed the anomalously low dark matter content of DF2 and DF4. A 2024 ultra-deep imaging study of DF2 and DF4 [Golini24] found faint tidal tails around DF4, providing evidence that tidal stripping by the nearby massive galaxy NGC 1035 is removing the dark matter from DF4. Both DF2 and DF4 are satellite galaxies of the larger NGC 1052 group, which means their low dark matter content may be environmental (a consequence of tidal interactions) rather than an intrinsic property of diffuse galaxies. A more recent candidate for a dark-matter-free dwarf is FCC 224 in the Fornax Cluster (discovered 2024), which would be a separate test case outside the NGC 1052 group. A 2024 study [Kravtsov24] showed that CDM-based galaxy formation models, with proper treatment of baryonic physics, can reproduce the dark matter content of UDGs including DF2 and DF4 suggesting that the DF2 is dark-matter-free anomaly may be less anomalous than originally thought. There is currently no consensus on the cause of the UDG dark matter deficit; candidate explanations include tidal stripping, modified gravity (MOND,  $f(R)$  gravity), baryonic feedback, and self-interacting dark matter.

Our models explanation. Our model offers a natural explanation that does not require tidal stripping: low dark matter in diffuse galaxies is a consequence of their low average energetic activity per unit volume. The chain of reasoning is:

1. Dark matter, in our model, is the current collective gravitational signature of 2D universes being created by 3+1 dimensional energetic events, weighted by the energy of each event.
2. The rate of energetic events per unit volume is proportional to the number density of particles (because the rate of collisions, decays, atomic transitions, and photon emissions all scale with the local particle density more particles in a region means more events of all kinds per unit time).
3. The rate of large energetic events per unit volume (supernovae, AGN outbursts) is set by the stellar density and the central black hole activity not by the total stellar mass. The average event energy per unit volume is also higher in denser galaxies.
4. Diffuse galaxies have low number density their stars and gas are spread out over a much larger volume than typical galaxies of the same total mass.
5. Therefore, diffuse galaxies have low rates of both small and large energetic events per unit volume, and low average event energies per unit volume.
6. Therefore, diffuse galaxies have low rates of 2D universe creation, weighted by event size, and hence low dark matter densities.

In short: more spread out means less energetic events because less collision, and therefore less dark matter. The model expects ultra-diffuse galaxies like DF2 to have low dark matter content.

What distinguishes our model from the tidal-stripping explanation. The tidal-stripping explanation (the dominant current interpretation) predicts that isolated UDGs (those without nearby

massive neighbors) should have normal dark matter content because there is no tidal force to strip it. Our model predicts that all diffuse galaxies should have low dark matter content, regardless of environment, because the low dark matter is a consequence of the galaxy's own low average activity per unit volume. This is a cleaner test of the model: identify a bona fide isolated UDG (no massive neighbor within several galaxy radii), measure its dark matter content, and compare to similar-mass non-UDG galaxies. Tidal stripping predicts normal dark matter; our model predicts low dark matter.

Other model-distinguishing predictions. The model also predicts that:

- Standard particle dark matter predicts that dark matter content correlates primarily with total stellar mass (more mass more dark matter halo).
- Our model predicts that dark matter content correlates with stellar density (or equivalently, with surface brightness and collision rate), not just total mass. Two galaxies of the same total mass but different densities should have different dark matter content in our model, with the more diffuse galaxy having less.

A direct observational test: select pairs of galaxies matched in total stellar mass but differing in stellar density (e.g., a compact dwarf vs. an ultra-diffuse galaxy of the same mass). Measure their dark matter content via rotation curves, velocity dispersions, or gravitational lensing. Standard CDM predicts similar dark matter content for mass-matched galaxies. Our model predicts less dark matter in the more diffuse galaxy. The same prediction would also hold for mass-matched pairs where one galaxy is in a dense environment and one is in a void (in our model, the void galaxy has less dark matter; in standard CDM, the environment should not matter as strongly).

This is testable with existing data from galaxy surveys (SDSS, DES) and would be a particularly clean test with upcoming data from LSST and Euclid. The correlation between surface brightness and dark matter content holding stellar mass and environment fixed is a sharp, model-distinguishing prediction.

Connection to other anomalies. The same logic applies to several other observed dark matter anomalies:

- Low surface brightness galaxies in general tend to have less dark matter than their high-surface-brightness counterparts of similar mass. Standard CDM has difficulty with this; our model predicts it as a natural consequence of the lower collision rates in diffuse systems.
- Dwarf spheroidal galaxies are often old, low-activity systems with low dark matter content. Our model expects this.
- Galaxies in cosmic voids (sparse regions of the universe) have low overall activity and may have less dark matter than galaxies in dense regions. Our model predicts this.

In each case, the energetic activity of the system (collision rate, star formation rate, gas dynamics) should correlate with its dark matter content. This is a single principle applied across multiple systems.

#### 1.5.9 4.9 Philosophical: dimensional structure and the block universe

This subsection is a philosophical interpretation, not a physical prediction. We include it for completeness, with the explicit understanding that it is interpretive rather than predictive.

If the proposed dimensional structure is real, then a hypothetical 4D observer would experience our 3+1 dimensional universe as a 4D structure in which our time is a spatial direction. From this perspective, the entire history of our universe is a static 4-dimensional structure the projection of



the 4D event laid out in space rather than time. This is the block universe interpretation of special relativity, extended to a 4D bulk perspective.

We note that this is a philosophical position, not a physical prediction. The block universe interpretation is debated within physics and philosophy of physics; many physicists accept it, many do not. It is not testable in the usual sense, and it is independent of the empirical content of the main model.

We include it because the dimensional structure implied by the model invites this kind of geometric reflection, but we explicitly do not claim that it is a prediction of the model.

#### 1.5.10 4.10 Speculative extension: black holes as windows into 4D

This subsection is a speculative extension, not a core claim of the model. We include it as a possible connection between the dimensional-cascade framework and black hole physics, with the explicit understanding that it is exploratory and not derived.

Black holes as voids in 3+1D space, or as 3+1D tears. A natural extension of the model is to consider black holes as regions where 3+1D spacetime has a void—the actual content of the black hole exists in 4D, not in 3+1D. From our 3+1D perspective, we observe the event horizon (the boundary of 3+1D geometry) and infer the singularity (the boundary of 3+1D itself). The gravity we attribute to the black hole is the projected gravity of the 4D content, in the same way that the 3+1D universes gravity is the projected gravity of the 4D event.

A complementary interpretation, which may be more aligned with the mainstream view of black holes as regions of extreme mass concentration, is to think of 3+1D space as having a kind of surface tension—it can be stretched and curved, but only up to a tear threshold. Beyond this threshold, 3+1D tears or opens into 4D, and the stuff inside the black hole is in 4D. In this view, the mass/energy concentration causes the curvature, and the excessive curvature is what causes the dimensional transition. The mass is the cause of the tear, but the tear itself is a structural failure of 3+1D space—not an infinite-density singularity in 3+1D.

Both interpretations are consistent with the dimensional-cascade framework. The void view is more radical; the tear view is more aligned with the mainstream view of black holes as mass-concentration-driven. In either case, the boundary of 3+1D spacetime is somewhere inside the event horizon, and the stuff of the black hole is in 4D. This is a speculative resolution of what the singularity might be, and is not derived from the model.

Information preservation. The black hole information paradox (does information that falls into a black hole survive?) is resolved in this interpretation: the information is not lost because it is not actually in 3+1D to begin with. The information is in 4D, where it can persist indefinitely. Hawking radiation would be the return of 4D information to 3+1D, leaking through the boundary. This is one of several proposed resolutions of the information paradox (others include holographic principle, ER=EPR, and firewall proposals), and it fits naturally with the dimensional-cascade framework.

A note on interior vs. exterior. Throughout this subsection, we use black holes are in 4D as shorthand for a more precise statement: the interior content of a black hole (the singularity, the stuff that has fallen in) is in 4D, while the exterior of the black hole (the event horizon, the gravitational field, the 2D universes created by the black holes energetic processes) is in 3+1D. The 2D universe creation associated with black holes happens at the event horizon (a 3+1D region), not at the singularity (a 4D region). This distinction resolves the apparent tension between this subsection (which says black holes create 2D universes) and the complete dimensional transition at the event horizon (where matter transitions fully to 4D): the content is in 4D, but the boundary is in 3+1D, and 2D universe creation is a boundary effect.

Time dilation as a dimensional effect. The gravitational time dilation observed near black holes

(well-established in general relativity) is given a new interpretation in this view: the time-dilation is because the clock near the black hole is partially in 4D space, where its causal structure is different from the 3+1D causal structure outside the event horizon. A clock near a black hole ticks slower in our 3+1D frame because part of its causal structure is in 4D, and the 4D-side dynamics are not fully projected into 3+1D. This is consistent with the dimensional time-dilation principle of §2.3: a brief moment in one frame can correspond to a vast duration in another, because the dimensional projection maps a short 4D duration to a long 3+1D duration (or vice versa, depending on the projection factor). The black holes interior (4D) and exterior (3+1D) experience different effective time scales, with the 4D interiors physical processes appearing in 3+1D as vastly dilated (i.e., the 4D process completes in brief 4D time, but projects to a very long 3+1D time). This is not because 4D time is faster than 3+1D time – that phrasing is misleading. It is because the rate of 4D-side processes, as observed from our 3+1D frame, is vastly slower than the rate of the same processes in 4D itself. The dimensional time-dilation factor between 4D and 3+1D is huge, which is why black hole evaporation takes  $\sim 10$  years for a solar-mass black hole: from the 4D frame, the evaporation is fast (relative to the 4D events full duration), but from our 3+1D frame, it is vastly slow because the dimensional time-dilation between 4D and 3+1D is vast.

Hawking radiation as diluted 4D energy. A natural extension: Hawking radiation is not a curved-spacetime quantum tunneling effect (as in standard semiclassical QFT on curved spacetime); it is actual 4D energy leaking through the dimensional boundary, but dilated by the dimensional time-dilation factor. Specifically, the true 4D energy of the black hole is some value  $E_4$ , and we observe a fraction  $E_3 = E_4 \cdot k$  where  $k$  is the dimensional projection factor. In the dimensional time-dilation picture (§2.3), the 4D event that contains the black hole is a spatially extended process with a finite duration in 4D time. From our 3+1D frame, we see only a brief slice of the 4D duration. A fast process in 4D (one that completes in brief 4D time) projects to a complete cosmic history in 3+1D (because the 4D duration is long compared to the 3+1D slice), and a slow process in 4D (one that takes much of the 4D duration) appears as a very slow 3+1D process (because the 3+1D slice is brief compared to the 4D duration). For Hawking radiation, the underlying 4D process is fast (relative to the 4D events full duration) but appears in 3+1D as a very slow process (the famous  $10^{67}$  years for solar-mass black hole evaporation). This is not because 4D time is faster than 3+1D time (a confusing earlier phrasing in this section) – it is because the rate of the underlying 4D process, as seen from our brief 3+1D slice, is vastly slower than the rate the 4D process would have if observed from a longer 3+1D slice. The information paradox is resolved in this view: the information is in 4D, and Hawking radiation is the slow leak of that information back to 3+1D, not a thermal emission that destroys information. The temperature of Hawking radiation is set by the dimensional time-dilation factor, not by the surface gravity in 3+1D alone. (We acknowledge that this is a speculative extension; the standard semiclassical derivation of Hawking radiation is well-established, and our 4D-energy-leakage interpretation is offered as a possible alternative rather than a replacement.)

Black holes as dominant dark matter contributors. In the dimensional-cascade framework, every energetic event creates a 2D universe. Black holes are the most energetic events in our universe. By the dimensional time-dilation rule (§2.3), a black hole with  $\ell_{event} \sim 3 \times 10^3$  m (stellar mass, Schwarzschild radius  $\sim 3$  km, or  $\sim 2.95$  km for a solar-mass BH) creates a 2D universe that lasts  $\sim 10^{-5}$  seconds in our frame, while a supermassive black hole with  $\ell_{event} \sim 1.2 \times 10^{10}$  m (Sagittarius A\* mass,  $\sim 4.3 \times 10^6 M_\odot$ , Schwarzschild radius  $\sim 1.18 \times 10^{10}$  m) creates a 2D universe that lasts  $\sim 40$  seconds in our frame. Numerical correction: an earlier version of this paragraph used  $\ell_{event} \sim 10^{10}$  m and a 2D lifetime of  $\sim 30$  s for Sgr A; these have been corrected to  $\sim 1.2 \times 10^{10}$  m and  $\sim 40$  s, based on the Sgr A mass and the dimensional time-dilation rule  $\ell/c$ . The 2D universes created by black holes are more energetic, longer-lived (in our frame), and more gravitationally significant than

those created by photon emissions or atomic transitions. Therefore, if black holes are still actively creating 2D universes in a galaxy (e.g., during AGN outbursts or stellar black hole formation events), those 2D universes would contribute disproportionately to the current dark matter in that galaxy. This is a current effect (per §4.2), not a historical one: only the 2D universes being created now contribute to the current dark matter density, weighted by their individual energies. Historical black hole activity contributes only via the current event rate (which depends on the current AGN activity, the current rate of stellar black hole formation, etc.). The model predicts that galaxies with active black holes should have somewhat higher dark matter content (per unit stellar mass) than galaxies with quiescent black holes, holding all other factors fixed.

A note on the event horizon vs. the black hole itself. Throughout this subsection, when we say black holes create 2D universes, we mean the event horizon creates 2D universes, not the black hole interior. The black hole interior (the singularity, the content that has fallen in) is in 4D, per the framing of §4.10. The event horizon is the 3+1D boundary a real 3+1D structure that exists in our universe. The 2D universe creation is a boundary effect at the event horizon, not an interior effect at the singularity. This is consistent with the §2.3 principle that every energetic event in our 3+1 dimensional universe creates a 2D universe: the event horizon is a 3+1D structure, and its energetic processes (the extreme curvature and quantum effects at the horizon) create 2D universes. The black hole interior (4D) does not directly create 2D universes; the black hole event horizon (3+1D) does.

Testable prediction: dark matter correlates with black hole activity, not just stellar mass. This is a sharper version of the §4.7 prediction. Two galaxies of the same total stellar mass but different black hole activity (e.g., one with an active galactic nucleus, one without) should have different dark matter content, even at fixed stellar density. The galaxy with more recent black hole activity should have more dark matter. This is testable with existing galaxy surveys: select pairs of mass-matched galaxies with different AGN activity, and compare their dark matter content inferred from rotation curves, velocity dispersions, or gravitational lensing. Standard CDM predicts similar dark matter content for mass-matched galaxies; this model predicts more dark matter in the more AGN-active galaxy.

Speculation: primordial black holes and dark matter. If primordial black holes (formed in the early universe) existed, they would have produced 2D universes that contributed to dark matter. In this model, primordial black holes could be the seed of dark matter structure. This is speculative but could be tested: if primordial black holes have a specific mass distribution, the model would predict a specific initial dark matter distribution that could be compared to cosmological observations.

The 4D event as the energy reservoir of the universe. In the dimensional-cascade framework, the 4D event is the parent of our 3+1D universe. The 4D events total energy is our universes total mass-energy (per the energy conservation of §2.2). The 4D events true energy is the integrated energy over its full 4D duration, which is vastly larger than the energy in our 3+1D frame (because we only see a brief slice of the 4D event). The concentration of this 4D energy at a black hole (a tear to 4D) could explain why black hole time dilation is so extreme: a black hole is connected to the entire 4D energy reservoir of the universe via the dimensional transition, which makes the local gravitational effect much stronger than the local 3+1D mass concentration alone would suggest.

Speculation: the speed of light as a dimensional projection. In standard brane-world physics, the fundamental speed is the higher-D speed, and the 3+1D speed of light  $c$  is the effective speed on the brane a projection of the higher-D causality. In this model, our 3+1D speed of light  $c$  would be the projection of the 4D events causal structure into 3+1D. Specifically,  $c$  in 3+1D might be  $c \approx c_4 \cdot k$  for some dimensionless projection factor  $k$  (where  $c_4$  is the natural 4D speed). The value of  $c$  in our universe is then not a fundamental constant but a consequence of the dimensional projection. The model does not currently derive the value of  $k$  from the geometry, but the framing

is consistent with brane-world physics. If the dimensional cascade continues (4D  $\rightarrow$  3+1D  $\rightarrow$  2D), the effective speed of light might differ at each level of the cascade. This is speculative but testable in principle: in a 2D universe created by an energetic event, the speed of light might differ from our 3+1D  $c$  by a factor related to the dimensional projection. Of course, we cannot directly observe 2D universes, so this prediction is not directly testable.

The 4D events causal structure and the speed of light. The 4D event is not a moving object – it is a spatially extended event with a finite duration in 4D time. The 4D speed  $c_4$  in the previous paragraph is the conversion factor between 4D spatial extent and 4D temporal duration: a 4D event with spatial extent  $\ell_{4D}$  has a full duration  $\Delta t_{4D} = \ell_{4D}/c_4$  in 4D time (per §2.2). The 3+1D speed of light  $c$  is not a projection of the 4D events velocity – the 4D event doesn't have a velocity. Instead,  $c$  is a property of the dimensional projection mechanism itself: the projection from 4D to 3+1D maps 4D causal structure to 3+1D causal structure, and the ratio of the projected causal speed to the native 3+1D speed is set by the projection factor  $k$  (with  $c = c_4 \cdot k$ ). The 3+1D sees a maximum causal speed  $c$  in its frame, set by the projection. The 4D event itself is not moving at any speed in 4D – it is a localized energetic process in 4D, with a finite spatial extent and finite duration, that projects into 3+1D as a spatially extended universe with a finite lifetime. The speed of light  $c$  in 3+1D is a property of the projection, not a property of the 4D events motion. We acknowledge that the §4.10 phrasing the 4D event can move at  $c_4 > c$  is misleading: the 4D event is not a moving object, and the relationship between  $c_4$  and  $c$  is a projection factor, not a relative velocity. (This is a correction to a confusing earlier phrasing; the substance of the speculation – that  $c$  is a consequence of the dimensional projection rather than a fundamental constant – is preserved.)

Honest acknowledgment. This subsection is highly speculative. The void in 3+1D interpretation of black holes is not derived from the model, and the connection between black hole activity and dark matter is a prediction that has not been tested. The mainstream view of black holes (as regions of extreme 3+1D spacetime curvature) is the default interpretation. We offer this subsection as a possible extension of the model, with appropriate caveats.

#### 1.5.11 4.11 Speculative extension: all fundamental constants as projections of the 4D event

This subsection is the most speculative part of the paper. It is offered as a philosophical/interpretive extension, not a derived claim. We include it because it follows naturally from the dimensional-cascade framework, but it should be read with appropriate skepticism.

The puzzle of the constants. Standard physics leaves many constants unexplained. The electron has a specific mass ( $\sim 511 \text{ keV}/c^2$ ). The speed of light has a specific value ( $c \approx 3 \times 10^8 \text{ m/s}$ ). Planck's constant has a specific value. The fine structure constant is  $\sim 1/137$ . The proton-to-electron mass ratio is  $\sim 1836$ . The cosmological constant has a specific (small) value. Absolute zero is exactly 0 K. The list goes on. These constants are measured, not derived. We use them in our equations, but we don't have a theory of why they have the values they do.

The dimensional-cascade interpretation. In the dimensional-cascade framework, all of these constants would be consequences of the specific 4D event that created our 3+1D universe. The 4D event has specific properties: a specific energy, a specific spatial structure, a specific duration, a specific set of internal dynamics. The dimensional projection of that specific event into 3+1D gives a specific set of constants. Different 4D events would give different 3+1D universes with different constants.

In this view: - The electron mass is a consequence of the 4D events specific energy spectrum, projected into 3+1D - The speed of light  $c$  is a consequence of the 4D events causal structure, projected into 3+1D - The Planck constant  $\hbar$  is a consequence of the 4D events action scale, projected into 3+1D - The fine structure constant  $\alpha \approx 1/137$  is a consequence of the dimensional

projection factor for the electromagnetic coupling - The gravitational constant  $G$  is a consequence of the bulk-brane cancellation factor  $\epsilon$  (§2.4, §2.6) - The cosmological constant (dark energy density) is the un-cancelled fraction of the inverted 4D gravity (§2.4)

Two mechanisms for constants are determined by the 4D event. Note that the phrase constants are determined by the 4D event can mean different things for different classes of quantities: - For 3+1D particles created during the Big Bang (electron, proton, photon, neutrino, etc.): the 4D event projects a Big Bang into our 3+1D brane, and during the Big Bang, these particles are created with specific masses, charges, and couplings (per Standard Model particle physics). The constants of the particles are set by the Standard Model (with the Standard Models free parameters ultimately being consequences of the 4D event). The neutrinos small mass, the electrons larger mass, the photons zero mass all are set by the 4D events specific energy spectrum, projected into 3+1D via the Big Bang. - For universal constants (speed of light, Planck constant, fine structure constant, gravitational constant, cosmological constant): the constants are set by the dimensional projection mechanism itself, not by specific particles. The speed of light, for example, is the projection of the 4D events causal structure into 3+1D; the fine structure constant is the projection factor for the electromagnetic coupling.

All these mechanisms lead to the same conclusion: the 4D event determines the constants of 3+1D physics. The specific mechanism differs (creation for particles, projection-mechanism-property for universal constants), but the result is the same: constants are not free parameters, they are consequences of the 4D event.

The constants are not fundamental. In this view, the fundamental constants are not free parameters of nature they are determined by the specific 4D event that created our universe. The fine-tuning problem (why do the constants have values that allow stars, planets, life?) is reframed: the constants arent tuned for us; we exist because our parent 4D event had these specific properties. Other 3+1D universes (from other 4D events) have different constants, and those universes might have their own fine-tuning for their specific physics.

Testable consequence: constants should be related. If all constants come from the same 4D event, they should be related to each other through the dimensional projection. The dimensionless constants (fine structure constant, electron-to-proton mass ratio, etc.) might be predictable from the geometry of the dimensional projection, not independent. The model does not currently derive these relations, but a specific implementation might be able to.

The multiverse by construction. The dimensional-cascade framework mechanistically generates a multiverse: each 4D event is a different parent, and each parent creates a 3+1D child universe with different constants. This is stronger than the standard string theory landscape (which is a theoretical construct): the dimensional cascade generates the multiverse through the dimensional projection mechanism.

Honest acknowledgment. This is the most speculative part of the paper. The claim that all fundamental constants are consequences of the dimensional projection is not derived from the model. The model provides a framing in which this is plausible, but the actual derivation of specific constant values from the geometry is left to future work. The mainstream view treats the constants as free parameters to be measured; this subsection offers an alternative framing in which the constants are determined by the dimensional projection. We offer this as a philosophical/interpretive extension, with appropriate skepticism.

1.5.12 4.12 (Section removed: the neutrino interpretation was speculative and introduced internal inconsistencies. Neutrino properties are taken as given in the Standard Model; the dimensional-cascade framework does not currently address them.)

1.5.13 4.13 Speculative extension: the weak force as a dimensional-projection effect

This subsection extends the dimensional-cascade framework to the weak nuclear force. It is offered as a conceptual extension that connects the model to the Standard Models parity-violating, flavor-changing, short-range force. As with the other speculative extensions, it should be read with appropriate skepticism.

The weak force in the Standard Model. The weak force is one of the four fundamental forces. It is mediated by the  $W$ 's and  $Z$  bosons (massive,  $\sim 8090$  GeV), it acts only on left-handed particles and right-handed antiparticles (parity violation), it can change particle flavor (e.g., neutron  $\rightarrow$  proton + electron + antineutrino in beta decay), and it has a very short range ( $\sim 10^{-16}$  m) due to the  $W/Z$  mass. The weak force is weak at long range because the massive mediators decay quickly into the vacuum, but at very short range it is comparable in strength to the electromagnetic force.

The dimensional-cascade interpretation. In the framework of §4.11 (all fundamental constants as projections of the 4D event), the weak forces constants would all be consequences of the specific 4D event that created our universe:

- $W/Z$  boson mass: a consequence of the dimensional projection factor for the weak-force mediator
- Higgs VEV: a consequence of the 4D events specific symmetry-breaking structure
- CKM and PMNS mixing angles: consequences of 4D mixing structures projected into 3+1D
- Weak coupling constant  $g_W$ : a consequence of the dimensional projection factor for the weak force
- Range of the weak force ( $r \sim \hbar/(m_W c)$ ): a consequence of the  $W/Z$  mass
- Strength at short range: a consequence of the coupling constant

In this view, the weak force is not unified with the dimensional cascade in a new way – it is described by the dimensional cascade in the same way as the other forces. The dimensional cascade doesn't add new forces; it gives a deeper origin for the existing ones.

Parity violation as a dimensional effect. The most interesting connection is parity violation – the weak forces left-handed-only coupling. In the Standard Model, this is a fundamental property of the weak interaction, but it is not derived from deeper principles. In the dimensional-cascade framework, a natural interpretation is that the 4D event has a specific chirality (handedness), and 3+1D particles that are left-handed in 4D project as left-handed in 3+1D. Right-handed 4D structures project as antiparticles in 3+1D (this is consistent with the Standard Model, where right-handed antiparticles exist). The 4D events chirality biases the creation of 3+1D particles toward left-handed particles, which is why the weak force couples only to left-handed particles. This would explain why the weak force is the only force that violates parity: it is the only force that is sensitive to the chirality of the 4D event. Photons and gluons are achiral in 4D (they don't have a handedness), so they couple equally to left- and right-handed 3+1D particles.  $W/Z$  bosons are chiral in 4D, so they couple only to one handedness in 3+1D. The graviton is a separate case: it is not achiral in 4D – it is inverted at the dimensional boundary (§2.4). The graviton couples to all particles (mass-energy), but its coupling is suppressed by the bulk-brane cancellation. So the graviton doesn't have a chirality in the simple sense; it has an inversion in the dimensional projection. This is a real idea in some string/brane theories: chirality in 4D is related to the orientation of strings/branes, and 3+1D parity violation is a consequence of how 4D structures project.

Flavor changing as a dimensional effect. The weak force is the only force that can change particle flavor. In the Standard Model, this is described by the CKM matrix (for quarks) and the PMNS matrix (for neutrinos), which encode the mixing between flavor and mass eigenstates. In the dimensional-cascade framework, these mixing matrices are projections of 4D mixing structures. The mixing angles are determined by the 4D event. Different 4D events would give different mixing angles. The CKM and PMNS matrices are not fundamental constants—they are consequences of the dimensional projection. A specific implementation of the model might derive the CKM/PMNS mixing angles from the geometry of the dimensional projection, but this is left to future work.

Short range as a consequence of the projection. The short range of the weak force is not a new effect in this model—it is a consequence of the W/Z mass, which is itself a consequence of the dimensional projection. In this view, the weak force is weak at long range because the W/Z are massive, and the W/Z are massive because the dimensional projection sets their mass. There is no new mechanism—just a deeper origin for the existing one.

The electroweak unification. In the Standard Model, the weak force is unified with electromagnetism as the electroweak force at high energies ( $\sim 100$  GeV). The Higgs mechanism breaks this symmetry at low energies, giving the W/Z their mass while leaving the photon massless. In the dimensional-cascade framework, the electroweak unification is a 3+1D phenomenon, and the Higgs mechanism is a 3+1D description of a deeper dimensional-projection process. The model does not replace the Higgs mechanism—it provides a deeper origin for why the Higgs mechanism works.

Unification with the other forces? A natural extension: in some grand-unified theories (GUTs), the strong, weak, and electromagnetic forces are unified at very high energies ( $\sim 10^{16}$  GeV). The dimensional-cascade framework could potentially provide a deeper origin for GUT-scale unification, but this is not part of the current model. The model is consistent with GUTs, but does not predict them. We leave this as an open question for future work.

Honest acknowledgment. This subsection is highly speculative. The claim that the weak forces constants are consequences of the dimensional projection is not derived from the model. The claim that parity violation is a dimensional effect is not derived from the model. The claim that flavor changing is a dimensional effect is not derived from the model. All three are interpretive extensions that connect the dimensional-cascade framework to known phenomena. The mainstream view treats the weak force as described by the Standard Model with the Higgs mechanism. We offer this subsection as a conceptual extension, with appropriate skepticism.

#### 1.5.14 4.14 Speculative extension: the strong force as a dimensional-projection effect

This subsection extends the dimensional-cascade framework to the strong nuclear force. It is the fourth and final\* force to be addressed. The strong force is the hardest to unify with the dimensional cascade, because it does not have a unique feature that maps directly to 4D physics the way gravity (weakness), electromagnetism (the speed of light), and the weak force (parity violation) do. We include it for completeness, with the explicit understanding that the connections are less direct than for the other forces.\*

The strong force in the Standard Model. The strong force is mediated by gluons (8 of them, all massless). It couples to color charge (three types: red, green, blue, with corresponding anti-colors). It only acts on quarks and gluons (not on leptons). The strong force has asymptotic freedom (the coupling gets weaker at short distances) and confinement (quarks cannot be isolated; they are always bound into hadrons). At low energies, the strong force has the largest coupling constant of the four forces ( $\sim 1$ , compared to EMs  $1/137$ ). The strong force holds quarks together inside protons and neutrons, and holds protons and neutrons together inside atomic nuclei.

The dimensional-cascade interpretation. In the framework of §4.11 (all fundamental constants

as projections of the 4D event), the strong forces constants would all be consequences of the specific 4D event that created our universe:

- Gluon mass (0): a consequence of the dimensional projection for massless mediators
- Strong coupling constant  $\alpha_s$ : a consequence of the dimensional projection factor for the strong force
- Color charge (3 types): the number 3 might be related to the 3 spatial dimensions of 3+1D, but this is not derived from the dimensional cascade
- Confinement scale  $\Lambda_{QCD} \sim 200$  MeV: a consequence of the dimensional projection factor
- Asymptotic freedom: a consequence of gluon-loop anti-screening in 3+1D, which is not directly addressed by the dimensional cascade

In this view, the strong force is not unified with the dimensional cascade in a new way it is described by the dimensional cascade in the same way as the other forces.

The hierarchy of force strengths. The relative strengths of the four forces at low energies are: strong ( $\sim 1$ ), EM ( $\sim 1/137$ ), weak ( $\sim 10^{-5}$ ), gravity ( $\sim 10^{-38}$ ). The huge range (38 orders of magnitude) is one of the deepest puzzles in physics. In the Standard Model, the hierarchy is unexplained we measure the couplings and accept the values. In the dimensional-cascade framework, the hierarchy is a consequence of the dimensional projection: each forces coupling is set by a different projection factor, and the specific 4D event determines the relative magnitudes. Gravity is the weakest because of the bulk-brane cancellation (g2.4). The strong force is the strongest at low energies because the dimensional projection factor for the strong force is largest. The EM and weak forces are intermediate. This is not a derivation of the hierarchy it is a reframing of the hierarchy as a consequence of the dimensional projection.

Asymptotic freedom and confinement. Asymptotic freedom (the strong force gets weaker at short distances) is due to gluon-loop anti-screening in 3+1D. Confinement (quarks cannot be isolated) is a consequence of the strong forces running coupling the force gets stronger at long distances, so quarks cannot be pulled apart without creating new quark-antiquark pairs. In the dimensional-cascade framework, asymptotic freedom and confinement are 3+1D phenomena, not directly addressed by the dimensional projection. The model is consistent with asymptotic freedom and confinement, but does not derive them.

Color charge (3 types) and 3 spatial dimensions. The strong force has three color charges (red, green, blue). Our universe has three spatial dimensions. This numerical coincidence is suggestive in some string theories, the number of colors is related to the number of compactified dimensions. In the dimensional-cascade framework, the three colors might be a consequence of the 3+1 dimensional structure, but this is not derived. We note the coincidence but do not claim it as a prediction of the model.

The unification of all four forces. The dimensional-cascade framework does not unify the four forces in the sense of grand-unified theories (GUTs). It is consistent with GUTs (the couplings would unify at some high energy, set by the 4D event), but it does not predict the specific unification scale or the specific GUT group. The model is a framework for thinking about the origins of the forces properties, not a theory that derives them. The unification offered by the dimensional cascade is a conceptual unification (all four forces are consequences of the same 4D event), not a quantitative unification (the couplings don't necessarily merge at a specific energy in this model).

Honest acknowledgment. This subsection is highly speculative. The claim that the strong forces constants are consequences of the dimensional projection is not derived from the model. The hierarchy of force strengths is reframed but not derived. Asymptotic freedom and confinement are 3+1D phenomena, not directly addressed by the dimensional cascade. The number 3 for color charge



is suggestive but not derived. The strong force is the hardest of the four forces to unify with the dimensional cascade, because it lacks a unique feature (like parity violation for the weak force) that maps directly to 4D physics. The mainstream view treats the strong force as described by quantum chromodynamics (QCD) with the specific structure of  $SU(3)$  color. We offer this subsection as a conceptual extension, with appropriate skepticism.

#### 1.5.15 4.15 Speculative extension: what Einstein was missing unification in 4D, not 3+1D

This subsection is a historical and philosophical note that places the dimensional-cascade framework in the context of Einsteins lifelong quest for a unified field theory. We include it because the dimensional-cascade model offers a specific diagnosis\* of why Einsteins program failed, and a specific alternative unification in 4D rather than 3+1D. As with the other speculative extensions, this is interpretive rather than derived.\*

Einsteins unified field theory program. From the 1920s until his death in 1955, Einstein worked on a unified field theory a program to merge gravity and electromagnetism into a single geometric framework. He sought to derive the electromagnetic field from the geometry of spacetime, in the same way that general relativity derives gravity from spacetime curvature. Einsteins program failed. The unified field theory he sought was never found.

Why Einsteins program failed. In retrospect, Einsteins program failed for several reasons:

1. He didnt know about the weak and strong forces. These were discovered later (1930s Fermi for weak, 1970s QCD for strong). His goal of unifying just gravity and EM was too limited.
2. He rejected quantum mechanics. Einstein famously declared God does not play dice. This was a problem because electromagnetism is fundamentally quantum (QED). A purely geometric unification of gravity and EM cannot work, because EM is not purely geometric it has quantum features (photons, vacuum fluctuations, etc.).
3. He didnt know about the dark sector. Dark matter and dark energy were not on the radar in Einsteins time. A complete unification would need to account for them.
4. He worked in 3+1D. Einsteins framework was general relativity in 3+1D. He did not consider that the unification might be in a higher dimensional structure, with the four forces being different projections of that higher-D structure.

The dimensional-cascade diagnosis. In the dimensional-cascade framework, gravity and EM are unified in 4D, not in 3+1D. In 3+1D, gravity and EM look like different forces with different properties: gravity is geometric (curvature of spacetime), EM is vector (the electromagnetic potential), gravity couples to mass-energy, EM couples to electric charge, gravity is purely attractive at long range, EM has both attraction and repulsion. These differences are real in 3+1D but in 4D, they are projections of the same underlying structure. The 4D structure is one; the 3+1D projections are different. This is why Einstein could not unify them in 3+1D: the unification is not in 3+1D.

Why gravity and EM look so different in 3+1D. The differences between gravity and EM in 3+1D geometric vs. vector, tensor vs. vector field, attractive vs. attractive/repulsive, classical vs. quantum are consequences of the dimensional projection. The 4D structure is projected into 3+1D in different ways for the two forces, giving different mathematical structures, different symmetries, and different quantum vs. classical behavior. Einstein sought to derive these differences from a single 3+1D geometry, but the differences come from the projection, not from the underlying 3+1D structure.

Implication for the weak and strong forces. The same diagnosis applies to the weak and strong forces: they are unified with gravity and EM in 4D, not in 3+1D. In 3+1D, the four forces look

like four different forces with different properties (mediators, ranges, couplings, parity behaviors). In 4D, they are all projections of the same 4D structure. The differences in 3+1D are consequences of the projection. Einsteins program of unifying the forces in 3+1D was therefore destined to fail: the unification is not in 3+1D.

Connection to modern unification attempts. Modern unification attempts (GUTs, string theory, loop quantum gravity) take different approaches:

- GUTs unify the strong, weak, and EM forces in 3+1D at very high energies ( $\sim 10^{16}$  GeV). They do not include gravity.
- String theory unifies all four forces in 10 or 11 dimensions, with the extra dimensions compactified. The 3+1D forces are projections of the higher-D strings.
- Loop quantum gravity quantizes 3+1D gravity directly, without unifying the other forces.

The dimensional-cascade framework is closest to string theory in spirit: both rely on higher-D structures projecting into 3+1D. The key difference is that the dimensional-cascade framework does not require compactification of extra dimensions the 4D event is a brief slice of a higher-D process, and the 3+1D universe is a projection of that slice. This is a different interpretation of how the higher-D structure relates to 3+1D.

Why string theory needs 10 dimensions. A natural question: why does string theory require 10 dimensions (or 11 in M-theory) when the dimensional-cascade framework requires only 4? The answer lies in the ambition of the two frameworks. String theory attempts to derive all of physics from a single mathematical object (vibrating strings). For the quantum theory to be mathematically consistent (anomaly-free), it requires a specific number of dimensions. Bosonic string theory requires 26 dimensions; superstring theory requires 10 (with supersymmetry); M-theory requires 11. The 10 dimensions are compactified to 3+1D at the Planck scale ( $\sim 10^{-35}$  m), with the extra 6 dimensions curled up in Calabi-Yau manifolds or similar structures. The complexity of string theory comes from this requirement: the extra 6 dimensions must be compactified in specific ways, and there are vastly many possible compactifications ( $\sim 10^{500}$  to  $10^{1000}$ , the landscape). Choosing the right compactification is the landscape problem, and string theory has not solved it.

The dimensional-cascade framework is simpler by design. The dimensional-cascade framework is less ambitious than string theory. It does not attempt to derive the Standard Model from first principles. It is a thought experiment that reinterprets existing physics (the dark sector, the four forces) through a dimensional-cascade lens. It does not require 10 dimensions, does not require compactification, does not require supersymmetry, and does not have a landscape problem. The model is conceptually simpler: a 4D event projects into 3+1D, the cascade is scale-invariant, gravity inverts at the dimensional boundary, and the dark sector is the cumulative effect. The price of this simplicity is that the model is less quantitative than string theory: it does not derive the specific values of the Standard Model parameters. The model is a framework for thinking about the dark sector and the four forces, not a theory that derives them from first principles.

A philosophical note. The complexity of string theory reflects the ambition of the program: deriving all of physics from a single mathematical object. The simplicity of the dimensional-cascade framework reflects its modesty: it is a thought experiment that reinterprets the dark sector, not a theory of everything. Both approaches have value. String theory is a mathematical framework that may eventually yield testable predictions (or may not). The dimensional-cascade framework is a conceptual framework that yields testable predictions now (RAR, DF2/DF4, no direct detection) but is less mathematically rigorous. We do not claim that one is better than the other; we offer the dimensional-cascade framework as a complementary approach, useful for thinking about the dark sector even if it does not replace more fundamental theories.

The broader landscape of unification attempts. String theory is the most famous unification attempt, but it is not the only one. Other major programs include: (1) Loop Quantum Gravity (LQG), which quantizes 3+1D spacetime using loops and spin networks, but does not include the Standard Model forces; (2) Causal Set Theory, which treats spacetime as a discrete set of causally-related events, addressing the quantum gravity problem but with limited contact to particle physics; (3) Causal Dynamical Triangulations (CDT), a numerical approach that builds spacetime from simplices and has shown 4D spacetime emerges from the construction; (4) Asymptotic Safety, which proposes that gravity has a fixed point at high energies, making it renormalizable without new structures; (5) Twistor Theory (Penrose), a mathematical reformulation of spacetime that has yielded real results in scattering amplitudes; (6) Noncommutative Geometry (Connes), which replaces continuous spacetime with noncommutative algebra and has reproduced the Standard Model + gravity in some versions; (7) Kaluza-Klein Theory, the original 5D unification of gravity and EM, whose principles live on in string theory; and (8) Brane-World Scenarios (Randall-Sundrum, ADD), which place our universe on a 3+1D brane in a higher-D bulk, and are conceptually closest to the dimensional-cascade framework.

Positioning within this landscape. The dimensional-cascade framework is closest to brane-world scenarios (Randall-Sundrum, ADD), which are referenced in §2.1 and §2.4. Both rely on a higher-D bulk and a 3+1D brane projection. The key difference is the downward perceptual inversion principle (per §2.4): the dimensional-cascade framework postulates that the bulks gravity is perceived as inverted by the child universes brane (the projected contribution is repulsive from the branes perspective), while the underlying gravity in the bulk remains attractive (standard GR). The physical mechanism for this perceptual inversion is grounded in the standard GR  $\rho + 3P < 0$  mechanism for negative effective gravitating density (per §2.4): the bulk-brane coupling translates the bulks ordinary attractive matter into a brane-perceived effective gravitating density with the opposite sign, in the same way that an inflaton field or the cosmological constant has negative effective gravitating density in our universe. The upward back-projection (child → parent) does not invert the perception; the parent perceives the childs net attractive gravity as attractive = dark matter. Standard brane-world models do not make this perceptual-inversion claim; they describe the branes effective gravity as suppressed by geometric dilution (ADD) or warping (RS), but with the same sign as the bulk. The cascades claim is that the specific bulk-brane coupling produces a negative effective gravitating density on the brane (via the standard  $\rho + 3P < 0$  mechanism), which is a stronger (more specific) version of standard brane-world models. The dimensional-cascade framework is also distinct from LQG (which works in 3+1D, not 4D projection), causal set theory (discrete vs. continuous), and the various algebraic reformulations (noncommutative geometry, twistor theory). The model is not a replacement for any of these programs; it is a thought experiment that offers a different interpretation of the dark sector and the four forces, with testable predictions that other programs do not currently make.

What is unique about the dimensional-cascade framework. Among all these unification attempts, the dimensional-cascade framework has several unique features: (1) scale-invariant cascade every energetic event creates lower-D universes, not just the original Big Bang; (2) downward perceptual inversion principle downward dimensional projection is perceived by the child as inverted (the underlying gravity in the bulk remains attractive; the inversion is a feature of the projection mechanism, not a violation of GR), upward back-projection is not perceived as inverted; (3) dark sector as direct consequence dark matter and dark energy are direct consequences of the cascade, not added assumptions; (4) testable predictions now the model makes specific testable predictions (RAR, DF2/DF4, no direct detection, activity-dependence) without requiring new physics; (5) conceptual simplicity the framework is intuitive (dimensional cascade, energy conservation, scale-invariance, directional perceptual inversion), not mathematically heavy. These features distinguish the model

from string theory (which is mathematically heavy but not testable now), LQG (which is mathematically rigorous but limited to gravity), and the other unification programs. We do not claim the dimensional-cascade framework is better than these programs; we claim it is different, with a focus on the dark sector and testable predictions rather than on mathematical rigor or unification of all forces.

Einstein's intuition was correct, but he was looking in the wrong place. Einstein's intuition that the four forces should be unified is shared by the dimensional-cascade framework. The difference is where the unification is sought. Einstein sought it in 3+1D geometry; the dimensional-cascade framework seeks it in 4D structure, with 3+1D as a projection. The unified field theory Einstein wanted is not in 3+1D; it is in the 4D event that projects into 3+1D.

Honest acknowledgment. This is a historical and philosophical note, not a physical claim. We do not claim to have derived Einstein's unified field theory; we offer a diagnosis of why his program failed, in the language of the dimensional-cascade framework. The mainstream view treats Einstein's program as a historical dead end, replaced by the Standard Model + general relativity + quantum field theory. The dimensional-cascade framework is a speculative alternative that places the unification in 4D. We offer this as a philosophical extension, with appropriate caveats.

#### 1.5.16 4.16 Positioning within the unified-dark-sector landscape

This subsection positions the dimensional-cascade framework within the specific\* niche of attempts to unify the dark sector (dark matter + dark energy + the hierarchy problem) with the four fundamental forces. We include it because there is a growing literature on this topic, and the dimensional-cascade framework has both overlaps with and distinctions from existing programs.\*

Major unified-dark-sector attempts. A wide variety of programs attempt to unify the dark sector with the four forces or to replace the dark sector with modifications of known physics. Major examples include: (1) MOND [Milgrom83] and its relativistic extensions (TeVeS, BIMOND, etc.) modify Newton's second law at low accelerations to replace dark matter, but have difficulties with CMB and gravitational lensing; (2) Verlinde's Emergent Gravity [Verlinde16] derive MOND-like phenomenology from entropic gravity, but limited to static situations; (3) Superfluid Dark Matter [Berezhiani15] dark matter is a real particle that forms a superfluid at galactic scales, combining CDM and MOND successes; (4) Unified Dark Matter / Chaplygin Gas [Kamenshchik01, Bento02] a single fluid acts as both dark matter and dark energy, but is strongly constrained by data; (5) CDM + Baryonic Feedback [Kravtsov24] the mainstream alternative, using better simulations of the standard model; (6) Scalar-Tensor /  $f(R)$  Gravity geometric modifications of gravity that can mimic dark sector effects; (7) Massive Gravity / Bi-Gravity [deRham11, Hassan12] the graviton has a mass, modifying gravity at large scales; (8) Mirror Matter [Foot95] a parallel Standard Model sector provides dark matter candidates; (9) Dark Fluid / Negative Mass [Farnes18] a fluid with negative mass that creates both dark matter and dark energy effects.

What is unique about the dimensional-cascade framework. Among all these programs, the dimensional-cascade framework has several unique features for unifying the dark sector with the four forces:

1. Both dark sector products as direct consequences of the cascade. Dark matter is the cumulative gravity of 2D universes (§2.5); dark energy is the un-cancelled fraction of inverted 4D gravity (§2.4). Both are automatic consequences of the dimensional projection, not added assumptions.
2. Same mechanism for hierarchy + dark matter. The hierarchy problem (gravity is weak) and the dark matter problem (dark matter is weak) are both consequences of bulk-brane cancellation

at different cascade levels (§2.6). This unifies two otherwise-distinct problems.

3. Testable predictions that distinguish from other programs. The model predicts that RAR scatter should correlate with galaxy activity (MOND predicts no such correlation, since MOND has no activity-dependence); DF2/DF4 type tests where dark matter depends on stellar activity, not just stellar mass; and no direct dark matter detection (since dark matter is 2D universes, not particles). These are distinguishing predictions, not shared with other unified-dark-sector programs.
4. Conceptual origin from a single 4D event. All four forces + the dark sector come from the same 4D event. Different forces and dark-sector products are different projections of the same underlying structure. This is more economical than most other approaches, which typically treat the dark sector as a separate phenomenon.
5. Consistent with prior work, but not a replacement. The model is compatible with brane-world scenarios (RS, ADD), MOND-like phenomenology, and Verlinde-like emergent gravity. It does not replace these programs – it provides a deeper origin for the same phenomenology, with a dimensional-cascade mechanism rather than a modification of gravity or an entropic force.

Connections to specific programs. The model has closest connections to: (1) Brane-world scenarios [ADD98, RS99] – both use higher-D bulk + 3+1D brane; the dimensional cascade adds the downward inversion principle (downward projection inverts, upward back-projection does not) and the scale-invariant cascade; (2) Verlinde’s Emergent Gravity [Verlinde16] – both derive MOND-like phenomenology without dark matter particles; the dimensional cascade uses cumulative 2D universe gravity, not entropic gravity; (3) Superfluid Dark Matter [Berezhiani15] – both are hybrid approaches (real matter, MOND-like effects); the dimensional cascade uses 2D universes, not a superfluid; (4) MOND [Milgrom83] – both reproduce the Radial Acceleration Relation (§4.1); the dimensional cascade adds the activity-dependence prediction that MOND lacks.

What the model is not. The dimensional-cascade framework is not: (1) a particle dark matter model (it has no WIMPs, axions, or other particles); (2) a modified gravity model (it doesn’t modify Einstein’s equations, just reinterprets the gravitational field as the projected 4D gravity); (3) a string theory or M-theory (it doesn’t use vibrating strings or 10/11 dimensions); (4) a loop quantum gravity (it doesn’t quantize 3+1D spacetime); (5) a  $f(R)$  or scalar-tensor theory (it doesn’t modify the Einstein-Hilbert action). The model is a thought experiment that reinterprets existing physics through a dimensional-cascade lens, with testable predictions and a clear physical interpretation. It is less mathematically rigorous than string theory, LQG, or  $f(R)$  gravity, but more testable and more conceptually clear.

Honest acknowledgment. This is a positioning subsection, not a derivation. The model is not better than MOND, Verlinde, superfluid dark matter, or any other program. The model is different: it offers a dimensional-cascade origin for the dark sector, with testable predictions and a clear physical interpretation. We do not claim that the dimensional-cascade framework is the correct unification – we claim it is a useful thought experiment that may illuminate the dark sector, even if it is not the final theory. Other programs (MOND, Verlinde, superfluid dark matter, etc.) are legitimate alternatives, and the empirical data will ultimately decide between them.

## 1.6 5. Lower-dimensional universes and the dark matter connection

In the scale-invariant version of the model (§2.3), all energetic events in our 3+1 dimensional universe create 2-dimensional universes as their aftermath, in 2-dimensional subspaces of our 3D space. Each such event creates a tiny 2D universe with its own brief lifetime, set by the events energy. The 2D universe has its own physics, its own dark sector (in 2D), and its own lifetime (the brief aftermath of the originating 3+1 dimensional event, as experienced from inside the 2D universe).

This is not a separate speculation in the scale-invariant model – it is the same mechanism that creates our universe from the 4D event, applied in the other direction. The principle is symmetric: higher-dimensional events create lower-dimensional universes, and lower-dimensional universes are real, with their own physics and their own brief existences.

The 2D universes created by 3+1 dimensional events have lifetimes in our 3D frame that scale with the events spatial extent, via the dimensional time-dilation rule  $\Delta t_{2D} \sim \Delta x_{event} / c$  (per §2.3 line 105). A small event (LHC collision,  $\sim \text{TeV}$  scale,  $\Delta x_{event} \sim 10^{-16} \text{ m}$ ) creates a 2D universe that lasts  $\sim 3 \times 10^{-27}$  seconds in our frame; a large event (supernova,  $\Delta x_{event} \sim 10^6 \text{ m}$  for the expanding photosphere,  $\sim 10^{44} \text{ eV}$  of visible-light energy) creates a 2D universe that lasts  $\sim 30$  seconds in our frame. Numerical correction: an earlier version of this section said  $\Delta t_{2D} = 10^{-27} \text{ s}$  for a supernova and  $10^{-27} \text{ s}$  for the LHC; these have now been corrected to  $\sim 30 \text{ s}$  and  $\sim 3 \times 10^{-27} \text{ s}$ , respectively, based on the dimensional time-dilation rule. The 2D universes from small events are essentially instantaneous in our frame; the ones from large events are not. From the perspective of each tiny universe, that brief moment is the entirety of its cosmic history.

We emphasize: the 2D universes created by 3+1 dimensional events are real in the same sense that our 3+1 dimensional universe is real. They are not mathematical abstractions. They have their own spacetime, their own physics, and (in principle) their own observers. We simply cannot interact with them from our 3D vantage point, because the dimensional barrier prevents it.

The key implication, developed in §2.3, is that dark matter is the collective gravitational signature of all these 2D universes. Each tiny universe has its own gravity (a small replica of the dimensional-projection mechanism), and the cumulative gravitational effect of countless such universes is what we observe as dark matter.

---

## 1.7 6. Falsification

A thought experiment must be falsifiable to be useful. We identify the following observations that would refute the model:

1. Direct detection of dark matter as a normal particle. If dark matter is detected with standard particle interactions (electromagnetic, strong, or weak), it is a substance in 3+1 dimensional space and not the collective gravitational signature of 2D universes. The model would be refuted.
2. Dark matter self-interaction or coupling to visible matter is detected. The Bullet Cluster observation shows visible matter (gas) concentrating in the collision center while dark matter passes through, naturally explained if dark matter is a collisionless substance. In our model, the dark matter at the original location is the current collective gravity of 2D universes being created now in the cluster (i.e., by the clusters ongoing activity – residual gas, low-level accretion, etc.). The model predicts that the dark matter stays at the original location even as the visible matter (gas) interacts and slows, because the 2D universes are being created

where the clusters activity is, not where the visible matter goes. The Bullet Cluster result therefore supports the model. A future observation showing dark matter strongly interacting with itself or with visible matter (so that dark matter and gas stay together after cluster collisions) would be in tension with this model. The model as currently stated makes no specific prediction about dark matter self-interaction, beyond the general statement that dark matter is the cumulative gravity of 2D universes being created by the current activity.

3. Dark matter density does not track the radial acceleration relation. If dark matter is observed to be uncorrelated with visible matter in galaxies, the model is refuted.
4. Sub-millimeter gravity follows  $1/r\check{s}$  down to the Planck length. If gravity is measured at sub-millimeter scales with no deviation from Newtons law, any extra dimensions in the model must be smaller than experimental reach ( $\sim 10$  m). The dimensional-projection mechanism in this model does not by itself predict a specific size for the extra dimensions, so this constraint does not directly apply to the inversion part of the model but if the model includes ADD-style geometric suppression as part of its mechanism, the extra dimensions would need to be smaller than  $\sim 10$  m. A future implementation of the model with a specific geometry would need to be consistent with the sub-millimeter constraint, and the absence of any deviation down to the smallest measurable scale would be in tension with ADD-style interpretations.
5. The dimensional-projection mechanism is shown to be impossible. If a general argument (not based on this specific model, but on broader principles) shows that the kind of dimensional projection proposed here cannot exist (for example, if the inversion of gravitational sign under dimensional projection is mathematically forbidden in a consistent way), the model would be refuted. Note that this is different from finding a different explanation for the cosmological constant problem; the model could still be correct on its own terms even if other explanations exist.
6. Dark energy density is not approximately constant. The model predicts that dark energy density is approximately constant in our 3+1 dimensional frame, matching standard CDM behavior. If future surveys (Euclid, LSST, Roman Space Telescope, SKA) detect a changing dark energy with  $w$  significantly different from  $-1$ , or a measurable variation in the dark energy density, the model would be in tension. A very slow variation (corresponding to slow evolution of the 4D events antigravity output over its full duration) would be expected but probably undetectable.
7. CMB / primordial element constraints are violated by any 4D event implementation. The model requires the 4D event to be spatially extended and approximately homogeneous (see §4.5) to reproduce the near-scale-invariance of the observed CMB power spectrum. A specific 4D event implementation must satisfy this homogeneity constraint. If every physically reasonable 4D event geometry leads to a CMB power spectrum that is inconsistent with observations (e.g., not nearly scale-invariant, or with too much anisotropy on large scales), the model would be refuted. The model does not currently derive a specific CMB prediction, so this criterion can only be tested when a specific 4D event geometry is proposed.
8. Gravity varies significantly with cosmic time. If  $G$  is observed to vary at a level inconsistent with the models prediction ( $G$  is approximately constant, because the bulk-brane cancellation is approximately constant during our universes brief slice of the 4D events full duration), the model is refuted.

9. Dark matter density does not correlate with energetic event rates on galaxy scales. If precise measurements show that mass-matched galaxy pairs with different stellar densities do not have different dark matter content, the scale-invariant version of the model is refuted. (The basic version of the model, in which the 4D event is special, would be unaffected by this test.) The model also predicts that dark matter density should not be enhanced in the Suns interior, near nuclear reactors, near particle accelerators, or in the Earths core stellar-scale and sub-stellar-scale effects are too small to be detectable. Detectable enhancement at any of these locations would be in tension with the model.

We acknowledge that the model is currently difficult to falsify in a clean way. The dimensional structure is undetermined, the bulk field content is unspecified, and the coupling constants are not derived. The model is best understood as a geometric hypothesis in need of theoretical development before it can be precisely tested.

---

## 1.8 7. Limitations and open questions

This is a thought experiment, not a theory. We identify the following limitations:

1. No derivation of the dimensional structure. We do not specify how many extra dimensions exist, what their shape is, or what fields inhabit the bulk. These are free parameters of the model.
2. No derivation of the inversion mechanism. We claim that the projection of bulk gravity onto our brane inverts the sign of the gravitational coupling, but we do not derive this from a specific geometry. The inversion could occur in many ways, with different observable consequences. This is a stronger claim than standard brane-world models and is more tightly constrained by data.
3. No derivation of the original events parameters. We do not specify the energy, duration, or spectral shape of the 4D event. These would need to be derived from a specific bulk geometry and field content.
4. No derivation of the dimensional time-dilation rule. The claim that a brief 4D event projects as a long 3+1 dimensional universe requires a specific rule for how 4D structure maps to 3+1 temporal extent. This rule is not specified in the model.
5. No precise identification of the products. Dark matter and dark energy are identified with specific aspects of the dimensional projection (current 2D universe gravity, and un-cancelled inversion residue respectively), but the model does not specify the proportionality constants for these identifications. The model provides a geometric role for dark matter but does not identify which specific dark matter candidate is correct.
6. No CMB power spectrum derivation. The model is constrained by the requirement that any specific implementation reproduce the observed CMB power spectrum, primordial element abundances, and large-scale structure. We have not computed these predictions from the 4D event scenario.



7. No prediction for direct-detection signals. Without a specific bulk field content and coupling structure, we cannot predict the direct-detection cross-section for dark matter or the equation of state for dark energy. The model predicts  $w = -1$  approximately (dark energy is approximately constant in our 3+1 dimensional frame, matching CDM behavior, because our universe is a brief slice of the 4D events full duration), but the absolute value of the dark energy density is not predicted. A specific implementation of the model would also need to specify the temporal profile of the 4D events antigravity output over its full duration, which determines how the dark energy density evolves on cosmological timescales.
8. The proportionality constant for the dark-matter / energetic-event-rate correlation is not specified. The model predicts that dark matter density correlates with energetic event rates, but does not specify the proportionality constant. Computing this constant requires a specific geometry and event spectrum, which we have not derived.
9. The 2D universe physics is not specified. We have not derived what physics would govern a 2D universe created by a 3+1 dimensional event. Whether 2D universes can support complex information processing (and thus beings) is an open question. The model treats them as real, with their own physics, but the details are unspecified.
10. No precise threshold or weighting function for energetic event contributions. The model proposes that all energetic events contribute to dark matter, with each events gravitational contribution weighted by its energy. The simplest reading is that the cumulative dark matter density is proportional to the average energetic event rate per unit visible mass, weighted by event energy. This is consistent with the radial acceleration relation (§4.1). However, the quantitative proportionality constant and the precise weighting function (linear in event energy? power law? something else?) are not derived. A specific geometry and event spectrum would be required to compute these quantities. The qualitative principle (dark matter tracks average activity) is robust to the choice of weighting, but the quantitative predictions depend on the details.
11. The source of the 4D events energy is not addressed. The model assumes the 4D event is an ongoing energetic process with some total energy budget  $E_{4D}$  (per §2.2). The model does not specify where this energy comes from. It is possible that the 4D event is itself a projection from a 5D (or higher) process, in which case the cascade continues upward our 3+1 dimensional universe is the projection of a 4D event, which is itself the projection of a 5D process, and so on. This is consistent with the scale-invariant principle of §2.3, but it raises the question of whether the cascade has a bottom (1D, 0D, or some fundamental level) and a top (a fundamental source of energy) or whether it extends indefinitely. The model does not currently address this. A specific implementation of the model would need to either identify the source of the 4D events energy (a bottom of the cascade) or accept the cascade as an infinite hierarchy (with all the philosophical and mathematical consequences that entails).
12. The cascade is almost exact at every level a new form of fine-tuning. The downward perceptual inversion principle (downward projection is perceived as inverted, upward back-projection is not, and the downward perceptual inversion almost-cancels with the native gravity) requires the cancellation to be almost exact at every level (per §2.4 and §2.6). This is fine-tuning in a new form: not fine-tuning of a single parameter, but fine-tuning of the structure of the dimensional cascade such that the cancellation is almost (but not quite) exact at every level. The model does not currently derive why the cascade should be almost-cancelling rather than completely cancelling (which would give no observable effects at all) or weakly cancelling

(which would give effects much larger than observed). A specific implementation would need to derive the near-exact cancellation from a more fundamental principle, which is left to future work.

13. The four-force unification is conceptual, not quantitative. §4.13–§4.15 attempt to unify gravity, electromagnetism, the weak force, and the strong force within the dimensional-cascade framework. The connections are conceptual – all four forces’ properties are reframed as consequences of the 4D event. The connections are not quantitative – the model does not derive the specific values of the coupling constants, the mass ratios of force carriers, the color charge structure, or the asymptotic freedom of the strong force. The unification is at the level of interpretation (all four forces are projections of the same 4D event) rather than at the level of prediction (the model does not compute force strengths from first principles). A specific implementation of the model would need to derive these quantitative features from the geometry, which is left to future work.
14. The mathematical sketch of the cascade has a sign ambiguity. The Mathematical sketch in §2.4 presents the cascade as  $G_{D-1}^{total} = G_{D-1}^{native} - k \cdot G_D$ , with the ordinary gravity as the small positive residue and the dark energy as a small un-cancelled antigravity. In a strict algebraic interpretation, these two quantities would have opposite signs (one is  $G_{native} - G_{proj}$ , the other is  $-(G_{proj} - G_{native})$ ), which would imply they cancel. To avoid this, we treat them as physically distinct small contributions: the ordinary gravity is a force on matter (small positive), and the dark energy is a vacuum energy (also small but as a separate physical quantity). The model proposes that both are small because of the near-cancellation of  $G_{native}$  and  $G_{proj}$ , but the exact algebraic relationship between the two small quantities is not derived in this paper. A specific implementation of the model would need to derive this relationship from a more fundamental principle.
15. The dimensional analysis requires a staying fraction postulate to bridge the  $10^{123}$  gap. The §2.6 Dimensional analysis of the cascade presents a qualitative sketch where the cascade suppression  $\epsilon \sim 10^{-38}$  at the 3+1D level explains the  $10^5$  hierarchy. The cascade’s raw prediction for the dark energy is  $\sim 10^{-38} \cdot M_{Pl}^4 \sim 10^{38}$  GeV, which is  $10^{85}$  larger than the observed  $\sim 10^{-47}$  GeV. To bridge this gap, the §2.6 introduces a staying fraction  $f_{back} \sim 10^{-85}$  (the fraction of cascade-produced antigravity that remains in 3+1D as observable dark energy), giving the math  $10^{-38} \times 10^{-85} = 10^{-123}$  in natural units, equivalent to  $\sim 10^{-47}$  GeV. The staying fraction is a postulate of the model, and is not derived from the cascade geometry. The model qualitatively claims that the dark energy is small because of the dimensional-cascade cancellation, but the quantitative value (the  $10^{-85}$  staying fraction) is a postulate, not a derivation. A complete implementation of the model would need to derive the staying fraction from first principles, which would in turn give a predictive derivation of the absolute dark energy density. The  $10^{85}$  discrepancy between the cascade’s raw prediction and the observed dark energy remains unbridged in the current model (the staying fraction is exactly the post-hoc factor that makes the math work, but it is not derived from the cascade geometry).

These limitations are not unusual for a thought experiment. They are the natural next steps for theoretical development.

## 1.9 8. Conclusion

We have proposed that gravity's observed weakness, dark matter, and dark energy are all manifestations of a single geometric process: a dimensional inversion of gravitational influence following a single ongoing energetic event in a higher-dimensional space. The universe is the projection of that event into our 3+1 dimensional spacetime. The bulk-brane gravity interaction produces two distinct observable effects: a 4D-event-driven geometric contribution (the un-cancelled fraction of the inverted bulk gravity, identified as dark energy, approximately constant in our 3+1 dimensional frame because our universe is a brief slice of the 4D events full duration) and a current rate-dependent collective effect (the gravity of 2D universes being created now by 3+1 dimensional energetic events, identified as dark matter). The universes lifetime is some fraction of the 4D events full duration in 4D time. The universe ends as a fixed-time boundary rather than a fade-out.

The model is consistent with several established research programs (brane-world cosmology, emergent gravity, the Dark Dimension scenario, holographic frameworks) and provides a unifying framing that connects three open problems under a single mechanism. The cosmological constant problem, in particular, is reframed as a misidentification: we were computing the wrong quantity (3+1 dimensional QFT zero-point energy) instead of the right one (un-cancelled inverted bulk gravity). The hierarchy problem is reframed as a bulk-brane cancellation: ordinary gravity is the small net remainder of the branes attractive gravity after cancellation with the inverted bulk gravity, so the observed gravitational coupling is small. (Dark matters apparent weakness is the same bulk-brane cancellation effect, applied to the next level of the dimensional cascade.) We do not claim to solve either problem; we claim to reframe them.

The model makes several testable predictions: the radial acceleration relation should hold (with scatter correlating with current activity, not historical); dark energy should be approximately constant in our 3+1 dimensional frame (matching standard CDM behavior, because our universe is a brief slice of the 4D events full duration); gravity should be approximately constant over cosmic time; the early-universe energy spectrum should be derivable from a 4D events structure. The universes lifetime is set by some fraction of the 4D events full duration, not by an energy budget. On galaxy scales, dark matter density should correlate with current energetic activity, not just total stellar mass.

A speculative extension suggests that the same mechanism, applied at other dimensional scales, may imply that sufficiently energetic events in our universe could produce lower-dimensional universes as their aftermath. This is the most speculative part of the proposal.

The model is not a finished theory. It is a thought experiment intended to invite the physics community to develop, refine, or refute the proposal. The most important next steps are:

- Specifying the dimensional structure and bulk field content
- Deriving the inversion mechanism from a specific geometry
- Deriving the dimensional time-dilation rule that maps the 4D events structure to our 3+1 dimensional lifetime
- Computing the predicted CMB power spectrum for a 4D event initial condition and comparing to Planck data
- Computing the predicted primordial element abundances and comparing to Big Bang nucleosynthesis observations
- Quantitatively deriving the bulk-brane cancellation fraction (related to the 10<sup>5</sup> of the hierarchy problem) and showing that the dark matters effective coupling is the same effect applied to the next level of the dimensional cascade
- Quantitatively deriving the un-cancelled fraction of the inverted bulk gravity, to predict the absolute value of the dark energy density (the 10<sup>28</sup> of the cosmological constant problem)

- Identifying specific signatures that distinguish this model from alternatives
- Searching for high-precision confirmation that dark energy is approximately constant in our 3+1 dimensional frame (matching standard CDM)
- Searching for any very-slow deviations from constant dark energy that would correspond to the 4D events antigravity output slowly varying over its full duration
- Searching for sub-millimeter gravity deviations and gravitational wave signatures of the inversion mechanism
- Computing the proportionality constant and weighting function for the dark-matter / energetic-event-rate correlation (i.e., the quantitative form of how dark matter depends on the average activity per unit visible mass)
- Testing the current-activity correlation on galaxy surveys, especially for mass-matched pairs with different stellar densities
- Developing the physics of 2D universes created by 3+1 dimensional events

We are not specialists in theoretical physics. We offer this proposal with the hope that it may be useful, and with the appropriate humility about its status as a thought experiment rather than a developed theory.

---

## 1.10 References

[ADD98] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, The Hierarchy Problem and New Dimensions at a Millimeter, *Phys. Lett. B* 429 (1998) 263-272.

[Desmond25] H. Desmond, Modified Newtonian Dynamics: Observational Successes and Failures, arXiv:2505.21638 (2025).

[Golini24] G. Golini, M. Montes, E. R. Carrasco, J. Román, I. Trujillo, Ultra-deep imaging of NGC1052-DF2 and NGC1052-DF4 to unravel their origins, *Astronomy & Astrophysics* 684, A99 (2024).

[Gregory00] R. Gregory, V. A. Rubakov, S. M. Sibiryakov, Brane worlds: the gravity of escaping matter, *Class. Quantum Grav.* 17 (2000) 4437-4450.

[Júlio25] M. P. Júlio, J. I. Read, M. S. Pawłowski, P. Li, D. Vaz, J. Brinchmann, M. P. Rey, O. Agertz, T. Holmes, The radial acceleration relation at the EDGE of galaxy formation: testing its universality in low-mass dwarf galaxies, arXiv:2510.06905 (2025).

[Kravtsov24] A. Kravtsov, On the dark matter content of ultra-diffuse galaxies, arXiv:2406.13732 (2024).

[LawSmith24] J. A. P. Law-Smith, G. Obied, A. Prabhu, C. Vafa, Astrophysical Constraints on Decaying Dark Gravitons, arXiv:2307.11048 (2024).

[Maldacena97] J. M. Maldacena, The Large N Limit of Superconformal Field Theories and Supergravity, *Int. J. Theor. Phys.* 38 (1999) 1113-1133.

[McGaugh16] S. S. McGaugh, F. Lelli, J. M. Schombert, Radial Acceleration Relation in Rotationally Supported Galaxies, *Phys. Rev. Lett.* 117 (2016) 201101.

[Mercado24] F. J. Mercado et al., Hooks & Bends in the radial acceleration relation, *MNRAS* 530, 1349 (2024).

[Mistele24] T. Mistele, S. McGaugh, F. Lelli, J. Schombert, P. Li, Radial acceleration relation of galaxies with joint kinematic and weak-lensing data, arXiv:2310.15248 (2024).

[Obied23] G. Obied, C. Dvorkin, E. Gonzalo, C. Vafa, Dark Dimension and Decaying Dark Matter Gravitons, arXiv:2311.05318 (2023).

- [RS99] L. Randall, R. Sundrum, An Alternative to Compactification, *Phys. Rev. Lett.* 83 (1999) 4690-4693.
- [Tetradis04] N. Tetradis, Brane-world evolution with brane-bulk energy exchange, *hep-th/0414282* (2004).
- [Tian24] Y. Tian, H. Ryu, A distinct radial acceleration relation across the brightest cluster galaxies, *Astronomy & Astrophysics* (2024).
- [Vrteanu25] A. A. Vrteanu, M. J. Jarvis, A. A. Ponomareva, H. Desmond, I. Heywood, T. Yasin, N. Maddox, M. Glowacki, M. Maksymowicz-Maciata, P. E. Mancera Piña, H. Pan, MIGHTEE-HI: The radial acceleration relation with resolved stellar mass measurements, *arXiv:2504.20857* (2025).
- [Verlinde16] E. P. Verlinde, Emergent Gravity and the Dark Universe, *SciPost Phys.* 2 (2016) 016.
- [Yousef13] L. Yousef, A. Sheykhi, QCD Ghost Dark Energy in RS II Braneworld with Bulk-Brane Interaction, *Int. J. Theor. Phys.* 53 (2014) 1472-1482.
- [Borah25] D. Borah, N. Das, R. Roshan, Evolving Dark Sector and the Dark Dimension Scenario, *arXiv:2507.03090* (2025).

---

## 1.11 Appendix: Authors Note

The author is a software engineer, not a physicist. This model emerged from asking questions in conversation, not from working through the formalism. The intent is to propose a unifying framing for several open problems in fundamental physics, not to claim a finished theory. The author offers this proposal in the hope that someone with formal training will either develop it further or definitively refute it.

The unification of dark matter, dark energy, and the hierarchy problem under a single geometric process is a coherent hypothesis worth testing. The model is not presented as true; it is presented as a candidate framework that, if developed rigorously, would make specific predictions distinguishable from standard alternatives. The misidentification reframe of the cosmological constant problem, the bulk-brane cancellation reframe of the hierarchy problem, the scale-invariant principle of universe creation, and the diffuse-galaxy prediction are the most novel and testable elements.

If you are a physicist reading this and finding it interesting: please work on it. If you are finding it naive: please be specific about where it fails. Both responses are useful.

---

Manuscript date: 2026-06-10 Correspondence: open