

## Differentiation Rules Part Two

---

```
1 caseInsensitive = function(a,b) {
2     return a.toLowerCase() == b.toLowerCase();
3 };
```

---

**Julia:** You know, some of those rules we learned were pretty useful, but some of these derivatives still suck! There **HAS** to be a better way!

**Dylan:** I'm sure there is, and I'm sure I know who could help us!

**James:** Did I hear my name?

**Dylan:** Not yet!

**Julia:** James!

**James:** There are more rules for differentiation that can make your life just a little bit easier!

## The Product Rule

**James:** From the last time we did this, what rule do you think would exist for the product of two functions?

**Julia:** Well, last time we added or subtracted the derivative of both functions, so I bet we multiply the derivative of both!

**Dylan:** Let's check!

Consider the functions  $f(x) = 2x$  and  $g(x) = 3x^3 + x^2$ .

Graph of  $f(x) = 2x, g(x) = 3x^3 + x^2$

**Question 1** Use Julia's guess to find the derivative of  $f(x) \cdot g(x)$ .

$18x^2 + 4x$

---

Learning outcomes:

**Definition 1.** The **derivative** of  $f(x)$  at  $a$  is defined by the following limit:

$$\left[ \frac{d}{dx} f(x) \right]_{x=a} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}.$$

Use the limit definition of the derivative to find the derivative of  $f(x) \cdot g(x)$ .

$$\boxed{24x^3 + 6x^2}$$

Was Julia right?

**Multiple Choice:**

- (a) Yes
- (b) No ✓

**Julia:** Darn! It didn't work!

**Dylan:** It must be a little harder than that...

**James:** That's right Dylan, but it is easier than the limit definition! All we have to do is use

$$\frac{d}{dx} (f(x) \cdot g(x)) = f(x) \cdot g'(x) + f'(x) \cdot g(x).$$

This is called the **Product Rule**.

**Question 2** Using the Product Rule, differentiate the products of the following functions:

$$f(x) = 6x^3, g(x) = 7x^4$$

$$\boxed{294x^6}$$

$$f(x) = \cos(x) + 4x, g(x) = 3x^2$$

$$\boxed{-3x^2 \sin(x) + 6x \cos(x)}$$

$$f(x) = x^2, g(x) = 3x^3 - 3x$$

$$\boxed{15x^4 - 9x^2}$$

$$f(x) = x^7, g(x) = 2x^{32}$$

$$\boxed{78x^{38}}$$

## The Quotient Rule

**Dylan:** Wow! That's gonna save a ton of time with products! Is there anything like it we can do with quotients?

**James:** There is! It's even called **the Quotient Rule!**

**Julia:** I bet it's a pain too though, just like the product rule.

**James:** Well, why don't you try using your intuition first rather than guessing?

**Dylan:** Alright, well, I guess I would divide the derivative of the numerator by the derivative of the denominator.

**Question 3** Consider the functions  $f(x) = x^3 + 1$  and  $g(x) = x$ .

Graph of  $f(x) = x^3 + 1, g(x) = x$

Use Dylan's guess to find the derivative of  $\frac{f(x)}{g(x)}$ .

Use the limit definition of the derivative to find the derivative of  $\frac{f(x)}{g(x)}$ .

Was Dylan right?

**Multiple Choice:**

- (a) Yes
- (b) No ✓

**Julia:** I knew it! It's never that easy!

**James:** Now calm down Julia, this rule is worse than the last one, but it's much better than going through by the limit definition:

$$\frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}.$$

**Question 4** Using the Quotient Rule, differentiate the products of the following functions to find  $\frac{d}{dx} \left[ \frac{f(x)}{g(x)} \right]$ :

$$f(x) = 3x - 1, g(x) = 2x + 1$$

$$\boxed{5/(2x + 1)^2}$$

$$f(x) = 1, g(x) = x + 10$$

$$\boxed{-1/(x + 10)^2}$$

$$f(x) = x^2, g(x) = 3x^3 - 3x$$

$$\boxed{(-x^2 - 1)/(3(x^2 - 1)^2)}$$

$$f(x) = x^7, g(x) = 2x^{32}$$

$$\boxed{-25/(2x^{26})}$$

## The Chain Rule

**James:** There's one last rule to learn today; the **Chain Rule**.

**Dylan:** That rule sounds pretty cool! When do we use it though? I thought we already covered the functions we need to know...

**Julia:** Yeah, what else is there?

**James:** We use the chain rule in composition of functions, like when we have  $\sin(2x)$  -  $2x$  is a function, and so is  $\sin(x)$

**Julia:** And how bad is the rule?

**James:** This one is a little more tricky -

$$\frac{d}{dx} f(g(x)) = f'(g(x)) \cdot g'(x).$$

**Dylan and Julia:** That's so gross.

**James:** Well, let's give it a try and see if you like it more than the limit definition!

**Question 5** Consider  $f(x) = \sqrt{x}$  and  $g(x) = \frac{1}{x}$

Graph of  $\text{sqrt}(x), 1/x$

Using the limit definition of derivative, evaluate the derivative of  $f(g(x))$ .

$$\boxed{-1/2x^{3/2}}$$

Now, evaluate the same limit using the chain rule. Notice you get the same answer. Yay.

**Question 6** Find the composition  $f(g(x))$ , then using the Chain Rule, differentiate  $f(g(x))$  for the following functions:

$$f(x) = 3x + x^2, g(x) = x^4 + 7x$$

$$f(g(x)) = \boxed{3(x^4 + 7x) + (x^4 + 7x)^2} \quad \frac{d}{dx} \left[ f(g(x)) \right] = \boxed{8x^7 + 70x^4 + 12x^3 + 98x + 21}$$

$$f(x) = \cos(x), g(x) = \sin(x)$$

$$f(g(x)) = \boxed{\cos(\sin(x))} \quad \frac{d}{dx} \left[ f(g(x)) \right] = \boxed{-\cos(x) \sin(\sin(x))}$$

$$f(x) = \cos(x), g(x) = x^3$$

$$f(g(x)) = \boxed{\cos(x^3)} \quad \frac{d}{dx} \left[ f(g(x)) \right] = \boxed{-3x^2 \sin(x^3)}$$

**Question 7** Using the Chain Rule, differentiate the compositions  $g(f(x))$  for the following functions:

$$f(x) = 3x + x^2, g(x) = x^4 + 7x$$

$$g(f(x)) = \boxed{(3x + x^2)^4 + 7(3x + x^2)} \quad \frac{d}{dx} \left[ g(f(x)) \right] = \boxed{(2x + 3)(4x^3(x + 3)^3 + 7)}$$

$$f(x) = \cos(x), g(x) = \sin(x)$$

$$g(f(x)) = \boxed{\sin(\cos(x))} \quad \frac{d}{dx} \left[ g(f(x)) \right] = \boxed{-\cos(\cos(x)) \sin(x)}$$

$$f(x) = \cos(x), g(x) = x^3$$

$$g(f(x)) = \boxed{\cos(x)^3} \quad \frac{d}{dx} \left[ g(f(x)) \right] = \boxed{-3 \cos(x)^2 \sin(x)}$$