

Differentiation Rules!

Julia: Hmm...I don't think differentiation rules, it takes so long and I hate using that long limit definition!

Dylan: No no Julia, it's differentiation *rules*!

Julia: Ohhhh, that makes more sense!

The Power Rule

Julia: I hate how long it takes to differentiate powers!

Dylan: Yeah, it takes forever! I feel like there was some sort of pattern to it, but I couldn't figure anything out.

James: Sounds like you guys need my help again?

Julia and Dylan: Help us James!

James: There *is* a pattern! Check out this table I made!

$f(x)$	$\frac{d}{dx} f(x)$
x^2	$2x^1$
x^3	$3x^2$
x^4	$4x^3$

Question 1 What pattern do you notice in James' table? Generalize this pattern in terms of x^n .

Multiple Choice:

- (a) $n \cdot x^{n-1}$ ✓
- (b) $n - 1 \cdot x^{n-1}$
- (c) $n \cdot x^n$
- (d) $n - 1 \cdot x^n$

Learning outcomes:

Question 2 Using the limit definition of a derivative, compute the derivative for x^3 .

$$\frac{d}{dx}x^5 = \boxed{5x^4}$$

Feedback (correct): Notice that your answer fits the same pattern as before!

Question 3 Use the power rule to differentiate the following functions.

$$f(x) = x^{10} \quad \frac{d}{dx}f(x) = \boxed{10x^9}$$

$$f(x) = 3x^2 \quad \frac{d}{dx}f(x) = \boxed{6x}$$

Hint: The value $\frac{1}{x}$ can be represented by x^{-1} .

$$f(x) = \frac{5}{x} \quad \frac{d}{dx}f(x) = \boxed{-5x^{-2}}$$

The Constant Rule

Dylan: Wow! That's neat!

Julia: I wish we could use rules like this all over the place though, it would really save me time.

James: There are plenty of places with rules like this! Why don't we look at a function like $y = 3$?

Consider $y = c$, where c is some arbitrary constant.

Question 4

Differentiate this function using the limit definition.

$$\frac{d}{dx}c = \boxed{0}$$

What can you generalize about the derivative of $y = c$ based on this?

Multiple Choice:

(a) $\frac{d}{dx}c = 2c$

(b) $\frac{d}{dx}c = 0$ ✓

(c) $\frac{d}{dx}c = x$

(d) $\frac{d}{dx}c = \frac{c}{2}$

Question 5 Using what you found in the previous problem, compute the following derivatives:

$f(x) = 2$ $\frac{d}{dx}f(x) = \boxed{0}$

$f(x) = 100$ $\frac{d}{dx}f(x) = \boxed{0}$

$f(x) = 0$ $\frac{d}{dx}f(x) = \boxed{0}$

The Constant Multiple Rule

Julia: James! Show us more! These things are going to save me so much time on my homework!

James: Alright alright, calm down Julia. We can look at a function like $y = 3x$ next.

Consider $y = k \cdot x$, where k is some arbitrary constant.

Question 6 $\frac{d}{dx}(k \cdot x) = \boxed{k}$

What can you generalize about the derivative of $y = kx$ based on this?

Multiple Choice:

(a) $\frac{d}{dx}kx = 2k$

$$(b) \frac{d}{dx} kx = kx^2$$

$$(c) \frac{d}{dx} kx = k \checkmark$$

$$(d) \frac{d}{dx} kx = x$$

Question 7

Using what you found in the previous problem, compute the following derivatives:

$$f(x) = 4x \qquad \frac{d}{dx} f(x) = \boxed{4}$$

$$f(x) = 10x \qquad \frac{d}{dx} f(x) = \boxed{10}$$

$$f(x) = \frac{1}{5}x \qquad \frac{d}{dx} f(x) = \boxed{\frac{1}{5}}$$

The Sum and Difference Rules

Dylan: Wow, this stuff is awesome! Is there any way to put it all together? Like, is there an easy way to tell what the derivative of $f(x) = 3x + 4$ is?

James: There is Dylan!

Question 8 Consider the differentiable functions $f(x)$ and $g(x)$. We will define a function $j(x) = f(x) + g(x)$.

Hint: In $j(x + h)$, the $(x + h)$ will replace x in the component functions as well.

Take the derivative of $j(x)$ using the limit definition.

$$j'(x) = \boxed{\frac{j(x+h) - j(x)}{h}}$$

What can you generalize based on this?

Multiple Choice:

$$(a) \frac{d}{dx} f(x) + g(x) = f'(x) - g'(x)$$

$$(b) \frac{d}{dx} f(x) + g(x) = g'(x) + f'(x)$$

$$(c) \frac{d}{dx} f(x) + g(x) = f(x) \cdot g'(x) - g(x) \cdot f'(x)$$

$$(d) \frac{d}{dx} f(x) + g(x) = f'(x) + g'(x) \checkmark$$

Question 9 Using what you found in the previous problem, compute the following derivatives:

$$f(x) = 3x^2 - 5x + 2, g(x) = x^2 + 3x$$

$$\frac{d}{dx} j(x) = \boxed{8x - 2}$$

$$f(x) = x^2 - 4x + 2, g(x) = -4x^2 + 3$$

$$\frac{d}{dx} j(x) = \boxed{-6x - 4}$$

$$f(x) = 5x^3 + 3x, g(x) = 2x^2 - 13x$$

$$\frac{d}{dx} j(x) = \boxed{15x^2 + 4x - 10}$$

Question 10 Julia wonders if a similar rule exists for $m(x) = f(x) - g(x)$. Using the limit definition of derivative, determine if there is a pattern. Then, if there is a rule, use it to solve the 1a, 1b, and 1c. If there is not, do them using the limit definition.

$$f(x) = 3x^2 - 5x + 2, g(x) = x^2 + 3x$$

$$\frac{d}{dx} m(x) = \boxed{4x - 8}$$

$$f(x) = x^2 - 4x + 2, g(x) = -4x^2 + 3$$

$$\frac{d}{dx} m(x) = \boxed{10x - 4}$$

$$f(x) = 5x^3 + 3x, g(x) = 2x^2 - 13x$$

$$\frac{d}{dx} m(x) = \boxed{15x^2 - 4x + 16}$$

In Summary

We've covered a lot of differentiation rules in this lab, to help you out we've made the following table.

Question 11

Power Rule	$\frac{d}{dx} x^n = \boxed{n} \boxed{*} \boxed{x^{n-1}}$, where n is any real number.
Constant Rule	$\frac{d}{dx} c = \boxed{0}$
Constant Multiple Rule	$\frac{d}{dx} (c \cdot f(x)) = \boxed{c} \boxed{*} \frac{d}{dx} \boxed{f(x)}$
Sum Rule	$\frac{d}{dx} (f(x) + g(x)) = \frac{d}{dx} \boxed{f(x)} \boxed{+} \frac{d}{dx} \boxed{g(x)}$
Difference Rule	$\frac{d}{dx} (f(x) - g(x)) = \frac{d}{dx} \boxed{f(x)} \boxed{-} \frac{d}{dx} \boxed{g(x)}$