

Implicit Differentiation - Finish solutions

Dylan: Woah! What's up with this?

Julia: I didn't know functions were explicit!

Dylan: The x and y are on the same side of the equation! I can't deal with this.

James: Functions can be explicit or implicit! And it not the way you're thinking Julia...

Introduction

So far we have dealt only with explicitly defined functions, where $y = f(x)$. Functions where there are both x and y on one side or both sides of the equation are called **implicit functions**.

Guided Example

Question 1 Which of the following equations are defined implicitly?

Select All Correct Answers:

(a)

$$y = x^2 + 5x - 7$$

(b)

$$y = \sin(x)$$

(c)

$$x^2 + y^2 = 1$$

✓

(d)

$$y = \text{sqrt}(x - 3)$$

Learning outcomes:

(e) $x^2y^3 + y = 5x + 8y$ ✓

Question 2 Graph the following implicitly defined function below,

$$x^2 + y^2 = 1$$

Graph of

Now, in the following sage cell, solve the function for y. For help using the solve command refer to the [documentation](#) here.

SAGE

```

1 x,y = var("x, y")
2 #eqn = x**2+y**2==1, this sets eqn to the unit circle
3 #use the solve command to solve eqn for y

```

Graph the two explicit equations on the same axis below.

Graph of

Which of the following are true?

Select All Correct Answers:

(a)

$$x^2 + y^2 = 1$$

is a function

(b)

$$-/\sqrt{1-x^2} \text{ is a function}$$

✓

(c)

$$/\sqrt{1-x^2} \text{ is a function}$$

✓

Question 3 Using the function(s) you found, find the slope of the tangent lines at a point with two values on your graph.

Free Response:

Implicit Differentiation Using Substitution

Consider the equation $-x^2 \cdot y^3 + y^5 - 32 = 0$.

Question 1 Using the method shown in the previous section, evaluate the function for y . Does this equation look easy to differentiate? No

Instead, let's treat our equation as an expression - because it equals zero, we don't have to worry about moving anything over. Now consider y as $y(x)$, a function of x , and differentiate with respect to x . Each y term will gain $\frac{dy}{dx}$. Then, set the expression equal to zero, and solve for $\frac{dy}{dx}$. What does this represent?

Free Response:

Question 2 Using your result in the previous section, evaluate $\frac{dy}{dx}$ at $x = 3$ and $x = 7$.

$x = 3$:

$x = 7$:

1 **Question 3** 2 SAGE

Now use Sage Math to find the slope of $\sin(x^2) = \cos(xy^2)$ at any point. Look here for information on implicit differentiation in Sage

☐

Perpendicular at a Point

Julia: Wow, implicit differentiation is rough.

Dylan: You're telling me... I've been doing this for hours! I wish we could at least do a little more with it if I have to learn it.

James: Did I hear that you guys want to know more about using implicit differentiation?

Julia and Dylan: James! Tell us more!

James: Alright guys, you can use implicit differentiation with implicit functions to tell if two functions are perpendicular at a point!

Julia: But how?

Dylan: Yeah, I don't see how that helps.

James: It's easy - all we have to do is see if the tangent lines are perpendicular at that point, and if they are, then so are the curves!

Question 4 Graph $3x - 2y + x^3 - x^2y = 0$ and $x^2 - 2x + y^2 - 3y = 0$ on the same set of axes.

Graph of

Do they look perpendicular anywhere?

Question 5 Hint: Remember, two perpendicular lines will have their slopes be the negative inverse of one another.

Prove the two curves are (or are not) perpendicular at the origin.

Free Response:

In Summary

There are two main methods to solve implicit equations

- (a) Solve for y and then differentiate.
- (b) Treat y as $y(x)$ and differentiate for x , eventually solving for $\frac{dy}{dx}$ to give the value of the derivative at any point.