

Differentiation Rules

Julia: Hmm...I don't think differentiation rules, it takes so long and I hate using that long limit definition!

Dylan: No no Julia, it's differentiation *rules*!

Julia: Ohhhh, that makes more sense!

The Power Rule

Julia: I hate how long it takes to differentiate powers!

Dylan: Yeah, it takes forever! I feel like there was some sort of pattern to it though. I couldn't figure anything out though.

James: Sounds like you guys need my help again?

Julia and Dylan: Help us James!

James: There *is* a pattern! Check out this table I made!

| $f(x)$ | $f'(x)$ |
|--------|---------|
| x^2 | $2x^1$ |
| x^3 | $3x^2$ |
| x^4 | $4x^3$ |

Question 1 What pattern do you notice in James' table?

Free Response: This is the model solution

Question 2 Generalize this pattern in terms of x^n

$$\frac{\partial}{\partial x} x^n = \boxed{n * x^{(n-1)}}$$

Question 3 Using the limit definition of a derivative, compute the derivative for x^5

$$\frac{\partial}{\partial x} x^5 = \boxed{5x^4}$$

The Constant Rule

Dylan: Wow! That's neat!

Julia: I wish we could use rules like this all over the place though, it would really save me time.

James: There are plenty of places with rules like this! Why don't we look at a function like $y = 3$?

Consider $y = c$, where c is some arbitrary constant.

Question 4 *Derive this function using the limit definition. What does your answer mean?*

Free Response: *This is the model solution*

Question 5 *Using what you found in the previous problem, compute the following derivatives:*

(a) $f(x) = 2$

(b) $f(x) = 100$

(c) $f(x) = 0$

$$\frac{\partial}{\partial x}(x = 2) = \boxed{2x \sin(y)}$$

Free Response: *This is the model solution*

Free Response: *This is the model solution*

Free Response: *This is the model solution*

The Constant Multiple Rule

Julia: James! Show us more! These things are going to save me so much time on my homework!

James: Alright alright, calm down Julia. We can look at a function like $y = 3x$ next.

Consider $y = kx$, where k is some arbitrary constant.

Question 6

Derive this function using the limit definition. What does your answer mean?

Free Response: This is the model solution

Question 7

Using what you found in the previous problem, compute the following derivatives:

$$f(x) = 4x$$

$$f(x) = 10x$$

$$f(x) = \frac{1}{5}x$$

Free Response: This is the model solution

The Sum and Difference Rules

Dylan: Wow, this stuff is awesome! Is there any way to put it all together?

Like, is there an easy way to tell what the derivative of $f(x) = 3x + 4$ is?

James: There is Dylan!

Consider the differentiable functions $f(x)$ and $g(x)$. We will define a function $j(x) = f(x) + g(x)$.

- (a) Take the derivative of $j(x)$ using the limit definition. What does your answer mean? *Hint: In $j(x+h)$, the $(x+h)$ will replace x in the component functions as well.*
- (b) Using what you found in the previous problem, compute the following derivatives:
 - (i) $f(x) = 3x^2 - 5x + 2$, $g(x) = x^2 + 3x$
 - (ii) $f(x) = x^2 - 4x + 2$, $g(x) = -4x^2 + 3$
 - (iii) $f(x) = 5x^3 + 3x$, $g(x) = 2x^2 - 13x$

- (c) Julia wonders if a similar rule exists for $j(x) = f(x) - g(x)$. Using the limit definition of derivative, determine if there is a pattern. Then, if there is a rule, use it to solve the 1a, 1b, and 1c. If there is not, do them using the limit definition.

In Summary

We've covered a lot of differentiation rules in this lab, to help you out we've made the following table.

| | |
|------------------------|---|
| Power Rule | $\frac{d}{dx}(x^n) = n * x^{(n-1)}$, where n is any real number besides 0. |
| Constant Rule | $\frac{d}{dx}(c) = 0$ |
| Constant Multiple Rule | $\frac{d}{dx}(cf(x)) = c \frac{d}{dx}f(x)$ |
| Sum Rule | $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$ |
| Difference Rule | $\frac{d}{dx}(f(x) - g(x)) = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$ |