LECTURE 16: NONLINEAR AND POLYNOMIAL MODELS

ECON 480 - ECONOMETRICS - FALL 2018

Ryan Safner

November 26, 2018



Polynomial Functions

Quadratic Model

Determining if (Larger) Polynomials are Necessary




 $\cdot\,$ OLS is commonly known as "linear regression" as it fits a straight line to data points




- · OLS is commonly known as "linear regression" as it fits a straight line to data points
- · Often, data and relationships between variables may not be linear




Nonlinearities? Example




NONLINEARITIES? EXAMPLE: QUADRATIC FIT




NONLINEARITIES? EXAMPLE: NO OUTLIERS



NONLINEARITIES? EXAMPLE: QUADRATIC FIT



NONLINEARITIES? EXAMPLE: LOGARITHMIC FIT



• Despite being "linear regression", OLS can handle this with an easy fix



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$\checkmark$$



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$Y = \beta_0 + \beta_1^2 X$$

$$\checkmark$$

$$Y = \beta_0 + \beta_1^2 X$$



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$Y = \beta_0 + \beta_1^2 X$$

$$Y = \beta_0 + \beta_1 \sqrt{X}$$



- Despite being "linear regression", OLS can handle this with an easy fix
- · OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$Y = \beta_0 + \beta_1^2 X$$

$$Y = \beta_0 + \beta_1 \sqrt{X}$$

$$Y = \beta_0 + \sqrt{\beta_1} X$$

$$X$$



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$Y = \beta_0 + \beta_1^2 X$$

$$Y = \beta_0 + \beta_1 \sqrt{X}$$

$$Y = \beta_0 + \sqrt{\beta_1} X$$

$$X$$

$$Y = \beta_0 + \beta_1 X_1 * X_2$$



- · Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2 \qquad \qquad \checkmark$$

$$Y = \beta_0 + \beta_1^2 X \qquad \qquad x$$

$$Y = \beta_0 + \beta_1 \sqrt{X} \qquad \qquad \checkmark$$

$$Y = \beta_0 + \sqrt{\beta_1} X \qquad \qquad x$$

$$Y = \beta_0 + \beta_1 X_1 * X_2 \qquad \qquad \checkmark$$

$$Y = \beta_0 + \beta_1 \ln(X) \qquad \qquad \checkmark$$



- Despite being "linear regression", OLS can handle this with an easy fix
- OLS requires all parameters (i.e. the β 's) to be linear, the regressors (X's) can be nonlinear:

$$Y = \beta_0 + \beta_1 X^2$$

$$Y = \beta_0 + \beta_1^2 X$$

$$Y = \beta_0 + \beta_1 \sqrt{X}$$

$$Y = \beta_0 + \sqrt{\beta_1} X$$

$$Y = \beta_0 + \beta_1 X_1 * X_2$$

$$Y = \beta_0 + \beta_1 \ln(X)$$

· In the end, X will always be just a number, OLS can always estimate parameters for numbers



• Plotting the points (X, \hat{Y}) can result in a curve for nonlinear X's

• The effect of $X \rightarrow Y$ may be nonlinear if:



• The effect of $X \to Y$ may be nonlinear if:

1. $X \rightarrow Y$ is different for different levels of X



- The effect of $X \to Y$ may be nonlinear if:
- 1. $X \rightarrow Y$ is different for different levels of X
 - e.g. Diminishing returns: \(\gamma\) X increases Y at a decreasing rate




- The effect of $X \rightarrow Y$ may be nonlinear if:
- 1. $X \rightarrow Y$ is different for different levels of X
 - e.g. Diminishing returns: \(\gamma\) X increases Y at a decreasing rate
 - e.g. Increasing returns: $\uparrow X$ increases Y at an increasing rate




- The effect of $X \to Y$ may be nonlinear if:
- 1. $X \rightarrow Y$ is different for different levels of X
 - e.g. Diminishing returns: \(\gamma\) X increases Y at a decreasing rate
 - e.g. Increasing returns: † X increases Y at an increasing rate
- 2. $X \rightarrow Y$ depends on the value of X_2 e.g. interaction terms (last lesson)






Linear: slope $eta_1 = rac{\Delta Y}{\Delta X}$ is constant








Quadratic: slope eta_1 depends on value of X







POLYNOMIAL FUNCTIONS OF X

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \dots + \hat{\beta}_r X_i^r + \epsilon_i$$

• r is the highest power X_i is raised to (e.g. quadratic r = 2, cubic r = 3, etc)



POLYNOMIAL FUNCTIONS OF X

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \dots + \hat{\beta}_r X_i^r + \epsilon_i$$

- r is the highest power X_i is raised to (e.g. quadratic r=2, cubic r=3, etc)
 - \cdot The graph of an $r^{ ext{th}}$ -degree polynomial function has r-1 bends



POLYNOMIAL FUNCTIONS OF X

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \dots + \hat{\beta}_r X_i^r + \epsilon_i$$

- r is the highest power X_i is raised to (e.g. quadratic r = 2, cubic r = 3, etc)
 - \cdot The graph of an r^{th} -degree polynomial function has r-1 bends
- Just another multivariate OLS regression model



$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

• Quadratic model has X and X^2 variables in it (yes, need both!)



$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

- Quadratic model has X and X^2 variables in it (yes, need both!)
- · How to interpret coefficients?



$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

- Quadratic model has X and X^2 variables in it (yes, need both!)
- · How to interpret coefficients?
 - \cdot β_0 as "intercept" and β_1 as "slope" makes no sense



$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

- Quadratic model has X and X^2 variables in it (yes, need both!)
- · How to interpret coefficients?
 - $\cdot \beta_0$ as "intercept" and β_1 as "slope" makes no sense
 - β_1 as effect $X_i \to Y_i$ holding X_i^2 constant makes no sense



$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

- Quadratic model has X and X^2 variables in it (yes, need both!)
- · How to interpret coefficients?
 - $\cdot \beta_0$ as "intercept" and β_1 as "slope" makes no sense
 - β_1 as effect $X_i \to Y_i$ holding X_i^2 constant makes no sense
 - Note: this is *not* a multicollinearity problem! Correlation only measures *linear* relationships!



QUADRATIC MODEL

$$\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i + \hat{\beta}_2 X_i^2 + \epsilon_i$$

- Quadratic model has X and X^2 variables in it (yes, need both!)
- · How to interpret coefficients?
 - $\cdot \beta_0$ as "intercept" and β_1 as "slope" makes no sense
 - β_1 as effect $X_i \to Y_i$ holding X_i^2 constant makes no sense
 - · Note: this is *not* a multicollinearity problem! Correlation only measures *linear* relationships!
 - Calculate marginal effects by calculating predicted \hat{Y}_i for different X_i



QUADRATIC MODEL: CALCULATING MARGINAL EFFECTS

· What is the effect of $\Delta X_i \rightarrow \Delta Y_i$?



QUADRATIC MODEL: CALCULATING MARGINAL EFFECTS

- What is the effect of $\Delta X_i \rightarrow \Delta Y_i$?
- Take the **derivative** of Y_i with respect to X_i :

$$\frac{dY_i}{dX_i} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$



QUADRATIC MODEL: CALCULATING MARGINAL EFFECTS

- What is the effect of $\Delta X_i \rightarrow \Delta Y_i$?
- Take the **derivative** of Y_i with respect to X_i :

$$\frac{dY_i}{dX_i} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$

• Marginal effect of a 1 unit change in X_i is a $\hat{\beta}_1 + 2\hat{\beta}_2 X_i$ unit change in Y



QUADRATIC MODEL: EXAMPLE

Example

Life
$$\widehat{\mathsf{Expectancy}}_i = \hat{eta}_0 + \hat{eta}_1 \mathsf{GDP}_i + \hat{eta}_2 \mathsf{GDP}_i^2$$

- Life Expectancy
- GDP per Capita (GDP for short)



 These coefficients will be very large, let's first transform gdpPercap into \$1,000s, call it gdp.t ¹



 $^{^{1}\}mbox{Note I}$ am using \mbox{dplyr} and %>% here for efficiency, I loaded them before without showing it

 These coefficients will be very large, let's first transform gdpPercap into \$1,000s, call it gdp.t ¹

```
gapminder <- gapminder %>%
  mutate(gdp.t=gdpPercap/1000)
```



 $^{^{1}\}mbox{Note I}$ am using \mbox{dplyr} and %>% here for efficiency, I loaded them before without showing it

 These coefficients will be very large, let's first transform gdpPercap into \$1,000s, call it gdp.t ¹

```
gapminder <- gapminder %>%
  mutate(gdp.t=gdpPercap/1000)
```

Let's also make the quadratic term by squaring gdp.t and calling it gdp.sq



 $^{^{1}\}mbox{Note I}$ am using \mbox{dplyr} and %>% here for efficiency, I loaded them before without showing it

• These coefficients will be very large, let's first transform gdpPercap into \$1,000s, call it gdp.t ¹

```
gapminder <- gapminder %>%
mutate(gdp.t=gdpPercap/1000)
```

Let's also make the quadratic term by squaring gdp.t and calling it gdp.sq

```
gapminder <- gapminder %>%
  mutate(gdp.sq=gdp.t^2)
```



¹Note I am using dplyr and %>% here for efficiency, I loaded them before without showing it

QUADRATIC MODEL: EXAMPLE REGRESSION IN R

- Can "manually" run regression with ${\tt gdp.t}$ and squared term ${\tt gdp.sq}$



QUADRATIC MODEL: EXAMPLE REGRESSION IN R

· Can "manually" run regression with gdp.t and squared term gdp.sq

Signif codes: 0 '+++' 0 001 '++' 0 01 '+' 0 05 ' ' 0 1 ' 1 1

```
reg1<-lm(lifeExp~gdp.t+gdp.sq, data=gapminder)</pre>
summary(reg1)
##
## Call:
## lm(formula = lifeExp ~ gdp.t + gdp.sq, data = gapminder)
##
## Residuals:
##
       Min
                 10 Median
                                  30
                                          Max
## -28.0600 -6.4253 0.2611 7.0889 27.1752
##
## Coefficients:
##
                Estimate Std. Error t value Pr(>|t|)
## (Intercept) 50.5240058 0.2978135 169.65 <2e-16 ***
## gdp.t 1.5509911 0.0373735 41.50 <2e-16 ***
## gdp.sq -0.0150193 0.0005794 -25.92 <2e-16 ***
## ---
```

QUADRATIC MODEL: EXAMPLE REGRESSION IN R II

 $\boldsymbol{\cdot}$ Or can use one of R's shortcuts for polynomial models:



QUADRATIC MODEL: EXAMPLE REGRESSION IN R II

- Or can use one of R's shortcuts for polynomial models:
- first term is normal, x, second term use $I(x^2)$ to add squared term



QUADRATIC MODEL: EXAMPLE REGRESSION IN R II

- Or can use one of R's shortcuts for polynomial models:
- first term is normal, x, second term use $I(x^2)$ to add squared term

```
reg1<-lm(lifeExp~gdp.t*I(gdp.t^2), data=gapminder)
summary(reg1)</pre>
```

```
##
## Call:
## lm(formula = lifeExp ~ gdp.t + I(gdp.t^2), data = gapminder)
##
## Residuals:
       Min
                 10 Median
                                  30
                                          Max
## -28.0600 -6.4253 0.2611
                             7 8889 27 1752
##
## Coefficients:
                Estimate Std. Error t value Pr(>|t|)
## (Intercept) 50.5240058 0.2978135 169.65 <2e-16 ***
## gdp.t
          1.5509911 0.0373735 41.50 <2e-16 ***
## I(gdp.t^2) -0.0150193 0.0005794 -25.92 <2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 8.885 on 1701 degrees of freedom
## Multiple R-squared: 0.5274, Adjusted R-squared: 0.5268
## F-statistic: 949.1 on 2 and 1701 DF, p-value: < 2.2e-16
```



QUADRATIC MODEL: EXAMPLE REGRESSION IN R III

 \cdot Another shortcut is just to use the ${\tt poly()}$ command



QUADRATIC MODEL: EXAMPLE REGRESSION IN R III

- Another shortcut is just to use the poly() command
- · Instead of any x variables, just add poly(x,2) where 2 is the highest power desired²

²R gives different coefficient estimates for this version unless you add , raw=TRUE inside the poly() function, to ensure the polynomials are not computed orthogonally.



QUADRATIC MODEL: EXAMPLE REGRESSION IN R III

- Another shortcut is just to use the poly() command
- Instead of any x variables, just add poly(x,2) where 2 is the highest power desired²

²R gives different coefficient estimates for this version unless you add , raw=TRUE inside the poly() function, to ensure the polynomials are not computed orthogonally.





$$\frac{dY}{dX} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$



$$\frac{dY}{dX} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 + 2(-0.02) \text{GDP}$$



$$\frac{d Y}{d X} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 + 2(-0.02) \text{GDP}$$

$$= 1.55 - 0.04 \text{GDP}$$



$$\frac{d Y}{d X} = \hat{\beta}_1 + 2\hat{\beta}_2 X_i$$

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 + 2(-0.02) \text{GDP}$$

$$= 1.55 - 0.04 \text{GDP}$$



- Positive, with diminishing returns
- Effect on Life Expectancy of increasing GDP depends on initial value of GDP!



Life Expectancy_i =50.52+ 1.55 GDP-
$$0.02 GDP^2$$
 (0.30) (0.04) (0.00)

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04(5)$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04(5)$$
$$= 1.55 - 0.20$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04(5)$$
$$= 1.55 - 0.20$$
$$= 1.35$$



• Marginal effect of GDP if GDP= \$5 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04(5)$$
$$= 1.55 - 0.20$$
$$= 1.35$$

• i.e. for every addition \$1 (thousand) in GDP per capita, average life expectancy increases by 1.35 years





$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (25)$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (25)$$
$$= 1.55 - 1.00$$



$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (25)$$
$$= 1.55 - 1.00$$
$$= 0.55$$



• Marginal effect of GDP if GDP= \$25 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (25)$$
$$= 1.55 - 1.00$$
$$= 0.55$$

• i.e. for every addition \$1 (thousand) in GDP per capita, average life expectancy increases by 0.55 years





$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$



• Marginal effect of GDP if GDP= \$50 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (50)$$



• Marginal effect of GDP if GDP= \$50 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (50)$$
$$= 1.55 - 2$$



• Marginal effect of GDP if GDP= \$50 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (50)$$
$$= 1.55 - 2$$
$$= -0.45$$



Marginal effect of GDP if GDP= \$50 (thousand):

$$\frac{d \text{ Life Expectancy}}{d \text{ GDP}} = 1.55 - 0.04 \text{GDP}$$
$$= 1.55 - 0.04 (50)$$
$$= 1.55 - 2$$
$$= -0.45$$

• i.e. for every addition \$1 (thousand) in GDP per capita, average life expectancy *decreases* by 0.45 years



Initial GDP Marginal Effect of +\$1,000 GDP



Initial GDP	Marginal Effect of +\$1,000 GDP
\$5,000	1.35 years




Initial GDP	Marginal Effect of +\$1,000 GDP
\$5,000	1.35 years
\$25,000	0.55 years



Initial GDP	Marginal Effect of +\$1,000 GDP
\$5,000	1.35 years
\$25,000	0.55 years
\$50,000	−0.45 years



QUADRATIC MODEL: EXAMPLE



· For a quadratic model, we can also find the predicted maximum or minimum of $\hat{Y_i}$



- · For a quadratic model, we can also find the predicted maximum or minimum of \hat{Y}_i
- $\boldsymbol{\cdot}$ By calculus, a minimum or maximum occurs where:



- · For a quadratic model, we can also find the predicted maximum or minimum of \hat{Y}_i
- · By calculus, a minimum or maximum occurs where:

$$\frac{dY}{dX} = 0$$



- · For a quadratic model, we can also find the predicted maximum or minimum of \hat{Y}_i
- By calculus, a minimum or maximum occurs where:

$$\frac{dY}{dX} = 0$$
$$\beta_1 + 2\beta_2 X = 0$$



- · For a quadratic model, we can also find the predicted maximum or minimum of \hat{Y}_i
- By calculus, a minimum or maximum occurs where:

$$\frac{dY}{dX} = 0$$
$$\beta_1 + 2\beta_2 X = 0$$
$$2\beta_2 X = -\beta_1$$



- · For a quadratic model, we can also find the predicted **maximum** or **minimum** of \hat{Y}_i
- By calculus, a minimum or maximum occurs where:

$$\frac{dY}{dX} = 0$$

$$\beta_1 + 2\beta_2 X = 0$$

$$2\beta_2 X = -\beta_1$$

$$X^* = -\frac{1}{2} \frac{\beta_1}{\beta_2}$$



Life
$$\widehat{\text{Expectancy}}_i = 50.52 + 1.55 \, GDP - 0.015 \, GDP^2$$
(0.30) (0.04) (0.00)



Life Expectancy_i =50.52+ 1.55
$$GDP$$
-0.015 GDP ² (0.30) (0.04) (0.00)

$$X^* = -\frac{1}{2} \frac{\beta_1}{\beta_2}$$



Life Expectancy_i =50.52+ 1.55
$$GDP$$
-0.015 GDP ² (0.30) (0.04) (0.00)

$$X^* = -\frac{1}{2} \frac{\beta_1}{\beta_2}$$

$$GDP^* = -\frac{1}{2} \frac{(1.55)}{(-0.015)}$$



Life Expectancy_i =50.52+ 1.55
$$GDP$$
-0.015 GDP ² (0.30) (0.04) (0.00)

$$X^* = -\frac{1}{2} \frac{\beta_1}{\beta_2}$$

$$GDP^* = -\frac{1}{2} \frac{(1.55)}{(-0.015)}$$

$$X^* = -\frac{1}{2} (-103.333)$$



Life Expectancy_i =50.52+ 1.55
$$GDP$$
-0.015 GDP ² (0.30) (0.04) (0.00)


$$X^* = -\frac{1}{2} \frac{\beta_1}{\beta_2}$$

$$GDP^* = -\frac{1}{2} \frac{(1.55)}{(-0.015)}$$

$$X^* = -\frac{1}{2} (-103.333)$$

$$X^* = 51.67$$





DETERMINING IF (LARGER) POLYNOMIALS

ARE NECESSARY

• Do we *need* a quadratic model?



- · Do we need a quadratic model?
- We can determine if \hat{eta}_{2} is statistically significant:



- · Do we need a quadratic model?
- We can determine if \hat{eta}_{2} is statistically significant:

•
$$H_0$$
: $\beta_2 = 0$



- · Do we need a quadratic model?
- We can determine if \hat{eta}_{2} is statistically significant:
 - H_0 : $\beta_2 = 0$
 - · H_1 : $\beta_2 \neq 0$



- · Do we need a quadratic model?
- · We can determine if \hat{eta}_{2} is statistically significant:

•
$$H_0$$
: $\beta_2 = 0$

·
$$H_1$$
: $\beta_2 \neq 0$

$$t = \frac{\hat{\beta}_2 - 0}{SE(\hat{\beta}_2)} = \frac{0.015}{0.0005} = 30$$



Life Expectancy_i =50.52+ 1.55 GDP- 0.015
$$GDP^2$$
 (0.30) (0.04) (0.0005)

- · Do we need a quadratic model?
- · We can determine if \hat{eta}_2 is statistically significant:
 - · H_0 : $\beta_2 = 0$
 - · H_1 : $\beta_2 \neq 0$

$$t = \frac{\hat{\beta}_2 - 0}{SE(\hat{\beta}_2)} = \frac{0.015}{0.0005} = 30$$

- \cdot Statistically significant \implies we should keep the quadratic model
 - · If we only ran a linear model, it would be biased!



HIGHER-ORDER POLYNOMIALS: CUBIC REGRESSION

Example

· Should we keep going up in polynomials?



HIGHER-ORDER POLYNOMIALS: CUBIC REGRESSION

Example

· Should we keep going up in polynomials?

Life
$$\widehat{\text{Expectancy}}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{GDP}_i + \hat{\beta}_2 \text{GDP}_i^2 + \hat{\beta}_3 \text{GDP}_i^3$$




HIGHER-ORDER POLYNOMIALS: CUBIC REGRESSION

Example

· Should we keep going up in polynomials?

Life
$$\widehat{\text{Expectancy}}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{GDP}_i + \hat{\beta}_2 \text{GDP}_i^2 + \hat{\beta}_3 \text{GDP}_i^3$$




HIGHER-ORDER POLYNOMIALS

• In general, should have a compelling theoretical reason why data or relationships should "change direction" multiple times




HIGHER-ORDER POLYNOMIALS

- In general, should have a compelling theoretical reason why data or relationships should "change direction" multiple times
- · Or clear data patterns that have multiple "bends"



A SECOND EXAMPLE

ExampleTest Scores: does school district's average income matter?




A SECOND EXAMPLE II

Example

Test Scores: does school district's average income matter?

$$TestScore_i = \hat{\beta_0} + \hat{\beta_1}Income_i$$




A SECOND EXAMPLE III

Example

Test Scores: does school district's average income matter?

TestScore_i =
$$\hat{\beta}_0 + \hat{\beta}_1$$
Income_i + $\hat{\beta}_2$ Income²_i



A SECOND EXAMPLE IV

 $\cdot\,$ Let's manually generate a squared term, ${\tt avgincsq}$



A SECOND EXAMPLE IV

 $\cdot\,$ Let's manually generate a squared term, avgincsq

```
CASchool <- CASchool %>%
  mutate(avgincsq=avginc^2)
```



A SECOND EXAMPLE V

##

```
regsc<-lm(testscr~avginc+avgincsq, data=CASchool)</pre>
summarv(regsc)
##
## Call:
## lm(formula = testscr ~ avginc + avgincsq, data = CASchool)
##
## Residuals:
      Min
##
           10 Median
                              30
                                     Max
## -44.416 -9.048 0.440
                            8.348 31.639
                                                      Test Score = 607.30 + 3.85 Income = 0.04 Income<sup>2</sup>
##
                                                                   (3.05) (0.30)
                                                                                          (0.01)
## Coefficients:
               Estimate Std. Error t value Pr(>|t|)
##
## (Intercept) 607.30174 3.04622 199.362 < 2e-16 ***
## avginc 3.85100 0.30426 12.657 < 2e-16 ***
## avgincsq -0.04231 0.00626 -6.758 4.71e-11 ***
## ---
```


Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1

A SECOND EXAMPLE VI

Example

Test Scores: does school district's average income matter?

$$\textit{TestScore}_i = \hat{\beta}_0 + \hat{\beta}_1 \textit{Income}_i + \hat{\beta}_2 \textit{Income}_i^2 + \hat{\beta}_3 \textit{Income}_i^3$$



A SECOND EXAMPLE VII

 $\cdot\,$ Let's manually generate a cubic term, avginc3:



A SECOND EXAMPLE VII

• Let's manually generate a cubic term, avginc3:

```
CASchool <- CASchool %>%
  mutate(avginc3=avginc^3)
```



A SECOND EXAMPLE VIII

##

```
regcu<-lm(testscr~avginc+avgincsg+avginc3, data=CASchool)</pre>
summary(regcu)
##
## Call:
## lm(formula = testscr ~ avginc + avgincsg + avginc3, data = CASchool)
##
## Residuals:
##
     Min
          10 Median 30
                               Max
## -44.28 -9.21 0.20 8.32 31.16
##
## Coefficients:
##
              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 6.001e+02 5.830e+00 102.937 < 2e-16 ***
## avginc 5.019e+00 8.595e-01 5.839 1.06e-08 ***
## avgincsq -9.581e-02 3.736e-02 -2.564 0.0107 *
## avginc3 6.855e-04 4.720e-04 1.452 0.1471
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
```

1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?



- 1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?
- 2. Plot your data: does a straight line fit well enough?



- 1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?
- 2. Plot your data: does a straight line fit well enough?
- 3. Specify a polynomial function of a higher power (start with 2) and estimate OLS regression



- 1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?
- 2. Plot your data: does a straight line fit well enough?
- 3. Specify a polynomial function of a higher power (start with 2) and estimate OLS regression
- 4. Use *t*-test to determine if higher-power term is significant



- 1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?
- 2. Plot your data: does a straight line fit well enough?
- 3. Specify a polynomial function of a higher power (start with 2) and estimate OLS regression
- 4. Use *t*-test to determine if higher-power term is significant
- 5. Interpret effect of change in *X* on *Y*



- 1. Are there good theoretical reasons for relationships changing (e.g. increasing/decreasing returns)?
- 2. Plot your data: does a straight line fit well enough?
- 3. Specify a polynomial function of a higher power (start with 2) and estimate OLS regression
- 4. Use *t*-test to determine if higher-power term is significant
- 5. Interpret effect of change in *X* on *Y*
- 6. Repeat steps 3-5 as necessary

