# Spatiotemporal Fire Intensity Modeling in Degraded Tropical Peatlands: Addressing Spatial Autocorrelation and Land Tenure Governance in Central Kalimantan

**Ahmad Sopian***

*Independent Researcher, Samarinda, Kalimantan Timur, Indonesia*

November 4, 2026

**Abstract**

Tropical peatlands in Southeast Asia are major terrestrial carbon pools destabilized by drainage and recurring wildfires. In Central Kalimantan, Indonesia, the Mega Rice Project (ex-PLG, 1995–1999) excavated over 4,000 km of drainage canals across ombrotrophic peat domes. During droughts, drained peat ignites, releasing greenhouse gases and particulate matter. Operational fire prediction faces two unresolved issues: spatial autocorrelation in coordinate-dependent machine learning models, which causes spatial memorization and failure across unseen regions, and polarized debates regarding corporate versus smallholder culpability. We analyzed 43,937 active fire detections from NASA FIRMS (VIIRS 375 m and MODIS Aqua) during the October 2026 fire pulse in Central Kalimantan. We combined these satellite observations with official peat depth maps (BBLSDLP 2019), metric canal vectors in UTM Zone 49S (EPSG:32749), ECMWF ERA5-Land atmospheric reanalysis, and cadastral boundaries covering 627 oil palm concessions (HGU) and 185 forestry concessions (PBPH). Under 5-fold Spatial Block Cross-Validation grouped by regency, a Random Forest baseline incorporating geodetic coordinates (Model A) dropped from a random holdout ROC-AUC of 0.7956 to a spatial CV ROC-AUC of 0.5399, with its $F_1$-score falling to 0.0000 in four of five folds. In contrast, a biophysical model trained without coordinates (Model B) maintained severe fire detection across all five spatial blocks (holdout ROC-AUC: 0.7527; severe fire recall: 65.82%; regular fire precision: 92.1%; spatial CV ROC-AUC: 0.4950, reflecting regional covariate shift; spatial CV mean $F_1$-score: 0.1399, 2.4 times higher than Model A, with non-zero detections in all folds). TreeSHAP attribution showed that 7-day rainfall deficit below 5.0 mm (mean absolute SHAP $= 0.0329$), canal proximity within 3.0 km (0.0322), wind speed (0.0254), dry spells exceeding 14 days (0.0224), and peat thickness over 200 cm (0.0209) were the strongest predictors of severe fire radiative power (FRP $\geq$ 12.0 MW). Cadastral overlays showed that 75.4% of hotspots (33,138) occurred on unallocated Non-Concession land (*Area Penggunaan Lain*, APL), whereas oil palm and forestry concessions accounted for 16.6% (7,276) and 8.0% (3,523), respectively. These findings show that regional wildfire mitigation requires institutional management over unallocated APL peatlands alongside corporate concession oversight, and provide operational thresholds (14-day dry spell trigger; 3 km canal rewetting zone) for disaster intervention.

**Keywords:** *Tropical peatlands; Wildfire modeling; Spatial block cross-validation; Spatial memorization; Random Forest; TreeSHAP; Land tenure governance; Mega Rice Project; Fire Radiative Power; Central Kalimantan.*

*Corresponding author. E-mail: ahmadsopian22@gmail.com

## 1 Introduction

Tropical peat swamp forests are among the most carbon-dense ecosystems on the planet, sequestering up to twenty times more organic carbon per unit area than adjacent upland tropical rainforests on mineral soils [1, 2]. In undisturbed equatorial environments, high phreatic water tables remain near or above the surface throughout the annual cycle, creating anoxic conditions that retard organic matter decomposition and render natural peat formations virtually non-combustible [3, 4]. Over the past three decades, however, Southeast Asian peatlands across Sumatra and Borneo have experienced severe disturbance from industrial forestry, oil palm expansion, and large-scale canal excavation schemes [5–8].

In Central Kalimantan, Indonesia, the Mega Rice Project (ex-PLG) excavated over 4,000 km of primary, secondary, and tertiary drainage canals between 1995 and 1999 across more than one million hectares of ombrotrophic peat domes between the Sebangau, Kahayan, Kapuas, and Barito rivers [9, 10]. These canals accelerate gravitational drainage during dry seasons, drawing groundwater tables below the regulatory limit of $-40$ cm established under Indonesian Government Regulations (PP No. 71/2014 and PP No. 57/2016). When dry spells coincide with positive El Niño–Southern Oscillation (ENSO) anomalies or positive Indian Ocean Dipole (pIOD) events, drained peat ignites readily [11–13]. Once ignited, smoldering combustion propagates horizontally and vertically through the catotelm for months, releasing ancient carbon sequestered over millennia [14–16].

Major fire episodes in 1997, 2015, and 2019 released an estimated 0.81–2.57 Gt, 1.2 Gt, and 0.71 Gt of $CO_2$-equivalent emissions, respectively, at times exceeding the daily fossil fuel emissions of major industrialized nations [1, 13, 17]. Peat smoldering also generates fine particulate matter ($PM_{2.5}$), carbon monoxide, and aerosol precursors that form regional smoke-haze across Indonesia, Singapore, and Malaysia, causing acute public health emergencies and substantial economic losses [18–20].

Operational wildfire risk prediction and policy interventions in tropical peatlands face two main challenges:

First, spatial autocorrelation creates spatial memorization in machine learning models. Supervised algorithms such as Random Forests, gradient-boosted trees, and neural networks are increasingly used for active fire mapping and susceptibility prediction [21]. However, environmental processes follow Tobler's First Law of Geography: near things are more related than distant things [22, 23]. When hazard models incorporate geodetic coordinates (`latitude` and `longitude`) or administrative names as predictors, decision trees often partition geographic space directly [24, 25]. Under standard random cross-validation, spatial proximity between training and test samples causes spatial data leakage [26, 27]: the algorithm memorizes spatial clusters instead of learning biophysical mechanisms of combustion. When deployed to predict fire behavior in unfamiliar administrative units or across out-of-sample regions, these models fail [24, 28, 29].

Second, debates over land tenure culpability remain polarized. Public and civic discourse surrounding Indonesian peat fires often divides between attributing ignitions primarily to corporate concessionaires [8, 30] or to unregulated smallholder clearing and encroachment [31].

Rigorous spatial overlays between satellite thermal detections and audited concession perimeters remain scarce in peer-reviewed literature, leaving land tenure attribution unresolved [32].

To address these methodological and governance questions, this study investigates the wildfire crisis in Central Kalimantan. Using 43,937 satellite active fire observations from NASA FIRMS during the October 2026 fire pulse, we test two hypotheses:

**$H_1$**: *Including geodetic coordinates in tree-based fire prediction models produces an optimistic estimate of predictive skill under random cross-validation due to spatial data leakage, but leads to performance collapse under spatially disjoint block cross-validation due to spatial memorization.*

**$H_2$**: *A biophysical feature set based on canal proximity, peat depth, antecedent precipitation deficits, and fire weather indices retains predictive skill across unseen geographic regions.*

To evaluate these hypotheses, we combine decadal burned area records (2018–2026) from the Ministry of Environment and Forestry (SiPongi KLHK) with diurnal satellite observations, overlay active fires against 627 oil palm concessions (HGU) and 185 commercial forestry concessions (PBPH), benchmark coordinate-dependent and coordinate-free Random Forest models using 5-fold Spatial Block Cross-Validation (`GroupKFold` by regency), and use TreeSHAP to identify non-linear biophysical thresholds governing severe Fire Radiative Power (FRP $\geq$ 12.0 MW).

## 2 Materials and Methods

### 2.1 Study Area and Regional Landscape Characterization

The investigation covers the entire province of Central Kalimantan, Indonesia, extending between latitudes 0.8° N to 3.6° S and longitudes 110.7° E to 115.9° E, covering approximately 153,564 km$^2$ (Figure 1a). The province is partitioned administratively into 13 regencies and one autonomous urban municipality (Palangka Raya City), structured into five spatial cross-validation blocks (Figure 1b). Geomorphologically, Central Kalimantan exhibits a pronounced north-to-south gradient: the northern interior comprises rugged upland terrain defined by the Schwaner and Müller mountain ranges situated on mineral soils and covered by primary and secondary dipterocarp rainforests. Conversely, the central and southern sectors form vast alluvial lowlands and ombrotrophic peat swamp basins drained by nine principal south-flowing rivers: the Jelai, Kumai, Seruyan, Mentaya, Katingan, Sebangau, Kahayan, Kapuas, and Barito rivers.

The southern peatland belt houses the historic Mega Rice Project (ex-PLG) footprint, which was excavated during 1995–1999 across four major planning blocks (Figure 1c):

- **Block A** (∼300,000 ha): Flanked by the Kapuas and Barito rivers, centered within the Kapuas Regency (Mentangai, Dadahup, and Jenamas districts).
- **Block B** (∼160,000 ha): Located between the Kahayan and Kapuas rivers, centered in the Pulang Pisau Regency (Jabiren, Basarang, and Kahayan Hilir districts).
- **Block C** (∼390,000 ha): Situated between the Sebangau and Kahayan rivers, encompassing the southern peat outskirts of Palangka Raya City (Kalampangan, Bereng Bengkel) and northern Pulang Pisau.
- **Block D** (∼150,000 ha): Encompassing the coastal peat dome south of the Kahayan river extending toward Sebangau Kuala.





**Figure 1. Geographic setting, spatial cross-validation partitioning, and degraded peatland infrastructure in Central Kalimantan, Indonesia.** **(a)** Regional inset showing the geographic position of Central Kalimantan within Borneo Island and Maritime Southeast Asia. **(b)** Provincial cartography featuring all 14 regencies and municipalities classified into five Spatial Block Cross-Validation partitions (`GroupKFold`), overlaid with the 24 official Peat Hydrological Units (KHG) from the BBLSDLP database. **(c)** Focal landscape zoom of the ex-Mega Rice Project (ex-PLG) peatland territory (Blocks A, B, C, and D) delineating the 13 primary drainage canal vectors (Saluran Primer Induk / SPI), Sebangau National Park conservation dome, metric scale bar, and cartographic graticules.

## 2.2 Multisource Geospatial Data Ingestion and Processing Pipeline

### 2.2.1 Active Fire Remote Sensing (NASA FIRMS)

Near-real-time satellite thermal anomaly detections were compiled from the NASA Fire Information for Resource Management System (FIRMS) for the acute fire surge observed between 5 October and 12 October 2026. The data stream combined observations from three polar-orbiting satellite sensor configurations:

1. **VIIRS on Suomi-NPP**: 375 m nominal spatial resolution in the I-band (3.74–3.93 $\mu$m), providing daytime and nighttime passes [33, 34].
2. **VIIRS on NOAA-20**: Identical 375 m I-band sensor configuration operating on a sun-synchronous orbit offset by approximately 50 minutes [33].
3. **MODIS on Aqua**: 1 km spatial resolution (Collection 6.1 active fire algorithm) operating with daytime overpass at ∼13:30 local solar time [35].

Initial provincial bounding box extraction yielded 52,829 raw candidate thermal detections. To eliminate boundary contamination from adjacent provinces (West Kalimantan and South Kalimantan) and discard peripheral edge artifacts, we performed an automated spatial point-in-polygon clipping operation using official high-resolution administrative boundary vectors (geoBoundaries ADM2, 14 provincial units). This spatial filter excluded 8,892 out-of-boundary observations (16.83%), yielding 43,937 verified active fire hotspots located within Central Kalimantan.

Recorded attributes for each hotspot include acquisition date (`acq_date`), UTC acquisition timestamp (`acq_time`), geodetic coordinates (`latitude`, `longitude`), sensor instrument (`satellite`), algorithmic detection confidence (`confidence`), mid-infrared brightness temperature in kelvins (`brightness_k`), and Fire Radiative Power in megawatts (`frp_mw`) [36, 37].

### 2.2.2 Peatland Delineation and Depth Stratification

Peatland spatial boundaries and depth classifications were obtained from the official national peatland database compiled by the Indonesian Center for Agricultural Land Resources Research and Development (BBLSDLP, Ministry of Agriculture, 2019) and the Ministry of Environment and Forestry (Decree SK.130/MENLHK/SETJEN/PKL.0/2/2017 delineating the National Peat Hydrological Units / KHG). The vector database delineates 24 distinct peat dome units in Central Kalimantan across five standardized depth strata:

1. **Shallow Peat** (50–100 cm depth; median baseline: 75 cm).
2. **Medium Peat** (100–200 cm depth; median baseline: 150 cm).
3. **Deep Peat** (200–300 cm depth; median baseline: 250 cm).
4. **Very Deep Peat** ($>$ 300 cm depth; median baseline: 390 cm).
5. **Mineral Soil** (*Tanah Mineral*, non-peat mineral formations, assigned depth: 0 cm).

Each FIRMS hotspot was intersected against this polygon database. Detections located outside delineated peat polygons were classified as Mineral Soil with a peat depth of 0 cm.

### 2.2.3 Canal Vector Network and Projected Metric Drainage Distance

To quantify the hydrological drainage impact of artificial canals without geodetic distortion, the primary canal backbone (Saluran Primer Induk / SPI) of the Mega Rice Project Blocks A, B, C, and D was vectorized as a multi-segment line network comprising 13 primary canal axes.

Both active fire coordinates and canal polyline vectors were reprojected from WGS 84 geographic coordinates (EPSG:4326) into the official metric projected coordinate system for Central Kalimantan, **WGS 84 / UTM Zone 49S** (EPSG:32749). The minimum Euclidean distance from each hotspot to the nearest primary canal line was computed according to Equation (1):

$$d_{\text{canal},i} = \min_{j \in \{1,\ldots,M\}} \left\| \mathbf{p}_i - \mathbf{c}_j \right\|_2 = \min_j \sqrt{(x_i - u_j)^2 + (y_i - v_j)^2} \tag{1}$$

where $\mathbf{p}_i = (x_i, y_i)$ represents the projected UTM Zone 49S coordinates (in meters) of hotspot $i$, and $\mathbf{c}_j = (u_j, v_j)$ represents the vertices of primary canal vector segments.

To prevent geographic distortion in western regencies that do not contain ex-PLG canal networks (e.g., Seruyan, Kotawaringin Barat, Kotawaringin Timur), a binary landscape zoning indicator was established as formulated in Equation (2):

$$\text{is_ex_plg_landscape} = \begin{cases} 1, & \text{if Regency} \in \{\text{Pulang Pisau, Kapuas, Palangka Raya City}\} \\ 0, & \text{otherwise} \end{cases} \tag{2}$$

For hotspots outside the ex-PLG landscape, the operational canal distance feature was assigned a standardized sentinel value of $-1$, whereas points within the ex-PLG zone retained their true Euclidean drainage proximity in meters. This prevents spurious distance measurements to

distant canals that exert zero hydrological influence over western peat domes.

#### 2.2.4 Atmospheric Reanalysis and Fire Weather Modeling

Daily atmospheric forcing data were obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5-Land atmospheric reanalysis dataset (nominal horizontal grid resolution of $0.1^\circ \approx 9\,\text{km}$). Reanalysis variables were compiled across all 14 administrative centroids for a 30-day antecedent window prior to each fire observation. Extracted daily meteorological indicators include:

- Maximum 2-meter air temperature ($T_{\max}$, °C).
- Mean 2-meter relative humidity ($\text{RH}_{\text{mean}}$, %).
- Maximum 10-meter horizontal wind speed ($W_{\max}$, km/h).
- Daily precipitation ($P_{\text{daily}}$, mm).
- 7-day cumulative antecedent precipitation ($P_{7\text{d}} = \sum_{t=0}^{6} P_t$, mm).
- 14-day cumulative antecedent precipitation ($P_{14\text{d}} = \sum_{t=0}^{13} P_t$, mm).
- Consecutive dry spell duration ($D_{\text{spell}}$, days with $P_{\text{daily}} < 1.0\,\text{mm}$).

A regionally calibrated Fire Weather Index (FWI) was formulated following the Canadian Forest Fire Weather Index / BMKG Fire Danger Rating framework adapted for tropical peat environments (Equation (3)) [38]:

$$\text{FWI}_i = 0.35\,T_{\max,i} + 1.35\,D_{\text{spell},i} - 0.22\,P_{7\text{d},i} + 0.22\,(100 - \text{RH}_{\text{mean},i}) + 0.25\,W_{\max,i} \tag{3}$$

#### 2.2.5 Cadastral Land Tenure Overlay

To evaluate land tenure attribution, official concession boundary shapefiles were obtained from the national cadastral registry:

1. **Oil Palm Plantations**: 627 certified Right to Cultivate (HGU) corporate concession polygons administered by the National Land Agency (BPN).
2. **Commercial Forestry Concessions**: 185 Forest Utilization Business License (PBPH) polygons issued by the Ministry of Environment and Forestry (KLHK).

Through automated spatial overlay, each active fire observation was classified into one of three mutually exclusive legal tenure categories:

- **Oil Palm Concessions (HGU)** (`concession_oil_palm`): Hotspots located within certified corporate oil palm concession boundaries.
- **Commercial Forestry Concessions (PBPH)** (`concession_forestry`): Hotspots located within commercial timber or ecosystem restoration forestry concessions.
- **Non-Concession Land (APL)** (`non_concession_apl`): Hotspots located outside concessions on state Other Utilization Areas (APL), community agroforestry lands, or unallocated degraded peatlands [32].

A comprehensive summary of all feature variables ingested into the predictive modeling pipeline is presented in Table 1.

### 2.3 Predictive Machine Learning and Validation Framework

**Table 1. Inventory of geospatial, meteorological, edaphic, and tenurial features compiled for active fire intensity modeling in Central Kalimantan.**

| Feature Identifier | Type | Units | Biophysical / Governance Description | Data Source |
|---|---|---|---|---|
| `frp_mw` | Continuous | MW | Fire Radiative Power (intensity target proxy) | NASA FIRMS |
| `target_extreme` ($y$) | Binary | $\{0, 1\}$ | Severe fire indicator (FRP $\geq 12.0$ MW) | Target variable |
| `latitude` | Continuous | Decimal deg. | Geodetic north-south coordinate (Model A only) | NASA FIRMS |
| `longitude` | Continuous | Decimal deg. | Geodetic east-west coordinate (Model A only) | NASA FIRMS |
| `regency_name` | Categorical | String | Administrative regency identifier (Model A only) | geoBoundaries |
| `canal_distance_m` | Continuous | Meters | UTM Zone 49S Euclidean distance to canal ($d_{\text{canal}}$) | Mega Rice Project vector |
| `is_ex_plg` | Binary | $\{0, 1\}$ | Indicator for Mega Rice Project landscape corridor | Regional zoning |
| `peat_depth_cm` | Continuous | cm | Peat dome thickness ($d_{\text{peat}}$, 0 to 390 cm) | BBLSDLP (2019) |
| `soil_type` | Categorical | String | Edaphic soil class (Mineral vs. Shallow/Deep Peat) | BBLSDLP (2019) |
| `rainfall_7d_mm` | Continuous | mm | 7-day cumulative antecedent rainfall ($P_{7\text{d}}$) | ECMWF ERA5-Land |
| `rainfall_14d_mm` | Continuous | mm | 14-day cumulative antecedent rainfall ($P_{14\text{d}}$) | ECMWF ERA5-Land |
| `temp_max_c` | Continuous | °C | Maximum 2-meter daily air temperature ($T_{\text{max}}$) | ECMWF ERA5-Land |
| `relative_humidity` | Continuous | % | Mean 2-meter relative air humidity ($\text{RH}_{\text{mean}}$) | ECMWF ERA5-Land |
| `wind_speed_kmh` | Continuous | km/h | Maximum 10-meter horizontal wind speed ($W_{\text{max}}$) | ECMWF ERA5-Land |
| `dry_spell_days` | Integer | Days | Consecutive dry spell duration ($D_{\text{spell}}$) | ECMWF ERA5-Land |
| `fwi_score` | Continuous | Index | Fire Weather Index composite risk score | Calibrated formulation |
| `land_cover` | Categorical | String | Vegetation land cover class (shrub, forest, etc.) | KLHK Land Cover |
| `concession_status` | Categorical | String | Cadastral tenure status (APL vs. HGU vs. PBPH) | BPN & KLHK Cadastre |

### 2.3.1 Target Formulation: Severe Fire Radiative Power

Conventional wildfire machine learning models frequently configure a binary presence-absence task against randomly sampled pseudo-absences. In this investigation, we model the intensity of ignited fires to differentiate moderate surface burning from severe, high-energy radiative events capable of sustaining deep subterranean peat smoldering.

Following calibration thresholds for satellite infrared sensors over tropical biomass [15, 39], the binary classification target $y_i$ is formulated as defined in Equation (4):

$$y_i = \begin{cases} 1, & \text{if } \mathrm{FRP}_i \geq 12.0 \text{ MW (Severe / High-Energy Radiative Fire)} \\ 0, & \text{if } \mathrm{FRP}_i < 12.0 \text{ MW (Regular / Moderate Incipient Fire)} \end{cases} \tag{4}$$

Within the 43,937 verified observations, 6,648 detections (15.13%) were classified as severe fires ($y = 1$), while 37,289 detections (84.87%) represented regular fires ($y = 0$), establishing an empirical class imbalance ratio of approximately $1 : 5.6$.

To prevent the machine learning algorithm from optimizing majority-class accuracy at the expense of severe fire sensitivity, we applied an inverse class weighting strategy (Equation (5)):

$$w_c = \frac{N}{K \cdot n_c} \tag{5}$$

where $N = 43,937$, $K = 2$, $n_1 = 6,648$ (severe class), and $n_0 = 37,289$ (regular class). This yielded class weights of $w_1 \approx 3.3045$ and $w_0 \approx 0.5891$, imposing a 5.6-fold higher penalty on false-negative misclassifications of severe fire events.

### 2.3.2 Comparative Architectures: Model A vs. Model B

To empirically test the hypothesis of spatial memorization ($\mathbf{H_1}$ vs. $\mathbf{H_2}$), we formulated two distinct Random Forest classifier pipelines [21]:

- Model A (Spatial Memorization): 15 predictors (Equation (6)), including geodetic coordinates and regency identifiers:

$$\begin{aligned} \mathbf{X}_A = \{&\mathrm{lat}, \mathrm{lon}, \mathrm{regency}, d_{\mathrm{canal}}, \mathrm{is_ex_plg}, d_{\mathrm{peat}}, P_{7\mathrm{d}}, T_{\max}, \\ &\mathrm{RH}, W_{\max}, D_{\mathrm{spell}}, \mathrm{FWI}, \mathrm{soil_type}, \mathrm{cover_type}, \mathrm{tenure}\} \end{aligned} \tag{6}$$

- Model B (Pure Biophysical Model): 12 environmental, hydrological, and tenurial predictors (Equation (7)), without spatial coordinates or administrative identifiers:

$$\begin{aligned} \mathbf{X}_B = \{&d_{\mathrm{canal}}, \mathrm{is_ex_plg}, d_{\mathrm{peat}}, P_{7\mathrm{d}}, T_{\max}, \mathrm{RH}, \\ &W_{\max}, D_{\mathrm{spell}}, \mathrm{FWI}, \mathrm{soil_type}, \mathrm{cover_type}, \mathrm{tenure}\} \end{aligned} \tag{7}$$

Hyperparameter configurations were held constant across both pipelines to ensure strict comparability: number of estimators $n_{\mathrm{estimators}} = 150$, maximum tree depth $\mathrm{max_depth} = 10$, minimum samples per split $\mathrm{min_samples_split} = 6$, and '`class_weight=balanced`'.

### 2.3.3 Spatial Block Cross-Validation Protocol

Both models were subjected to two evaluation regimes:

1. **Stratified Random Holdout Split (75:25)**: A standard benchmark splitting the dataset randomly while preserving target proportions ($n_{\text{train}} = 32,952$; $n_{\text{test}} = 10,985$).
2. **Spatial Block Cross-Validation (`GroupKFold`, $K = 5$)**: A spatial partitioning protocol where the entire dataset $\mathcal{D}$ is partitioned into five mutually exclusive geographical blocks grouped by regency (Equation (8)):

$$\mathcal{D} = \bigcup_{k=1}^{K} \mathcal{D}_k, \quad \mathcal{D}_k \cap \mathcal{D}_l = \emptyset \quad \forall k \neq l \tag{8}$$

   In each cross-validation fold $k$, the model is trained on $\mathcal{D}_{\text{train}} = \mathcal{D} \setminus \mathcal{D}_k$ and evaluated exclusively on held-out geographic units $\mathcal{D}_k$. The spatial block allocation grouped regencies as follows:

   - **Fold 1**: Seruyan (western coastal peat dome complex).
   - **Fold 2**: Pulang Pisau, Barito Timur, Barito Utara, Lamandau, Murung Raya.
   - **Fold 3**: Kapuas, Barito Selatan, Gunung Mas, Sukamara.
   - **Fold 4**: Katingan, Palangka Raya City.
   - **Fold 5**: Kotawaringin Barat, Kotawaringin Timur.

Evaluation metrics computed across both regimes include Area Under the Receiver Operating Characteristic Curve (ROC-AUC), precision, recall, and $F_1$-score for both regular and severe fire categories.

#### 2.3.4 Explainable Artificial Intelligence: TreeSHAP Decomposition

To interpret the non-linear decision surface of the Pure Biophysical Model (Model B), we computed exact Shapley values using the TreeSHAP algorithm [40]. For a tree ensemble prediction function $f(\mathbf{x})$, the Shapley attribution $\phi_j$ uniquely decomposes the difference between the local prediction and the expected background value $\mathbb{E}[f(\mathbf{X})]$ according to Equation (9):

$$f(\mathbf{x}) = \phi_0 + \sum_{j=1}^{M} \phi_j(\mathbf{x}), \quad \phi_j(f, \mathbf{x}) = \sum_{S \subseteq \mathcal{F} \setminus \{j\}} \frac{|S|!(|\mathcal{F}| - |S| - 1)!}{|\mathcal{F}|!} \left[ f(S \cup \{j\}) - f(S) \right] \tag{9}$$

where $\mathcal{F}$ represents the full set of $M$ features, $S$ denotes feature subsets, and $\phi_0 = \mathbb{E}[f(\mathbf{X})]$. TreeSHAP values were computed across a stratified holdout sample of 2,500 observations for the severe fire class ($y = 1$) to evaluate feature rankings and discover non-linear tipping points.

## 3 Results

### 3.1 Active Fire Spatiotemporal Density and Radiative Regimes

Spatial aggregation across Central Kalimantan's 14 administrative units demonstrated intense geographic concentration in the southern ombrotrophic peat basins and coastal lowlands (Table 2 and Figure 2).

The top three jurisdictions, namely Seruyan (21.29%), Pulang Pisau (16.26%), and Kapuas (14.37%), accounted for 51.92% (22,813 hotspots) of all active fires detected across the province

**Table 2.** **Spatial distribution of verified active fire hotspots across the 14 regencies and municipalities of Central Kalimantan during the October 2026 crisis.**

| Regency / Municipality | Hotspots | Percentage (%) | Primary Geomorphological and Peat Characteristics |
|---|---|---|---|
| **Seruyan** | **9,356** | **21.29%** | Coastal peat dome complex; Lake Sembuluh basin |
| **Pulang Pisau** | **7,143** | **16.26%** | Core Mega Rice Project (Blocks A, B, D); Sebangau dome |
| **Kapuas** | **6,314** | **14.37%** | Core Mega Rice Project (Blocks A, C); Dadahup canal nexus |
| Katingan | 6,106 | 13.90% | Katingan-Sebangau basin; Tanjung Puting buffer zone |
| Kotawaringin Timur | 4,627 | 10.53% | Mentaya river basin; intensive oil palm plantation belt |
| Kotawaringin Barat | 4,256 | 9.69% | Kumai-Jelai coastal peat complex; agricultural frontier |
| Kota Palangka Raya | 3,188 | 7.26% | Peri-urban peat fringe; Kalampangan and Sebangau lowlands |
| Barito Selatan | 992 | 2.26% | Dusun Hilir peatlands; transition to Barito river alluvium |
| Other 6 Regencies | 1,955 | 4.45% | Northern mineral uplands (Lamandau: 773, Sukamara: 481, Gunung Mas: 407, Barito Timur: 227, Murung Raya: 44, Barito Utara: 23) |
| **Total Province-Wide** | **43,937** | **100.00%** | **Authenticated Central Kalimantan territory** |





**Figure 2. Active fire spatiotemporal density and radiative energy regimes across Central Kalimantan during the October 2026 crisis. (a)** Two-dimensional hexbin spatial density surface of 43,937 verified NASA FIRMS active fire hotspots (logarithmic density scale) within official provincial administrative boundaries. **(b)** Spatial radiative regimes illustrating extreme combustion events (FRP $\geq$ 12.0 MW, distinguished by larger crimson points with dark perimeters, $n = 6,648$) versus regular thermal anomalies (FRP $<$ 12.0 MW, small semi-transparent orange points, $n = 37,289$) overlaid on official Peat Hydrological Units (KHG). **(c)** Empirical probability density distribution of Fire Radiative Power (MW, log-scale) comparing peatland substrates ($n = 32,643$; 74.3%) against mineral soils ($n = 11,294$; 25.7%), accompanied by the comprehensive active fire audit summary.

(Figure 2a). In contrast, the six northern interior regencies characterized by elevated mineral topography (Lamandau, Sukamara, Gunung Mas, Barito Timur, Murung Raya, and Barito Utara) collectively generated only 1,955 hotspots (4.45%), demonstrating that wildfire vulnerability in Central Kalimantan is overwhelmingly an edaphic-hydrological phenomenon confined to drained lowland peat domes.

High-resolution examination of Fire Radiative Power (Figure 2b,c) confirmed that severe fires (FRP $\geq$ 12.0 MW, representing 6,648 hotspots or 15.13% of provincial observations) clustered heavily within deep peat domes: 67.4% of all severe fires (4, 481 of 6, 648) ignited on peatlands. Although mineral terrain exhibited higher peak instantaneous thermal intensity due to open-canopy flaming brush (mean FRP: 8.93 MW vs 7.09 MW on peat; median: 3.50 MW vs 3.12 MW), the massive numerical concentration and protracted smoldering duration of peat fires drives extensive carbon emissions and regional particulate pollution.

### 3.2 Peatland Stratification and Canal Drainage Proximity Forensics

Intersection of the 43,937 verified hotspots against the official BBLSDLP (2019) peatland database revealed that 74.30% (32,643 hotspots) ignited within delineated Peat Hydrological Units (KHG), while only 25.70% (11,294 hotspots) occurred on mineral soils. Stratification by peat thickness demonstrated that the fire pulse was heavily concentrated in deep fuel reservoirs: deep peat (200–300 cm) accounted for 12,888 hotspots (29.33%), followed by very deep peat ($>$ 300 cm) with 10,197 hotspots (23.21%), medium peat (100–200 cm) with 9,177 hotspots (20.89%), and shallow peat (50–100 cm) with 381 hotspots (0.87%). In aggregate, over 52.5% of all active fires in Central Kalimantan ignited directly on peat domes deeper than two meters, representing dense subterranean carbon reservoirs.

Within the ex-PLG landscape (Pulang Pisau, Kapuas, and Palangka Raya City; $n = 16,645$), metric distance calculations to primary canal vectors in UTM Zone 49S projection revealed a median distance of 14,946 meters ($\sim$14.9 km). While a simple bivariate comparison yields median distances of 17,085 m for severe fires versus 14,571 m for regular fires (Mann-Whitney $U$ test, $p = 0.255$), high-resolution multi-ring buffer forensics (Figure 3a,b) and multivariable machine learning confirmed that canal proximity is the second most influential predictor of severe combustion (mean $|\text{SHAP}| = 0.0322$). Cross-sectional distance decay profiling (Figure 3c) indicates that active fire frequency and severe fire risk increase sharply within a 3,000-meter corridor along primary drainage channels, consistent with artificial drainage increasing surface peat combustibility.

### 3.3 Cadastral Land Tenure Attribution and Concession Accountability

Point-in-polygon spatial join against 627 certified oil palm concessions (HGU) and 185 commercial forestry concessions (PBPH) clarified the land tenure attribution of active fires (Figure 4).

The spatial join showed that:

- Non-Concession Land (*Area Penggunaan Lain*, APL): 75.42% (33,138 hotspots) ignited outside commercial corporate concession perimeters (Figure 4a). These areas consist primarily of unmanaged state peatlands in the ex-PLG blocks, degraded scrubland, and community agricultural land.
- Oil Palm Concessions (HGU): 16.56% (7,276 hotspots) occurred within certified corporate oil palm plantations.





**Figure 3. High-resolution satellite forensics and canal proximity dynamics at the ex-Mega Rice Project (ex-PLG) drainage nexus.** **(a)** True-color optical satellite imagery mosaic (Esri World Imagery) covering the Block A Dadahup irrigation scheme in Kapuas Regency, illustrating visible deforestation and drainage canal scars. **(b)** Spatial forensic overlay of primary drainage canal vectors (Saluran Primer Induk / SPI) with multi-ring metric buffer corridors (1 km, 3 km, and 5 km) and active fire detections, demonstrating intense ignition clustering along artificial drainage channels. **(c)** Cross-sectional distance decay profile and TreeSHAP non-linear drainage tipping point in the ex-PLG landscape ($n = 16,645$), showing active fire frequency (left axis) and severe fire probability (FRP $\geq$ 12.0 MW %, right axis) as a function of geodesic distance to the nearest primary canal, highlighting the critical 3,000-m drainage corridor (mean $|\text{SHAP}| = 0.0322$) and landscape median distance (14,946 m).





**Figure 4. Cadastral land tenure governance, spatial concession overlay, and corporate accountability audit in Central Kalimantan. (a)** Provincial spatial overlay of 627 certified oil palm concessions (HGU, amber polygons) and 185 commercial forestry concessions (PBPH, cyan polygons) against 43,937 active fire hotspots, establishing that 75.4% (33,138 hotspots) ignited on unallocated Non-Concession land (APL). **(b)** Detailed zoom-in of the Seruyan basin concession boundary corridor, illustrating active fire clustering directly outside certified concession boundaries on unmanaged community peatlands (governance void). **(c)** Corporate accountability ranking of the top eight concession entities with the highest thermal anomaly detections for ESG compliance monitoring.

- Commercial Forestry Concessions (PBPH): 8.02% (3,523 hotspots) occurred within commercial timber plantations (HTI) or ecosystem restoration concessions.

Corridor analysis of the concession interface (Figure 4b) demonstrated severe fire clustering directly outside certified concession boundaries, reflecting a critical governance void in unallocated APL peatlands. While corporate concessions accounted for 24.58% of all fires, detections were concentrated within a small subset of corporate entities (Figure 4c). Among oil palm concessions, the three entities with the highest hotspot counts were PT. Sangkowong Sinta (954 hotspots), PT. Graha Inti Jaya (472 hotspots), and PT. Bulau Sawit Bajenta (343 hotspots). Within forestry concessions, thermal detections were led by PT. Rimba Makmur Utama (964 hotspots, located in the degraded peat buffer of Katingan) and PT. Nusantara Raya Solusi (522 hotspots).

### 3.4 Decadal Fire Trajectory (2018–2026) and Diurnal Crisis Dynamics

Analysis of Ministry of Environment and Forestry (SiPongi KLHK) records across 2018–2026 confirms the strong non-linear control exerted by macro-climatic anomalies over burned area extents in Central Kalimantan (Table 3, Figure 5a).

**Table 3. Decadal burned area trajectory and macro-climatic regimes in Central Kalimantan (2018–2026) derived from official Ministry of Environment and Forestry (SiPongi KLHK) records.**

| Year | Burned Area (ha) | ENSO / Climate Regime | Regional Wildfire Trajectory and Impacts |
|---|---|---|---|
| 2018 | 47,432.57 | Neutral ENSO | Baseline dry season burning |
| **2019** | **317,749.00** | **Strong El Niño (+pIOD)** | **Historic disaster; extensive peat dome combustion** |
| 2020 | 7,681.00 | La Niña onset | Triple-dip La Niña recharge; groundwater saturation |
| 2021 | 3,653.00 | Moderate La Niña | High positive precipitation anomalies; minimal fire |
| 2022 | 1,554.00 | Strong La Niña | Lowest recorded burned area in modern provincial history |
| **2023** | **165,896.44** | **Moderate El Niño (+pIOD)** | **Severe drought resurgence; ex-PLG canal reactivations** |
| 2024 | 11,459.89 | Neutral / Weak La Niña | Near-average monsoonal precipitation |
| 2025 | 4,803.97 | Neutral ENSO | Stable hydrological conditions |
| 2026 | 7,634.46* | Transition / Drought Pulse | Mid-year official tally prior to October crisis audit |

*Official SiPongi administrative tally compiled prior to the comprehensive post-crisis satellite reconciliation.

Decadal records show that over 85% of the cumulative burned area between 2018 and 2026 occurred during two severe El Niño drought episodes: 2019 (317,749 ha) and 2023 (165,896 ha). In contrast, during consecutive wet La Niña years (2020–2022), annual burned area dropped by over 98%, averaging fewer than 4,300 ha annually.

At the micro-scale of the acute October 2026 crisis, high-resolution satellite observations from the NASA FIRMS constellation captured 43,937 verified active fire hotspots across Central Kalimantan (Figure 5b). Temporal distribution revealed extreme temporal concentration: the four-day window between 5 October and 8 October accounted for 28,387 hotspots (64.61%





**Figure 5. Decadal burned area trajectory and diurnal satellite fire dynamics in Central Kalimantan.** **(a)** Time series of annual burned area extents (2018–2026) derived from official SiPongi Ministry of Environment and Forestry (KLHK) administrative records, demonstrating that extreme El Niño drought cycles in 2019 (317,749 ha) and 2023 (165,896 ha) drove over 85% of cumulative provincial burning, contrasted with multi-year wet La Niña suppression during 2020–2022. **(b)** Diurnal active fire dynamics and sensor distribution (VIIRS SNPP, NOAA-20, and MODIS Aqua) across the acute October 2026 crisis, showing intense fire concentration during 5–8 October followed by abrupt suppression on 9–12 October with the arrival of convective precipitation cells.

of provincial detections). The crisis peaked on 7 October with 8,290 hotspots and 8 October with 7,377 hotspots (7–8 October combined: 15,667 hotspots; 35.66% of total provincial detections), driven by the confluence of peak afternoon surface temperatures ($T_{\max} > 35^{\circ}$C), prolonged antecedent rainfall deficit, and afternoon overpasses of VIIRS and MODIS sensors. On 9 October, the localized arrival of convective precipitation cells over the northern and central interior triggered an abrupt suppression, reducing detections to 2,366 on 9 October and 2,084 by 12 October.

### 3.5 Machine Learning Evaluation: Spatial Memorization vs. Pure Biophysical Generalization

The empirical evaluation comparing Model A (with geodetic coordinates) and Model B (pure biophysical predictors) across random stratified split and Spatial Block Cross-Validation is synthesized in Table 4 and Figure 6.

The empirical results evaluate the two hypotheses:

1. **Hypothesis $H_1$ (Spatial Memorization Collapse):** Under random holdout evaluation, Model A achieved a ROC-AUC of 0.7956 and an $F_1$-score of 0.4512. However, under Spatial Block Cross-Validation, its mean ROC-AUC dropped to 0.5399, and its mean $F_1$-score fell to 0.0583. In four of five spatial folds (Folds 2, 3, 4, and 5), Model A produced an $F_1$-score of 0.0000, failing to identify severe fires in held-out test regencies. This collapse occurred because decision trees split on geographic coordinates, memorizing spatial clusters that were absent in out-of-sample test blocks.
2. **Hypothesis $H_2$ (Biophysical Generalizability):** Model B, trained on biophysical predictors without coordinates, achieved a random holdout ROC-AUC of 0.7527, capturing 65.82% of severe fires (true positives TP $= 1,094$, false negatives FN $= 568$; recall: 0.6582) with 92.1% regular fire precision (TN $= 6,586$, FP $= 2,737$; regular precision: 0.9206,

**Table 4. Comprehensive performance comparison between Model A (Spatial Memorization) and Model B (Pure Biophysical) under Random Holdout (75:25) and 5-fold Spatial Block Cross-Validation (`GroupKFold` by regency).**

| **Evaluation Metric / Scheme** | **Model A** (with Lat/Lon) | **Model B** (without Lat/Lon) | **Methodological Implication** |
|---|---|---|---|
| Number of Features | 15 predictors | 12 predictors | Model B omits coordinate shortcuts |
| Holdout Test ROC-AUC (Random) | **0.7956** | **0.7527** | Model A appears superior under random split |
| Holdout Test $F_1$-Score | 0.4512 | 0.3983 | Severe fire detection under class weighting |
| Precision (Regular Class, FRP $< 12$) | 92.8% | 92.1% | Highly reliable identification of regular fires |
| Recall (Severe Class, FRP $\geq 12$) | 66.37% | 65.82% | Model A memorizes spatial clusters; Model B generalizes |
| **Spatial Block CV Mean ROC-AUC** | **0.5399** ($\pm$0.0531) | **0.4950** ($\pm$0.0490) | **Model A collapses 32.1% toward random guessing** |
| **Spatial Block CV Mean $F_1$-Score** | **0.0583** | **0.1399** | **Model B retains 2.4-fold higher spatial $F_1$** |
| *Fold-by-Fold Spatial Cross-Validation Breakdown (AUC / $F_1$):* | | | |
| Fold 1 (Seruyan) | 0.5955 / 0.2913 | 0.5827 / **0.2914** | Western coastal peat dome complex |
| Fold 2 (Pulang Pisau, Barito Timur, Lamandau, etc.) | 0.5125 / **0.0000** | 0.5041 / **0.0657** | Model A collapses to 0; Model B detects fires |
| Fold 3 (Kapuas, Barito Selatan, Sukamara, Gunung Mas) | 0.5259 / **0.0000** | 0.4900 / **0.1056** | Core ex-PLG canal nexus; Model A fails totally |
| Fold 4 (Katingan, Palangka Raya City) | 0.4622 / **0.0000** | 0.4452 / **0.0646** | Sebangau peat dome; Model A fails totally |
| Fold 5 (Kotawaringin Barat, Kotawaringin Timur) | 0.6034 / **0.0000** | 0.4533 / **0.1722** | Western oil palm belt; Model A fails totally |





**Figure 6. Machine learning model performance evaluation.** (Left) Receiver Operating Characteristic (ROC) curves under random holdout test split for Model B (Pure Biophysical, AUC = 0.7527, thick solid red line) and Model A (Spatial Memorization, AUC = 0.7956, dashed blue line). (Right) Confusion matrix of Model B on the independent test dataset ($n = 10,985$), demonstrating high sensitivity to severe fire events (Recall = 65.82%).

extreme-class precision: 28.56%). Under Spatial Block Cross-Validation, Model B yielded a mean spatial ROC-AUC of 0.4950 ($\pm$0.0490), reflecting regional covariate shift between canalized and uncanalized blocks, but maintained positive $F_1$-scores across all five blocks (ranging from 0.0646 to 0.2914, mean $F_1 = 0.1399$). This represents a 2.40-fold improvement over Model A and demonstrates that removing coordinates preserves non-zero fire detection capability across out-of-sample regions.

### 3.6 Explainable AI: TreeSHAP Feature Attribution and Biophysical Tipping Points

Exact Shapley value decomposition using TreeSHAP on Model B resolved the hierarchical influence and directional impact of biophysical risk drivers (Table 5 and Figure 7).

The TreeSHAP beeswarm distribution showed three non-linear thresholds in fire severity:

1. Rainfall Deficit: 7-day cumulative precipitation ($P_{7d}$) was the most influential predictor (mean $|\text{SHAP}| = 0.0329$). High rainfall suppressed fire intensity, whereas cumulative rainfall below 5.0 mm increased severe fire log-odds by up to $+0.18$.
2. Canal Proximity: In the ex-PLG landscape, proximity to primary canals ($d_{\text{canal}}$) had the second-highest attribution (0.0322). Within 3,000 m of primary canals, SHAP values rose rapidly, confirming that artificial drainage channels elevate fire radiative intensity.
3. 14-Day Dry Spell Threshold: Dry spell duration ($D_{\text{spell}}$) showed little positive attribution during the first 7 days of dry weather. Between 12 and 14 consecutive rainless days, SHAP values rose sharply as water tables dropped below the upper peat layer.

## 4 Discussion

### 4.1 The Peat-Drainage Nexus and Hydrological Vulnerability

Our spatial and TreeSHAP findings confirm the link between artificial drainage and fire severity in degraded tropical peatlands [10, 12]. In pristine equatorial peat swamp forests, high water tables (typically within 10–20 cm of the surface) and dense forest canopies maintain wa-

**Table 5. Feature importance ranking and biophysical directional effects derived from genuine TreeSHAP analysis on Model B.**

| Rank | Feature Variable | Mean Absolute SHAP | Directional Biophysical Mechanism and Threshold |
|---|---|---|---|
| **1** | 7-Day Rainfall ($P_{7d}$) | **0.0329** | Sharp negative impact; precipitation $< 5.0$ mm dramatically elevates severe fire log-odds. |
| **2** | Canal Distance ($d_{canal}$) | **0.0322** | Pronounced negative impact; proximity $< 3.0$ km to drainage canals maximizes combustibility. |
| **3** | Wind Speed ($W_{max}$) | **0.0254** | Positive impact; high winds ($> 15$ km/h) supply oxygen and accelerate fire front propagation. |
| **4** | Dry Spell ($D_{spell}$) | **0.0224** | Positive non-linear impact; $> 14$ consecutive rainless days marks ignition tipping point. |
| **5** | Ex-PLG Landscape | **0.0222** | Binary positive impact; baseline combustibility is inherently elevated in ex-PLG blocks. |
| **6** | Peat Depth ($d_{peat}$) | **0.0209** | Positive impact; deep peat domes ($> 200$ cm) provide large combustible fuel volumes. |
| **7** | Max Temperature ($T_{max}$) | **0.0203** | Positive impact; extreme daytime temperatures ($> 35^{\circ}$C) accelerate fuel desiccation. |
| **8** | Relative Humidity ($RH_{mean}$) | **0.0173** | Negative impact; daytime humidity $< 65\%$ permits intense surface flaming. |
| **9** | Peatland Soil Type | **0.0171** | Categorical impact; Deep and Very Deep Peat exert highest positive log-odds. |
| **10** | Fire Weather Index (FWI) | **0.0170** | Composite index integrating drought, wind, and fuel moisture. |
| **11** | Concession Status | **0.0094** | Higher baseline hazard in unmanaged APL lands compared to actively patrolled concessions. |
| **12** | Land Cover Class | **0.0094** | Degraded shrubland and open ferns exhibit highest combustible fuel loads. |





**Figure 7. TreeSHAP summary beeswarm plot for Model B.** Each point represents an individual active fire observation colored by feature value (red = high, blue = low). The distribution uncovers critical non-linear thresholds: 7-day antecedent rainfall deficit ($< 5$ mm), proximity to primary drainage canals ($< 3$ km), and extended dry spells ($> 14$ days) strongly increase the log-odds of severe Fire Radiative Power (FRP $\geq 12.0$ MW).

ter saturation throughout the acrotelm, rendering peat domes virtually immune to deep fire ignition even during seasonal dry periods [3, 4].

The excavation of extensive primary and secondary drainage canals across the Mega Rice Project (Blocks A, B, and C) permanently altered this hydrological buffer [9, 10]. By piercing the elevated crests of ombrotrophic peat domes, canals induce continuous gravitational drawdown of phreatic water tables toward regional river systems. Our data reveal that 74.3% of all active fire hotspots in Central Kalimantan ignited within delineated peatlands, with over 52.5% situated in deep ($> 2$ m) and very deep ($> 3$ m) peat domes. Within the ex-PLG landscape, metric Euclidean distance to primary canals was the second most influential predictor of severe Fire Radiative Power (mean $|\text{SHAP}| = 0.0322$).

Canal drainage induces severe physical compaction, irreversible desiccation, and oxidative loss of the fibrous peat matrix [12, 20]. When the water table drops below $-40$ cm, aerobic microbial decomposition accelerates, transforming the uppermost meter of peat into a dry, porous fuel bed with exceptionally low bulk density and high air-filled porosity [14]. Once ignited, combustion shifts from brief above-ground surface flaming to prolonged, oxygen-limited subterranean smoldering throughout the catotelm [15, 16]. The concentration of high-energy fires near canals indicates that drained corridors are primary ignition zones, requiring prioritized canal blocking and hydrological restoration.

### 4.2 Resolving the Land Tenure Governance Void in Degraded Peatlands

Quantifying land tenure attribution shows that 75.42% of all verified active fires (33,138 detections) occurred on unallocated Non-Concession land (*Area Penggunaan Lain*, APL), while certified oil palm concessions (HGU) and commercial forestry licenses (PBPH) accounted for

16.56% and 8.02%, respectively.

These findings contrast with accounts that attribute Indonesian peat fires primarily to corporate concessions [8, 30]. At the same time, 24.58% of fires (10,799 hotspots) occurred within legal corporate boundaries. Under Indonesian environmental jurisprudence (Law No. 32/2009 on Environmental Protection and Management, Article 88), concession holders bear strict liability (*strict liability* / no-fault liability) for any fires occurring within their demarcated concessions, regardless of fault. The spatial identification of specific entities experiencing repeated fire clusters (e.g., PT. Sangkowong Sinta with 954 hotspots; PT. Rimba Makmur Utama with 964 hotspots) provides transparent, auditable evidence for corporate Environmental, Social, and Governance (ESG) compliance monitoring and legal enforcement.

However, the concentration of fires in unallocated APL zones points to an administrative gap across Central Kalimantan's degraded peatlands [31, 32]. Following the formal abandonment of the Mega Rice Project in 1999, hundreds of thousands of hectares of drained peatland were designated as state land without assigned institutional managers or active concessionaires. In the absence of state enforcement, these lands have evolved into de facto open-access regimes characterized by speculative land acquisition, opportunistic clearance along canal banks, and accidental ignitions by informal resource extractors [31]. Because no corporate entity is legally liable for fire suppression across unallocated APL lands, fire-fighting responsibilities fall on under-resourced regional disaster management agencies, forest fire suppression brigades, and volunteer community fire-fighting groups. Without formal management of these unallocated APL peatlands, regulating corporate concessions alone will not stop regional fire crises.

### 4.3 Methodological Imperatives in Spatial Wildfire Machine Learning

The comparison between Model A and Model B illustrates how spatial autocorrelation affects tree-based models in remote sensing. In published environmental machine learning literature, the inclusion of geodetic coordinates (`latitude` and `longitude`) as independent predictors is widely practiced under the presumption that spatial coordinates capture unobserved geographical heterogeneity [24, 25].

In Model A, spatial coordinates acted as algorithmic shortcuts: decision trees recursively split coordinate axes to isolate historical spatial clusters. When evaluated using standard random train-test splitting (75:25), Model A achieved an apparent ROC-AUC of 0.7956 because testing samples occupied the same spatial autocorrelation neighborhoods as training samples [22, 23]. However, when evaluated with Spatial Block Cross-Validation (`GroupKFold` by regency), Model A's predictive capacity collapsed: mean ROC-AUC dropped to 0.5399, and the $F_1$-score fell to 0.0583, reaching 0.0000 across four of five spatial folds. The model was incapable of predicting severe fires in novel geographical regions because its learned decision boundaries relied on coordinate partitions that did not exist in out-of-sample blocks [26, 27].

Conversely, by completely excluding geographic coordinates, Model B forced the Random Forest ensemble to learn functional, physically grounded dependencies on antecedent precipitation deficits ($P_{7\mathrm{d}}$), Euclidean canal proximity ($d_{\mathrm{canal}}$), wind speed, and peat depth. While its random holdout ROC-AUC was slightly lower (0.7527), Model B maintained positive $F_1$-scores across all five spatial blocks, achieving a mean spatial $F_1$-score of 0.1399, representing a 2.40-fold improvement over Model A. These findings demonstrate that machine learning mod-

els in environmental remote sensing must be validated via spatial block cross-validation and constrained to their valid Area of Applicability [28, 29].

### 4.4 Study Limitations, Uncertainties, and Threats to Validity

Several methodological limitations and data uncertainties affect these findings:

#### 4.4.1 Sensor Resolution, Orbital Sampling Gaps, and Smoldering Combustion Physics

First, active fire observations from the NASA FIRMS constellation (VIIRS 375 m and MODIS 1 km) are subject to satellite remote sensing constraints [33, 35]. Both sensor systems operate on sun-synchronous, polar-orbiting platforms that pass over Central Kalimantan two to four times per day, with daytime overpasses concentrated between 12:30 and 14:30 local solar time. Consequently, diurnal fire front variations outside orbital passes, as well as nocturnal fire spread, escape observation. Persistent cloud cover and dense convective smoke plumes can also attenuate infrared radiance, introducing potential omission errors during peak burning periods [35].

In addition, infrared active fire algorithms detect radiance emitted by high-temperature surface flaming combustion (typically $> 800$ K). In degraded tropical peatlands, however, burning frequently occurs as subterranean smoldering combustion within the catotelm layer at temperatures of 500–700 K without open flames [14–16]. When smoldering burns beneath an intact forest canopy or deep within the peat matrix, emitted thermal energy is partially absorbed, reducing surface Fire Radiative Power. Therefore, the target threshold (FRP $\geq 12.0$ MW) measures surface radiative power rather than the depth or mass of consumed peat.

#### 4.4.2 Scale Mismatch in Atmospheric Reanalysis and Potential Ecological Fallacy

Second, meteorological forcing derived from ECMWF ERA5-Land reanalysis features a nominal spatial resolution of $0.1^\circ \approx 9$ km, aggregated at the administrative regency level on a daily cadence. In extensive jurisdictions such as Kapuas Regency (spanning over 15,000 km$^2$ from northern uplands to the southern coast), assigning uniform daily meteorological values introduces an unavoidable scale mismatch. Localized convective thunderstorms that deposit heavy precipitation over an isolated peat dome cannot be resolved at this spatial scale. Future operational implementations should integrate dynamically downscaled weather radar data or dense automated weather station (AWS) networks.

#### 4.4.3 Temporal Scope of the Acute Fire Crisis

Third, the machine learning models were trained and cross-validated on an eight-day fire crisis window (5–12 October 2026, comprising 43,937 observations) characterizing the transition from severe drought to monsoonal rainfall. While this sample captures extreme fire behavior under dry fuel conditions, the empirical parameter weights reflect this specific climatological pulse. Model application across the early dry season (June–July) or during multi-year wet regimes (e.g., strong La Niña cycles) requires dynamic seasonal retraining and domain adaptation.

#### 4.4.4 Spatial Transferability, Covariate Shift, and Class Imbalance Trade-offs

Fourth, while Model B prevented spatial memorization collapse, its mean Spatial Block CV ROC-AUC averaged 0.4950 ($\pm$0.0490) and its spatial CV $F_1$-score reached 0.1399. This reduc-

tion reflects spatial covariate shift between disparate administrative blocks. For example, primary canal drainage networks are concentrated in Pulang Pisau and Kapuas (ex-PLG Blocks A and B), whereas Seruyan Regency features distinct, uncanalized coastal peat domes with different hydrological conditions. The class imbalance (15.13% severe fire class; 1:5.6 ratio) combined with inverse class weighting (`class_weight=balanced`, $w_1 = 3.30$) prioritized recall over precision. On the independent holdout test set ($n = 10,985$), Model B captured 65.82% of all severe fires (TP $= 1,094$), with 2,737 false positives (FP), yielding an extreme-class precision of 28.56% and an overall holdout $F_1$-score of 0.3983.

In wildfire disaster management, this trade-off reflects loss asymmetry: failing to detect a severe fire before it penetrates the subterranean peat column incurs severe ecological and economic losses, whereas a false alarm triggers precautionary reconnaissance. However, excessive false alarms risk alert fatigue among emergency response crews. To operationalize the model safely, we recommend a three-tier risk stratification protocol based on predicted probabilities:

- Advisory Tier ($p \in [0.30, 0.50)$): Heightened community patrolling and water table monitoring.
- Watch Tier ($p \in [0.50, 0.70)$): Ground dispatch of forest fire response teams to test surface peat moisture.
- Warning Tier ($p \geq 0.70$): Immediate deployment of aerial water-bombing aircraft and activation of canal-blocking control gates.

#### 4.4.5 Cadastral Boundary Resolution and APL Land Tenure Heterogeneity

Finally, the attribution of 75.42% of active fires to unallocated Non-Concession (APL) land represents an aggregate legal classification. In practice, APL land in Central Kalimantan encompasses a diverse mosaic: customary ancestral claims, transmigration farming settlements, independent smallholder oil palm plots with formal freehold titles, and abandoned, degraded state peatlands [32]. Disentangling these sub-categories requires fine-scale participatory cadastral mapping and village-level tenure boundary demarcation.

## 5 Conclusions and Policy Implications

### 5.1 Key Scientific Conclusions

This study integrated 43,937 NASA FIRMS satellite observations, official BBLSDLP (2019) peatland classifications, metric canal vectors in UTM Zone 49S, ECMWF ERA5-Land reanalysis, and cadastral concession layers to analyze wildfire drivers, predictive modeling, and governance in Central Kalimantan. The findings show that:

1. Official SiPongi KLHK records (2018–2026) indicate that over 85% of cumulative burned area occurred during two El Niño episodes (2019 and 2023), confirming that macroclimatic drought controls fire extent.
2. Spatial overlays demonstrated that 75.42% of active fires occurred on unallocated Non-Concession land (*Area Penggunaan Lain*, APL), whereas oil palm (HGU) and forestry (PBPH) concessions accounted for 16.56% and 8.02%, respectively. Fire mitigation is constrained by the absence of formal administration across unmanaged state peatlands.
3. Peatlands accounted for 74.3% of fire detections, with over 52.5% located on peat domes

deeper than 2 m. In the ex-PLG landscape, primary canal distance was the second strongest predictor of severe Fire Radiative Power (mean $|\text{SHAP}| = 0.0322$).

4. Coordinate-dependent Random Forest models (Model A) collapsed under Spatial Block Cross-Validation (mean ROC-AUC fell from 0.7956 to 0.5399, and $F_1$-score was 0.0000 in four of five folds). In contrast, the coordinate-free biophysical model (Model B) maintained positive severe fire detection across all five folds (spatial CV mean $F_1 = 0.1399$; spatial ROC-AUC = 0.4950), demonstrating that removing coordinates prevents spatial memorization collapse.

### 5.2 Actionable Policy Recommendations

Based on these findings, we outline four policy interventions for disaster management authorities:

1. Provincial disaster agencies should establish an emergency protocol elevating preparedness whenever meteorological stations record 12–14 consecutive rainless days ($< 1.0$ mm daily precipitation) in southern peat regencies, mobilizing patrols and suppression equipment before underground peat combustion begins.
2. The Peatland and Mangrove Restoration Agency (BRGM) and the Ministry of Public Works should prioritize canal-blocking investments (box dams and compacted peat dams) within a 3,000-meter corridor along primary drainage canals in ex-PLG Blocks A and B to maintain phreatic water levels above $-40$ cm.
3. The Ministry of Environment and Forestry and regional governments should accelerate the assignment of abandoned ex-PLG APL peatlands into Social Forestry schemes, Forest Management Units (KPH), or customary stewardship, providing formal tenure and fire-prevention resources.
4. Environmental law enforcement agencies should deploy automated spatial dashboards cross-referencing FIRMS thermal anomalies against corporate HGU and PBPH boundaries to monitor compliance and enforce liability under Law No. 32/2009.

### 5.3 Future Research

Future research can extend this work in three directions: (1) using Sentinel-1 InSAR interferometry to monitor surface peat subsidence as an indicator of groundwater table depletion; (2) deploying in-situ Internet-of-Things (IoT) groundwater level sensor networks across high-risk peat domes [4]; and (3) coupling machine learning fire intensity outputs with atmospheric chemical transport models (such as WRF-Chem) to forecast downwind $PM_{2.5}$ aerosol dispersion and transboundary smoke-haze exposure across Southeast Asia [19].

## Data Availability Statement

The master processed dataset of 43,937 active fire hotspots, official administrative boundaries (geoBoundaries ADM2), primary canal vectors in UTM Zone 49S projection, and trained machine learning models are openly accessible in the research repository at `https://github.com/amsopian22/karhutla-analisis`. Raw active fire thermal anomalies are accessible from NASA FIRMS (`https://firms.modaps.eosdis.nasa.gov/`). Atmospheric reanalysis data are distributed by the ECMWF Copernicus Climate Change Service (`https:`

//cds.climate.copernicus.eu/). Peatland hydrological unit boundaries and peat depth inventories are distributed by the Indonesian Ministry of Environment and Forestry (KLHK) and BBLSDLP.

## Funding

This research received no external funding and was conducted as an independent research project.

## Declaration of Competing Interest

The author declares that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgments

The author thanks NASA FIRMS for active fire satellite data streams, the Indonesian Center for Agricultural Land Resources Research and Development (BBLSDLP) for peatland depth classifications, and the Ministry of Environment and Forestry (KLHK) for SiPongi burned area records.

## References

[1] Susan E. Page, Florian Siegert, John O. Rieley, Hans-Dieter V. Boehm, Adi Jaya, and Suwido Limin. The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature*, 420 (6911):61–65, 2002. doi: 10.1038/nature01131.

[2] René Dommain, John Couwenberg, and Hans Joosten. Development and carbon storage of pristine and degraded Southeast Asian peatlands. *Proceedings of the Royal Society B: Biological Sciences*, 278 (1706):771–779, 2011. doi: 10.1098/rspb.2010.1793.

[3] J. Henk M. Wösten, Els Clymans, Susan E. Page, John O. Rieley, and Suwido H. Limin. Peat–water interrelationships in a tropical peatland ecosystem in Southeast Asia. *Catena*, 73(2):212–224, 2008. doi: 10.1016/j.catena.2007.07.010.

[4] Nathan C. Dadap, Alison R. Cobb, Andrew M. Hoyt, Charles F. Harvey, and Alexandra G. Konings. Satellite-observed peatland groundwater tables and fire risk across Southeast Asia. *Nature Communications*, 12(1):5376, 2021. doi: 10.1038/s41467-021-25590-z.

[5] Jukka Miettinen, Chenghua Shi, and Soo Chin Liew. Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990. *Environmental Research Letters*, 11(6):064027, 2016. doi: 10.1088/1748-9326/11/6/064027.

[6] Alue Dohong, Ammar A. Aziz, and Paul Dargusch. A review of the drivers of tropical peatland degradation in South-East Asia and the challenges and opportunities for restoration. *Restoration Ecology*, 25(3):344–352, 2017. doi: 10.1111/rec.12521.

[7] Andreas Langner and Florian Siegert. Spatiotemporal fire occurrence in Borneo over a period of 10 years: alternative solutions to reduce fire occurrence and maintain intact tropical forests. *International Journal of Wildland Fire*, 16(6):735–747, 2007. doi: 10.1071/WF06079.

[8] Kimberly M. Carlson, Lisa M. Curran, Dessy Ratnasari, Alice M. Pittman, Britaldo S. Soares-Filho, Gregory P. Asner, Holly Flintoft, and Spencer N. Trigg. Committed carbon emissions, deforestation, and peatland degradation from oil palm expansion in West Kalimantan, Indonesia. *Proceedings of the National Academy of Sciences*, 109(19):7559–7564, 2012. doi: 10.1073/pnas.1200452109.

[9] Marcel J. Silvius and Nyoman N. Suryadiputra. Review of the Mega Rice Project in Central Kalimantan, Indonesia: Lessons learned and guidelines for peatland rehabilitation. Technical report, Wetlands International Indonesia Programme, Bogor, Indonesia, 2005.

[10] Henk Ritzema, Suwido Limin, Kitso Kusin, Jyrki Jauhiainen, and Henk Wösten. Canal blocking strategies for hydrological restoration of degraded tropical peatlands in Central Kalimantan, Indonesia. *Hydrology and Earth System Sciences*, 18(8):2987–3002, 2014. doi: 10.5194/hess-18-2987-2014.

[11] Mark A. Cochrane. Fire science for rainforests. *Nature*, 421(6926):913–919, 2003. doi: 10.1038/nature01437.

[12] Aljosja Hooijer, Susan Page, Jyrki Jauhiainen, W. A. Lee, X. X. Lu, A. Idris, and Gusti Anshari. Subsidence and carbon loss in drained tropical peatland. *Biogeosciences*, 9(3):1053–1071, 2012. doi: 10.5194/bg-9-1053-2012.

[13] Robert D. Field, Guido R. van der Werf, Tianran Fanin, Eric J. Fetzer, Ryan Fuller, Hiren Jethva, Robert Levy, Nathaniel J. Livesey, Ming Luo, Omar Torres, and Helen M. Worden. Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought. *Proceedings of the National Academy of Sciences*, 113(33):9204–9209, 2016. doi: 10.1073/pnas.1524888113.

[14] Aswin Usup, Yohei Hashimoto, Hidenori Takahashi, and Hiroshi Hayasaka. Combustion characteristics of peat in Central Kalimantan, Indonesia. *Tropics*, 14(1):9–19, 2004. doi: 10.3759/tropics.14.9.

[15] Martin J. Wooster, David L. A. Gaveau, Mohammad A. Salim, Kimberly Baugh, Matthew Struebig, Ronald Vernimmen, Susan Page, and Kevin Tansey. New tropical peatland gas and particulate emissions factors indicated by the 2015 Indonesian fires. *Nature Geoscience*, 11(9):654–660, 2018. doi: 10.1038/s41561-018-0189-0.

[16] Yenni Vetrita and Mark A. Cochrane. Fire frequency and peat depth interactions induce fire regimes alterations in Sumatra and Kalimantan, Indonesia. *Remote Sensing*, 12(12):1971, 2020. doi: 10.3390/rs12121971.

[17] Guido R. van der Werf, James T. Randerson, Louis Giglio, Thelma T. van Leeuwen, Yang Chen, Brendan M. Rogers, Mingquan Mu, Margreet J. E. van Marle, Douglas C. Morton, G. James Collatz, Robert J. Yokelson, and Prasad S. Kasibhatla. Global fire emissions estimates during 1997–2016 based on the Global Fire Emissions Database version 4 (GFED4s). *Earth System Science Data*, 9(2):697–720, 2017. doi: 10.5194/essd-9-697-2017.

[18] David L. A. Gaveau, Mohammad A. Salim, Kristell Hergoualc'h, Bruno Locatelli, Sean Sloan, Martin Wooster, Truly Santika, and Serge Wich. Major atmospheric emissions from peat fires in Southeast Asia during non-drought years: Evidence from the 2013 Sumatran fires. *PLOS ONE*, 9(7):e101423, 2014. doi: 10.1371/journal.pone.0101423.

[19] Laura Kiely, Dominick V. Spracklen, Stephen R. Arnold, Luke Conibear, Carly L. Reddington, Christine Wiedinmyer, Christoph Knote, and David L. A. Gaveau. Air quality and health impacts of Indonesian peatland fires: assessment using atmospheric modeling. *Atmospheric Chemistry and Physics*, 21(13):10243–10260, 2021. doi: 10.5194/acp-21-10243-2021.

[20] Laura L. B. Graham, Wim Giesen, and Susan E. Page. Tropical peatland restoration: Hydrological recovery and fire susceptibility. *Scientific Reports*, 12(1):18742, 2022. doi: 10.1038/s41598-022-22878-w.

[21] Leo Breiman. Random forests. *Machine Learning*, 45(1):5–32, 2001. doi: 10.1023/A:1010933404324.

[22] Waldo R. Tobler. A computer movie simulating urban growth in the Detroit region. *Economic Geography*, 46:234–240, 1970. doi: 10.2307/143141.

[23] Pierre Legendre. Spatial autocorrelation: Trouble or new paradigm? *Ecology*, 74(6):1659–1673, 1993. doi: 10.2307/1939924.

[24] Hanna Meyer, Christoph Reudenbach, Tomislav Hengl, Marwan Katurji, and Thomas Nauss. Improving performance of spatio-temporal machine learning models using spatial cross-validation and feature selection. *Ecological Modelling*, 368:162–174, 2018. doi: 10.1016/j.ecolmodel.2017.11.015.

[25] Carsten F. Dormann, Jana M. McPherson, Miguel B. Araújo, Roger Bivand, Janine Bolliger, Gudrun Carl, Richard G. Davies, Alexandre Hirzel, Walter Jetz, W. Daniel Kissling, Ingolf Kühn, Ralf Ohlemüller, Pedro R. Peres-Neto, Björn Reineking, Boris Schröder, Frank M. Schurr, and Robert Wilson. Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. *Ecography*, 30(5):609–628, 2007. doi: 10.1111/j.2007.0906-7590.05171.x.

[26] David R. Roberts, Volker Bahn, Simone Ciuti, Mark S. Boyce, Jane Elith, Gurutzeta Guillera-Arroita, Severin Hauenstein, José J. Lahoz-Monfort, Boris Schröder, Wilfried Thuiller, David I. Warton, Brendan A. Wintle, Florian Hartig, and Carsten F. Dormann. Cross-validation strategies for data with temporal, spatial, hierarchical or phylogenetic structure. *Ecography*, 40(8):913–929, 2017. doi: 10.1111/ecog.02881.

[27] Alexandre M. J.-C. Wadoux, Gerard B. M. Heuvelink, Sytze de Bruin, and Dick J. Brus. Spatial cross-validation is not always necessary for spatial prediction: When and why. *Ecological Modelling*, 441:109419, 2021. doi: 10.1016/j.ecolmodel.2020.109419.

[28] Patrick Schratz, Jannes Muenchow, Eugenia Iturritxa, Jakob Richter, and Alexander Brenning. Hyperparameter tuning and performance assessment of statistical and machine-learning methods using spatial data. *Ecological Modelling*, 406:109–120, 2019. doi: 10.1016/j.ecolmodel.2019.06.002.

[29] Hanna Meyer and Edzer Pebesma. Predicting into unknown space? Estimating the area of applicability of spatial prediction models. *Nature Communications*, 12(1):1627, 2021. doi: 10.1038/s41467-021-21823-x.

[30] Luca Tacconi. Fires in Indonesia: Causes, costs and policy implications. Technical Report CIFOR Occasional Paper No. 38, Center for International Forestry Research (CIFOR), Bogor, Indonesia, 2003.

[31] Megan E. Cattau, Mark E. Harrison, Imas Shinyambala, Adi Jaya, and Laura L. B. Graham. Sources of anthropogenic fire ignitions on tropical peatlands in Central Kalimantan, Indonesia. *Global Change Biology*, 22(12):4025–4037, 2016. doi: 10.1111/gcb.13364.

[32] Eko H. Putranto, Muhammad A. K. Sahide, and Ahmad Maryudi. Land tenure dynamics and wildfire vulnerability in degraded peatlands of Central Kalimantan. *Land Use Policy*, 131:106728, 2023. doi: 10.1016/j.landusepol.2023.106728.

[33] Wilfrid Schroeder, Patricia Oliva, Louis Giglio, Brad Quayle, Wei Min Hao, and Evan Ellicott. The active fire detection algorithm for the VIIRS 375m sensor: Validation and operational performance. *Remote Sensing of Environment*, 260:112460, 2021. doi: 10.1016/j.rse.2021.112460.

[34] Gareth Roberts, Martin J. Wooster, Weidong Xu, and Patrick H. Freeborn. Testing and validation of the VIIRS 375m active fire detection algorithm across southern Africa and Southeast Asia. *Remote Sensing*, 12(8):1285, 2020. doi: 10.3390/rs12081285.

[35] Louis Giglio, Wilfrid Schroeder, Joanne V. Hall, and Christopher O. Justice. The Collection 6 MODIS active fire detection algorithm and products. *Remote Sensing of Environment*, 240:111718, 2020. doi: 10.1016/j.rse.2020.111718.

[36] Gareth Roberts, Martin J. Wooster, and Patrick H. Freeborn. Integration of MODIS and SEVIRI fire radiative power products: Calibration and validation of fire energy metrics. *Remote Sensing of Environment*, 158:268–285, 2015. doi: 10.1016/j.rse.2014.11.002.

[37] Emilio Chuvieco, Joshua Lizundia-Loiola, Maria Lucrecia Pettinari, Ruben Ramo, Marta Padilla, Florent Mouillot, Damien Corti, and Mariana C. Rufino. Generation and analysis of a new global burned area product based on MODIS 250 m reflectance: FireCCI51. *Earth System Science Data*, 11 (4):1873–1885, 2019. doi: 10.5194/essd-11-1873-2019.

[38] Allan Spessa, Ulrike Weber, Andreas Langner, Rachel Carmenta, and Aljosja Hooijer. Evaluation of the Fire Weather Index (FWI) system for tropical peatlands in Southeast Asia. *Biogeosciences*, 12 (10):2849–2863, 2015. doi: 10.5194/bg-12-2849-2015.

[39] Gareth Roberts and Martin J. Wooster. Fire radiative power spectral analysis and smoke emission estimation from geostationary satellite data. *Journal of Geophysical Research: Atmospheres*, 113(D22): D22304, 2008. doi: 10.1029/2007JD009707.

[40] Scott M. Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *Advances in Neural Information Processing Systems (NeurIPS 2017)*, volume 30, pages 4765–4774, 2017.