

ANTICOAGULANT CONVERSION CHART

Newsletter by: PRIC Hospital Sg Buloh, Pua Fu Siang & Ooi Yu Xin

To From	Warfarin	Dabigatran	Rivaroxaban	Apixaban	Edoxaban	Parenteral
LMWH	Overlapped at least 5 days OR until 2 consecutive days with INR in therapeutic range. ^{1,2}	Start DOAC at the next scheduled dose of LMWH/Fondaparinux. ^{1,3,4,8}				Treatment: ² Stop LMWH/Fondaparinux; start IV UFH 1-2 hours before next dose of LMWH/Fondaparinux. Prophylactic: ⁵ Stop LMWH/Fondaparinux; start SC UFH at the next scheduled dose of LMWH/Fondaparinux.
Fondaparinux						
Continuous UFH		Start DOAC at time of discontinuation of continuous UFH. ^{1,3}				Discontinue the UFH infusion; Start Edoxaban 4 hours after discontinuation of UFH infusion. ⁸ Stop continuous UFH; start LMWH/ Fondaparinux within 1 hour. ²
Warfarin		Stop Warfarin; start Dabigatran when INR < 2.0. ^{1,3,6}	AF: ³ Stop Warfarin; start Rivaroxaban when INR ≤ 3.0. DVT or PE: ³ Stop Warfarin; start Rivaroxaban when INR ≤ 2.5.	Stop Warfarin; start Apixaban when INR < 2.0. ^{1,3,4}	Stop Warfarin; start Edoxaban when INR ≤ 2.5. ⁸	Stop Warfarin; start parenteral anticoagulant when INR < 2.0. ⁷
Dabigatran	Option 1: Overlapping ⁶ Start Warfarin, then if: CrCl ≥ 50ml/min: Stop Dabigatran 3 days later CrCl 30-50ml/min: Stop Dabigatran 2 days later CrCl 15-30ml/min: Stop Dabigatran 1 day later CrCl <15ml/min: Dabigatran not recommended. ²					CrCl ≥ 30 ml/min: ⁷ Stop Dabigatran; start parenteral agent 12 hours after last Dabigatran. CrCl < 30 ml/min: ⁷ Stop Dabigatran; start parenteral agent 24 hours after last Dabigatran. (Note: Fondaparinux contraindicated)
Rivaroxaban/ Apixaban	Option 1: Overlapping ^{2,6} Start Warfarin; Continued for at least two days then stop DOAC when INR in therapeutic range. Option 2: Bridging ^{2,6} Stop DOAC; start Warfarin with parenteral agent at next scheduled dose of DOAC. Cont. parenteral agent until INR in therapeutic range.	Stop current DOAC; start new DOAC at the next scheduled dose of previous DOAC. ^{6, 8}				
Edoxaban	Option 1: Overlapping ⁸ INR to be measured before Edoxaban is administered. Start Warfarin (loading dose not required); Reduce the dose of Edoxaban by 50% for up to 14 days and stop when INR in therapeutic range. Option 2: Bridging ⁸ Stop Edoxaban; Start Warfarin with parenteral agent at next scheduled dose of edoxaban. Cont. parenteral agent until INR in therapeutic range.					Stop Rivaroxaban/Apixaban/Edoxaban; start parenteral agent at the next scheduled dose of Rivaroxaban/Apixaban/Edoxaban. ^{1,7,8}

Only for internal circulation. For further enquiries, kindly contact ext 4126. Revised July 2024.

References

1. CPG Prevention and Treatment of Venous Thromboembolism. 2013; 1-168.
2. Lexicomp® Drug Information handbook. 27th edition. Wolters Kluwer; 2018.
3. Anticoagulation MTAC (AC-MTAC) Protocol. Pharmaceutical Services Program 2020; 2:1-86.
4. Lofgren B. How to Convert Between Anticoagulants. Pharmacy Times. 2015.
5. Gonzalez C, Franco-Martinez C. Transitioning Between Anticoagulants. University Health System. Jan 2014.
6. Leung LLK, Mannucci PM, Tirnauer JS. Direct oral anticoagulants (DOACs) and parenteral direct-acting anticoagulants: Dosing and adverse effects. UpToDate 2020.
7. Switching To and From Various Anticoagulants. Minneapolis Heart Institute®, Abbott Northwestern Hospital. Aug 2016.
8. Lixiana [Product Insert]. Germany: Daiichi Sankyo Europe; 13 Sept 2023.

