





DOAC DOSING GUIDE FOR VENOUS THROMBOEMBOLISM (VTE)

PREPARED BY PHARMACY DEPT HOSP SG BULOH. UPDATED JUNE 2024
FOR INTERNAL CIRCULATION ONLY

- Min 5 days of treatment with parenteral anticoagulant is required before initiation
- 110mg BD dose is not studied in clinical trial, only based on PKPD analyses



FOOTNOTE FOR ALL DOAC

- Not recommended in pregnancy/lactation
- Not recommended when liver enzyme elevated (ALT/AST >2x ULN)
- For extremes of body weight (< 50kg, > 120kg or BMI > 40kg/m²) please consult your pharmacist.
- For further enquiries, please contact Pharmacy.
- TE : Thromboembolic

- Min 5 days of treatment with parenteral anticoagulant is required before initiation

DOAC DOSING GUIDE FOR ATRIAL FIBRILLATION (AF)

PREPARED BY PHARMACY DEPT HOSP SG BULOH. UPDATED JUNE 2024

FOR INTERNAL CIRCULATION ONLY

DABIGATRAN

CrCl

< 30 mL/min

CONTRAINDICATED

30 – 50 mL/min

> 50 mL/min

- Age ≥ 80 years
- Concomitant Verapamil / Amiodarone

- Low TE risk
- High bleeding risk

110mg
BD

110mg
BD

150mg
BD

150mg
BD

110mg
BD

150mg
BD

110mg
BD

RIVAROXABAN

CrCl

< 15 mL/min

CONTRAINDICATED

15 – 49 mL/min

15mg OD

≥ 50 mL/min

20mg OD

FOOTNOTE

- Not recommended in pregnancy/lactation
- Not recommended when liver enzyme elevated (ALT/AST >2x ULN)
- For extremes of body weight (< 50kg, > 120kg or BMI > 40kg/m²) please consult your pharmacist.
- For further enquiries, please contact Pharmacy.
- TE : Thromboembolic

APIXABAN

CrCl

15 – 29 mL/min

2.5mg BD

< 15 mL/min

CONTRAINDICATED

* US-FDA approved for usage
in patient on **dialysis**
(based on PK studies)

≥ 30 mL/min

≥ 80 yrs old

≤ 60kg

SCr ≥ 133
umol/L

If ≥ 2 features

2.5mg BD

If ≤ 1 feature

5mg BD

EDOXYBAN

CrCl

15 – 50 mL/min

30mg OD

51 – 95 mL/min

> 95 mL/min

Not recommended

< 15 mL/min

CONTRAINDICATED

≤ 60kg

30mg OD

Concomitant Cyclosporin/Dronedarone
/Erythromycin/Ketoconazole

> 60kg

60mg OD