

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

Take home message:

- DAPT duration is dependent on the indication of use.
- DAPT duration may be individualized depending on patient’s ischaemic and bleeding risk.
- Newer generation stents (e.g. Resolute Onyx, Xience, Ultimaster, Orsiro, Biofreedom) are approved for short DAPT duration as short as 1 month only.

Disclaimer:

The recommendations listed in this document is intended for educational & informational purposes. It is NOT intended to take the place of physician judgement in diagnosing and treatment of particular patients. DAPT duration may be individualized depending on patient’s ischemic and bleeding risk.

Abbreviation & Description:

NSTE-ACS	Non-ST elevation acute coronary syndrome (Combination of unstable angina & NSTEMI)
CABG	Coronary artery bypass surgery
STEMI	ST elevation myocardial infarction
CAD	Coronary artery disease
Stable CAD	• Stable chest pain or other symptoms (e.g. dyspnoea) which are known to be due to CAD. • Patient who had a previous episode of ACS but stable now. • Post revascularization (by CABG or PCI). • Patient who are asymptomatic but are known to have CAD on non-invasive testing. This may occur in the absence or presence of ischemia and/or Left Ventricular (LV) dysfunction.
DAPT	Dual antiplatelet
OAC	Oral anticoagulant
AF	Atrial fibrillation
PCI	percutaneous coronary intervention
MI	Myocardial infarction
TIA	Transient ischaemic attack
P2Y12 Inhibitor	E.g. Clopidogrel, Ticagrelor, Ticlopidine, Prasugrel

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

NSTE-ACS^{[1], [2], [5], [7]}

Elective PCI with

bare metal stents/drug eluting stent/
drug coated balloon

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

High
Bleeding Risk
(PRECISE-DAPT≥25)

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

High
Bleeding Risk
(PRECISE-DAPT≥25)

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

High
Bleeding Risk
(PRECISE-DAPT≥25)

Medical Therapy Only

CABG

(Coronary Artery Bypass Surgery)

3 - 6 months*

3 - 6 months

To complete
total duration
of **12 months**
of DAPT
after CABG
(including
DAPT duration
before CABG)

To complete
total duration
of **6 months**
of DAPT
after CABG
(including
DAPT duration
before CABG)

After completed DAPT, re-assess the risks and benefits for prolong DAPT by using DAPT score.

DAPT score <2: Stop DAPT

DAPT score ≥2: Prolong DAPT

Long term aspirin monotherapy

Clopidogrel monotherapy if aspirin is not tolerated.

* = DAPT duration can be as short as 1 - 3 months depending on type of stent.

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

STEMI ^{[2], [3], [5], [7]}

Elective PCI with

bare metal stents/drug eluting
stent/drug coated balloon

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

≥12 months

High
Bleeding Risk
(PRECISE-DAPT≥25)

≥6 months*

Fibrinolytic therapy

1 - 12 months

Medical Therapy Only

Ticagrelor is preferred over clopidogrel for 1st year of treatment

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

≥12 months

High
Bleeding Risk
(PRECISE-DAPT≥25)

3 - 6 months

CABG

(Coronary Artery Bypass Surgery)

Low-Moderate
Bleeding Risk
(PRECISE-DAPT<25)

To complete
total duration
of **12 months**
of DAPT
after CABG
(including
DAPT duration
before CABG)

High
Bleeding Risk
(PRECISE-DAPT≥25)

To complete
total duration
of **6 months**
of DAPT
after CABG
(including
DAPT duration
before CABG)

After completed DAPT, re-assess the risks and benefits for prolong DAPT by using DAPT score.

DAPT score <2: Stop DAPT

DAPT score ≥2: Prolong DAPT


Long term aspirin monotherapy

Clopidogrel monotherapy if aspirin is not tolerated.

* = DAPT duration can be as short as 1 - 3 months depending on type of stent.

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting




* = DAPT duration can be as short as 1 - 3 months depending on type of stent.

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

Patient on chronic oral anticoagulant
(e.g. AF, valvular heart disease)

[2], [3], [6], [7], [12]



Yellow box = Triple therapy (DAPT + OAC)

Red box = P2Y12 inhibitor + OAC
(Clopidogrel is preferred)


Notes:

- Triple therapy should be as short as possible.
- Consider target INR 2.0 - 2.5 if warfarin is used.
- Aspirin: Low dose (75 - 100mg) is preferred to minimize bleeding risk.

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

Ischaemic stroke ^{[4], [5], [9], [10], [13], [14]}



** = DAPT duration may based on interventional radiologist's recommendation

General Recommended Duration Dual AntiPlatelet Therapy (DAPT)

Prepared by PRIC, Sia Hee Peng & Pee Lay Ting

References:

- [1] CPG NSTE-ACS 3rd edition 2021
- [2] Levine, G. N., Bates, E. R., Bittl, J. A., Brindis, R. G., Fihn, S. D., Fleisher, L. A., ... Smith, S. C. (2016). 2016 ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: A report of the american college of cardiology/american heart association task force on clinical practice guidelines: An update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention, 2011 ACCF/AHA guideline for Coronary Artery Bypass Graft Surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease, 2013 ACCF/AHA guideline for the management of St-Elevation Myocardial Infarction, 2014 AHA/ACC guideline for the management of patients with non-st-elevation acute coronary syndromes, and 2014 ACC/AHA guideline on Perioperative Cardiovascular Evaluation and management of patients undergoing noncardiac surgery. *Circulation*, 134(10). Retrieved from <https://doi.org/10.1161/cir.0000000000000404>
- [3] CPG STEMI 4th edition 2019
- [4] CPG Ischaemic stroke 3rd edition 2020
- [5] CPG Primary & Secondary Prevention of Cardiovascular Disease 2017
- [6] Costa, F., & Valgimigli, M. (2019). Novel directions for the management of dual antiplatelet therapy in patients with coronary artery disease. *Cardiovascular Medicine*. Retrieved from <https://doi.org/10.4414/cvm.2019.02024>
- [7] Valgimigli, M., Bueno, H., Byrne, R. A., Collet, J.-P., Costa, F., Jeppsson, A., ... Kinnaird, T. (2017). 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with eacts. *European Heart Journal*, 39(3), 213–260. Retrieved from <https://doi.org/10.1093/eurheartj/ehx419>
- [8] Gargiulo, G., Valgimigli, M., Capodanno, D., & Bittl, J. (2017). State of the art: Duration of dual antiplatelet therapy after percutaneous coronary intervention and coronary stent implantation – past, present and future perspectives. *EuroIntervention*, 13(6), 717–733. Retrieved from <https://doi.org/10.4244/eij-d-17-00468>
- [9] Antiplatelet therapy for ischemic stroke and transient ischemic attack. *Canadian Stroke Best Practices*. (n.d.). Retrieved from <https://www.strokebestpractices.ca/recommendations/secondary-prevention-of-stroke/anti-platelet-therapy-in-ischemic-stroke-and-tia#:~:text=For%20patients%20with%20an%20acute,21%20days%20after%20the%20event%2C>
- [10] Kleindorfer, D. O., Towfighi, A., Chaturvedi, S., Cockroft, K. M., Gutierrez, J., Lombardi-Hill, D., ... Williams, L. S. (2021). 2021 guideline for the prevention of stroke in patients with stroke and transient ischemic attack: A guideline from the American Heart Association/American Stroke Association. *Stroke*, 52(7):e364-e467. Retrieved from <https://doi.org/10.1161/str.0000000000000375>
- [11] Writing Committee Members, Lawton, J. S., Tamis-Holland, J. E., Bangalore, S., Bates, E. R., Beckie, T. M., Bischoff, J. M., Bittl, J. A., Cohen, M. G., DiMaio, J. M., Don, C. W., Fremes, S. E., Gaudino, M. F., Goldberger, Z. D., Grant, M. C., Jaswal, J. B., Kurlansky, P. A., Mehran, R., Metkus, T. S., Jr, Nnacheta, L. C., ... Zwischenberger, B. A. (2022). 2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Journal of the American College of Cardiology*, 79(2), e21–e129. <https://doi.org/10.1016/j.jacc.2021.09.006>
- [12] Van Gelder, I. C., Rienstra, M., Bunting, K. V., Casado-Arroyo, R., Caso, V., Crijns, H. J. G. M., De Potter, T. J. R., Dwight, J., Guasti, L., Hanke, T., Jaarsma, T., Lettino, M., Løchen, M. L., Lumbers, R. T., Maesen, B., Mølgaard, I., Rosano, G. M. C., Sanders, P., Schnabel, R. B., Suwalski, P., ... ESC Scientific Document Group (2024). 2024 ESC Guidelines for the management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *European heart journal*, 45(36), 3314–3414. <https://doi.org/10.1093/eurheartj/ehae176>
- [13] Powers, W. J., Rabinstein, A. A., Ackerson, T., Adeoye, O. M., Bambakidis, N. C., Becker, K., Biller, J., Brown, M., Demerschalk, B. M., Hoh, B., Jauch, E. C., Kidwell, C. S., Leslie-Mazwi, T. M., Ovbiagele, B., Scott, P. A., Sheth, K. N., Southerland, A. M., Summers, D. V., Tirschwell, D. L., & American Heart Association Stroke Council (2018). 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*, 49(3), e46–e110. <https://doi.org/10.1161/STR.0000000000000158>
- [14] Greco, A., Occhipinti, G., Giacoppo, D., Agnello, F., Laudani, C., Spagnolo, M., Mauro, M. S., Rochira, C., Finocchiaro, S., Mazzone, P. M., Faro, D. C., Landolina, D., Ammirabile, N., Imbesi, A., Raffo, C., & Capodanno, D. (2023). Antithrombotic Therapy for Primary and Secondary Prevention of Ischemic Stroke: JACC State-of-the-Art Review. *Journal of the American College of Cardiology*, 82(15), 1538–1557. <https://doi.org/10.1016/j.jacc.2023.07.025>