



What is Dabigatran?

- Dabigatran is an anticoagulant medication.
- It prevents blood clot from occurring continuously in the body while allowing the existing ones to dissolve itself naturally
- It does not dissolve the existing blood clots but reduces the risk of more blood clot formation.

Why must I take Dabigatran?

- For stroke prevention or blood clot from irregular heartbeat (Atrial Fibrillation) OR
- For treatment or prevention of blood clot formation in leg (deep vein thrombosis) and lungs (pulmonary embolism).

Monitoring of Dabigatran:

- Dr will conduct kidney and liver function tests to ensure they are in optimal conditions.
- Ensure to come for clinic appointment on your given appointment date.

Duration:

- Treatment duration is 3 – 6 months, or longer, depending on your condition. **DO NOT STOP** your medication unless directed by your doctor.

Administration:

- Take **TWICE DAILY** (EVERY 12 hours), at the same time every day.
- Only take out a capsule from the blister pack each time. **DO NOT CRUSH, CHEW or BREAK** open the capsule before swallowing as this may increase the risk of bleeding.
- Swallow the **WHOLE capsule** with a full glass of water, with or without food.
- STOP any SUPPLEMENTS/HERBAL MEDICATIONS because of drug interactions.

Storage:

- Store the medication in a cool dry place, away from direct sunlight and in the original packaging in order to protect from moisture
- Store the capsules Do not put the capsules in pill boxes or pill organizers, unless capsules can be maintained in the original package.

What do you do when you missed a dose?

- If you missed a dose, take it as soon as you remember. A forgotten dose may still be taken if it is still more than 6 hours until your next scheduled dose
- If there is less than 6 hours until the next scheduled dose, the missed dose should be omitted and take the next dose as usual
- Do not take a double dose to make up for missed individual doses.

Special Precautions with Dabigatran

- Seek treatment if you have these symptoms:

Bleeding symptoms:

- Bruises
- Nose / Gum bleed
- Coughing or vomiting blood
- Stool with blood (blackish/red color)
- Blood in urine



Renal Dysfunction symptoms:

- Sudden weight gain
- Leg swelling
- Nausea and vomiting
- Skin rashes

Gastrointestinal Disorder symptoms:

- Bloating
- Abdominal pain
- Acid reflux



- You may bleed easily.
- Be careful when brushing teeth, use tooth brush with soft bristle. Use electric shaver instead of razor to prevent from injuries.
- It is important to tell your healthcare providers that you are taking dabigatran if you need to have any surgery, dental work or other medical procedures that increase your risk for bleeding. Dabigatran may need to be stopped temporarily. Do not stop taking your medication unless directed by your doctor.
- Avoid taking alcohol due to interaction with dabigatran which may increase risk of bleeding.



What is Apixaban?

- Apixaban is an anticoagulant medication.
- It prevents blood clot from occurring continuously in the body while allowing the existing ones to dissolve itself naturally
- It does not dissolve the existing blood clots but reduces the risk of more blood clot formation.

Why must I take Apixaban?

- For stroke prevention or blood clot from irregular heartbeat (Atrial Fibrillation) OR
- For treatment or prevention of blood clot formation in leg (deep vein thrombosis) and lungs (pulmonary embolism).

Monitoring of Apixaban:

- Dr will conduct kidney and liver function tests to ensure they are in optimal conditions.
- Ensure to come for clinic appointment on your given appointment date.

Duration:

- Treatment duration is 3 – 6 months, or longer, depending on your condition. **DO NOT STOP** your medication unless directed by your doctor.

Administration:

- Take **TWICE DAILY** (EVERY 12 hours), same time every day.
- Take the tablet with a glass of water.
- If you have difficulty swallowing, the tablet can be crushed and mixed with a small amount of water to be taken immediately
- This medication can be taken before or after meal



What do you do when you missed a dose?

- If you missed a dose, take it as soon as you remember. A forgotten dose may still be taken if it is still more than 6 hours until your next scheduled dose
- If there is less than 6 hours until the next scheduled dose, the missed dose should be omitted and take the next dose as usual
- Do not take a double dose to make up for missed individual doses.

Storage

- Store the medication in a cool dry place, away from direct sunlight.
- Keep the medication away from children.

Special Precautions with Apixaban

- Seek treatment if you have these symptoms:

Bleeding symptoms:

- Bruises
- Nose / Gum bleed
- Coughing or vomiting blood
- Stool with blood (blackish/red color)
- Blood in urine

Renal Dysfunction symptoms:

- Sudden weight gain
- Leg swelling
- Nausea and vomiting
- Skin rashes

Gastrointestinal Disorder symptoms:

- Bloating
- Abdominal pain
- Acid reflux



- You may bleed easily.
- Be careful when brushing teeth, use tooth brush with soft bristle. Use electric shaver instead of razor to prevent from injuries.
- It is important to tell your healthcare providers that you are taking apixaban if you need to have any surgery, dental work or other medical procedures that increase your risk for bleeding. Apixaban may need to be stopped temporarily. Do not stop taking your medication unless directed by your doctor.
- Avoid taking alcohol due to interaction with apixaban which may increase risk of bleeding.

Tab Edoxaban 30mg and 60mg

What is Edoxaban?

- Edoxaban is an anticoagulant medication.
- It prevents blood clot from occurring continuously in the body while allowing the existing ones to dissolve itself naturally
- It does not dissolve the existing blood clots but reduces the risk of more blood clot formation.

Why must I take Edoxaban?

- For stroke prevention or blood clot from irregular heartbeat (Atrial Fibrillation) OR
- For treatment or prevention of blood clot formation in leg (deep vein thrombosis) and lungs (pulmonary embolism).

Monitoring of Edoxaban:

- Dr will conduct kidney and liver function tests to ensure they are in optimal conditions.
- Ensure to come for clinic appointment on your given appointment date.

Duration:

- Treatment duration is 3 – 6 months, or longer, depending on your condition. **DO NOT STOP** your medication unless directed by your doctor.

Administration:

- Take **ONCE DAILY** (EVERY 24 hours), at the same time every day.
- Take the tablet with a glass of water.
- If you have difficulty swallowing, the tablet can be crushed and mixed with a small amount of water to be taken immediately
- This medication can be taken before or after meal



What do you do when you missed a dose?

- If you missed a dose, take it as soon as you remember.
- Do not take a double dose to make up for missed individual doses.

Storage

- Store the medication in a cool dry place, away from direct sunlight.
- Keep the medication away from children.

Special Precautions with Edoxaban

- Seek treatment if you have these symptoms:

Bleeding symptoms:

- Bruises
- Nose / Gum bleed
- Coughing or vomiting blood
- Stool with blood (blackish/red color)
- Blood in urine



Renal Dysfunction symptoms:

- Sudden weight gain
- Leg swelling
- Nausea and vomiting
- Skin rashes

Gastrointestinal Disorder symptoms:

- Bloating
- Abdominal pain
- Acid reflux



- You may bleed easily.
- Be careful when brushing teeth, use tooth brush with soft bristle. Use electric shaver instead of razor to prevent from injuries.
- It is important to tell your healthcare providers that you are taking edoxaban if you need to have any surgery, dental work or other medical procedures that increase your risk for bleeding. Edoxaban may need to be stopped temporarily. Do not stop taking your medication unless directed by your doctor.
- Avoid taking alcohol due to interaction with edoxaban which may increase risk of bleeding.