# **Formulas**

# Formulas and Theorem

#### 

If we have an Orthogonal Basis  $\{ ec{u}_1, \dots, ec{v}_n \}$  in  $\mathbb{R}^n$  then for any  $ec{w} \in \mathbb{R}^n$ ,

$$\vec{w} = c_1 \vec{u}_1 + \cdots + c_n \vec{v}_n$$

 $C_q$  can be found using,

$$c_q = rac{ec{w} \cdot ec{u}_q}{ec{u}_q \cdot ec{u}_q}$$

## 

Let non-zero  $\vec{u} \in \mathbb{R}^n$ , and  $\vec{y} \in \mathbb{R}^n$ . The orthogonal projection of  $\vec{y}$  onto  $\vec{u}$  is the vector in the span of  $\vec{u}$  that is closest to  $\vec{y}$ .

$$\mathrm{proj}_{ec{u}}ec{y} = rac{ec{y} \cdot ec{u}}{ec{u} \cdot ec{u}} ec{u}$$

Also,  $ec{y} = \hat{y} + ec{z}$  and,  $ec{z} \in W^{\perp}$ 

## **⊘** Best Approximation Theorem ∨

Let W be a subspace of  $\mathbb{R}^n, \vec{y} \in \mathbb{R}^n$ , and  $\hat{y}$  is the orthogonal projection of  $\vec{y}$  onto W. Then for any  $\vec{v} \neq \hat{y}, \vec{v} \in W$ , we have

$$||ec{y} - \hat{y}|| < ||ec{y} - ec{v}||$$

#### 

Let W be a subspace of  $\mathbb{R}^n$ . Then, each  $\vec{y} \in \mathbb{R}^n$  has a unique decomposition.

$$ec{y} = \hat{y} + z, \quad \hat{y} \in W, \quad z \in W^{\perp}$$

If  $\vec{u_1}, \dots, \vec{u_n}$  is the orthogonal basis for W,

$$\hat{y} = rac{ec{y} \cdot ec{u_1}}{ec{u_1} \cdot ec{u_1}} ec{u_1} + \cdots + rac{ec{y} \cdot ec{u_n}}{ec{u_n} \cdot ec{u_n}} ec{u_n}$$

 $\hat{y}$  is the orthogonal projection of  $\vec{y}$  onto W

## **⊘** QR Factorization ∨

For a  $m \times n$  matrix A linearly independent columns,

$$A = QR$$

Q is an  $m \times n$ , with columns are an orthonormal basis for ColA. R is  $n \times n$ , upper triangular, with positive entries on its diagonal.

## *⊘* **Normal Equation ∨**

$$A^T A \hat{x} = A^T ec{b}$$

Manipulating this we can get this,

$$\hat{x} = (A^TA)^{-1}A^T \vec{b}$$

## Eigenvectors orthogonality >

If A is a symmetric matrix, with eigenvectors  $\vec{v_1}$  and  $\vec{v_2}$  corresponding to two distinct eigenvalues, then  $\vec{v_1}$  and  $\vec{v_2}$  are orthogonal.

## 

An  $n \times n$  symmetric matrix A has the following properties,

- All eigenvalues of A are real.
- The eigenspaces are mutually orthogonal.
- A can be diagonalized as  $A = PDP^T$ , where D is diagonal and P is orthogonal.