Formulas-3

Fubini's Theorem

$$V = \iint_R f(x,y) dA = \int_c^d \int_a^b f(x,y) dx dy = \int_a^b \int_c^d f(x,y) dy dx$$

if R is defined by $a \leq x \leq b$ and $g_1(x) \leq y \leq g_2(x)$ then,

$$V=\int_a^b\int_{q_1(x)}^{g_2(x)}f(x,y)dydx$$

Notice: The functions $g_n(x)$ are evaluated first.

Properties

$$egin{aligned} &\iint_R cf(x,y)dA = c\iint_R f(x,y)dA \ &\iint_R f(x,y) \pm g(x,y)dA = \iint_R f(x,y)dA \pm \iint_R g(x,y)dA \ &\iint_R f(x,y)dA \geq 0 ext{ if } f(x,y) \geq 0 ext{ on } R \ &\iint_R f(x,y)dA \geq \iint_R g(x,y)dA ext{ if } f(x,y) \geq g(x,y) ext{ on } R \end{aligned}$$

If R_1 and R_2 are non-overlapping regions,

$$\begin{split} \iint_R f(x,y) dA &= \iint_{R_1} f(x,y) dA + \iint_{R_2} f(x,y) dA \\ A &= \iint_R dA = \int_{x_1}^{x_2} \int_{y_1}^{y_2} dy dx = \int_{\theta_1}^{\theta_2} \int_{r_1}^{r_2} r dr d\theta \\ V &= \iint_R f(x,y) dA \\ \text{Average Value} &= \frac{1}{\text{Area of } R} \iint_R f dA \quad \text{OR} \quad \frac{1}{\text{Volume of } D} \iiint_D F dV \\ V &= \iiint_D dV = \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} dz dy dx = \int_{\theta_1}^{\theta_2} \int_{r_1}^{r_2} \int_{z_1}^{z_2} r dz dr d\theta = \int_{\rho_1}^{\rho_2} \int_{\phi_1}^{\phi_2} \int_{\theta_1}^{\theta_2} \rho^2 \sin(\phi) d\theta d\phi d\rho \end{split}$$

Mass and First Moments of 3D Solids

Mass:
$$M = \iiint_{D} \delta dV$$

First Moments:
$$M_{yz} = \iiint_D x \delta dV, M_{xz} = \iiint_D y \delta dV, M_{xy} = \iiint_D z \delta dV$$

Center of mass:
$$ar{x} = rac{M_{yz}}{M}, ar{y} = rac{M_{xz}}{M}, ar{z} = rac{M_{xy}}{M}$$

Mass and First Moments of 2D Plates:

$$\text{Mass: } M = \iint_R \delta dA$$

First Moments:
$$M_y = \iint_R x \delta dA, M_x = \iint_R y \delta dA$$

Center of mass:
$$ar{x} = rac{M_y}{M}, ar{y} = rac{M_x}{M}$$

Moments of Inertia of 3D Solids:

About x-axis:
$$I_x = \iiint_D (y^2 + z^2) \delta dV$$

About y-axis:
$$I_y = \iiint_D (x^2 + z^2) \delta dV$$

About z-axis:
$$I_z = \iiint_D (x^2 + y^2) \delta dV$$

About a line L:
$$I_L = \iiint_D r^2(x,y,z) \delta dV$$

Moments of Inertia of 2D Plates:

About x-axis:
$$I_x = \iint_R y^2 \delta dA$$

About y-axis:
$$I_y = \iint_R x^2 \delta dA$$

About a line L:
$$I_L = \iint_R r^2(x,y) \delta dA$$

About the origin:
$$I_O = \iint_R (x^2 + y^2) \delta dA = I_x + I_y$$

Joint probability density function:

Conditions:

$$f(x,y) \ge 0$$

$$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty}f(x,y)dxdy=1$$

$$P((X,Y) \in R) = \iint_R f(x,y) dx dy$$

Mean and expected value:

$$\mu_X = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x,y) dx dy \ \mu_Y = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x,y) dx dy$$

Spherical Coordinates (ρ, ϕ, θ)

$$egin{aligned} r &=
ho \sin(\phi), \ x &= r \cos(heta) =
ho \sin(\phi) \cos(heta) \ z &=
ho \cos(\phi), \ y &= r \sin(heta) =
ho \sin(\phi) \sin(heta) \
ho &= \sqrt{x^2 + y^2 + z^2} = \sqrt{r^2 + z^2} \end{aligned}$$

Jacobian:

$$J(u,v) = egin{array}{c|c} rac{\partial x}{\partial u} & rac{\partial x}{\partial v} \\ rac{\partial y}{\partial u} & rac{\partial y}{\partial v} \end{array} = rac{\partial (x,y)}{\partial (u,v)} \ & \iint_R f(x,y) dx dy = \iint_G f(g(u,v),h(u,v)) \left| rac{\partial (x,y)}{\partial (u,v)}
ight| du dv \ & J(u,v,w) = \left| rac{rac{\partial x}{\partial u} & rac{\partial x}{\partial v} & rac{\partial x}{\partial w} \ rac{\partial y}{\partial u} & rac{\partial y}{\partial v} & rac{\partial y}{\partial w} \ rac{\partial z}{\partial u} & rac{\partial z}{\partial v} & rac{\partial z}{\partial w} \ \end{array}
ight| = rac{\partial (x,y,z)}{\partial (u,v,w)} \ = \iint_R f(g(u,v,w),h(u,v,w),k(u,v,w)) \left| rac{\partial (x,y,z)}{\partial (u,v,w)}
ight| du dv dw \ .$$