

# **A Couple More Practice Problems**

## **Concepts of Programming Languages**

# S-Combinator

$$\{x : \tau_1, y : \tau_2\} \vdash \lambda z^{\top}. xz(yz) : \top \rightarrow \top$$

*Determine types  $\tau_1$  and  $\tau_2$  such that the above judgment is derivable in STLC, then given the derivation. Are there multiple choices for  $\tau_1$  and  $\tau_2$ ? If so, give another choice and justify your answer*

$$\frac{}{\Gamma \vdash \bullet : \top}$$

$$\frac{(x : \tau) \in \Gamma}{\Gamma \vdash x : \tau}$$

$$\frac{\Gamma, x : \tau \vdash e : \tau'}{\Gamma \vdash \lambda x^{\tau}. e : \tau \rightarrow \tau'}$$

$$\frac{\Gamma \vdash e_1 : \tau \rightarrow \tau' \quad \Gamma \vdash e_2 : \tau}{\Gamma \vdash e_1 e_2 : \tau'}$$

# CBN vs. CBV (Part 1)

*Determine an expression in the **untyped** lambda calculus that terminates using CBN evaluation but does not terminate using CBV evaluation*

# CBN vs. CBV (Part 1)

*Determine an expression in the **untyped** lambda calculus that terminates using CBN evaluation but does not terminate using CBV evaluation*

$$\frac{e_1 \longrightarrow e'_1}{e_1 e_2 \longrightarrow e'_1 e_2} \quad \frac{e_2 \longrightarrow e'_2}{(\lambda x. e_1) e_2 \longrightarrow (\lambda x. e_1) e'_2}$$

$$\frac{}{(\lambda x. e)(\lambda y. e') \longrightarrow [(\lambda y. e')/x]e}$$

small-step call-by-value

# CBN vs. CBV (Part 1)

*Determine an expression in the **untyped** lambda calculus that terminates using CBN evaluation but does not terminate using CBV evaluation*

$$\frac{\frac{e_1 \longrightarrow e'_1}{e_1 e_2 \longrightarrow e'_1 e_2} \quad \frac{e_2 \longrightarrow e'_2}{(\lambda x. e_1) e_2 \longrightarrow (\lambda x. e_1) e'_2}}{(\lambda x. e)(\lambda y. e') \longrightarrow [(\lambda y. e')/x]e}$$

small-step call-by-value

$$\frac{e_1 \longrightarrow e'_1}{e_1 e_2 \longrightarrow e'_1 e_2} \quad \frac{}{(\lambda x. e) e' \longrightarrow [e'/x]e}$$

small-step call-by-name

# CBN vs. CBV (Part 2)

*Let  $e^k$  be the untyped  $\lambda$ -calculus expression*

$$\frac{\text{k times}}{(\lambda x . x) \dots (\lambda x . x)}$$

*How many steps does it take to evaluate  $e^k$  using CBN and CBV? What is its value?*

*Using this expression, determine an expression that take 500 steps to evaluate using CBN and 104 steps to evaluate using CBV*