E-ISSN: 3139-0668
Volume: I, Issue: II

https://kalingauniversity.ac.in/International-Journal-of-Research-and-Studies

INTERNATIONAL JOURNAL OF RESEARCH AND STUDIES

Month/Year : April-June, 2026 | DOI: https://doi.org/10.5281/zenodo.23655844

RECEIVED 28.04.2025

ACCEPTED 06.05.2025

PUBLISHED 30-06-2026

IN SILICO-GUIDED DEVELOPMENT AND PHYSICOCHEMICAL CHARACTERIZATION OF CINNAMON EXTRACT-LOADED TRANSDERMAL PATCHES FOR POSTPRANDIAL HYPERGLYCEMIA MANAGEMENT: A COMPREHENSIVE REVIEW

Chinmay Kumar Pradhan - Research Scholar

Department of Pharmacy, Kalinga University, Raipur, Chhattisgarh

Khushboo Gupta - Assistant Professor

Faculty of Pharmacy, Kalinga University, Raipur, Chhattisgarh

ABSTRACT:

Postprandial hyperglycemia is a significant factor in the advancement of type 2 diabetes mellitus T2DM and associated vascular disorders. Cinnamon extracts have demonstrated notable antihyperglycemic effects by blocking carbohydrate digesting enzymes. However traditional oral delivery presents difficulties due to extensive hepatic first pass metabolism low bioavailability and brief biological half lives. This review outlines current findings on shifting from oral formulations to cinnamon loaded transdermal drug delivery systems TDDS. It evaluates the physicochemical characteristics of matrix type patches the application of Fourier transformed spectroscopy FTIR and reversed phase high performance liquid chromatography RP HPLC for quality control and the efficiency of multi target molecular docking against alpha glucosidase alpha amylase and matrix metalloproteinase 8 MMP 8. Additionally it addresses the integration of safety evaluations using tools such as ProTox to ensure the practical application of these herbal polymeric systems.

Keywords: *Postprandial Hyperglycemia, Cinnamon Extract, Transdermal Patches, Molecular Docking, RP-HPLC Validation, FTIR Characterization.*

1. Introduction

Diabetes mellitus often described as honey urine has developed from a historical clinical curiosity into a major global health concern. It is a chronic metabolic disorder that results in elevated blood glucose levels that persist. This can occur when pancreatic beta cells fail to produce sufficient insulin or when the body does not respond effectively to insulin. Postprandial hyperglycemia which refers to the rapid rise in blood glucose levels one to two hours after a meal is an early sign of diabetes and a major contributor to oxidative stress cardiovascular disease and microvascular damage. Managing these sudden glucose fluctuations is frequently more critical for preventing endothelial dysfunction than simply controlling prolonged high blood sugar levels.

Conventional pharmacological treatment often relies on oral antihyperglycemic agents and insulin therapy however these approaches face considerable limitations. Oral drugs such as Repaglinide are restricted by extremely short biological half lives approximately one hour and significant hepatic first pass metabolism leading to a peak and valley pattern in plasma concentrations and frequent gastrointestinal discomfort. Recent clinical studies suggest that the antihyperglycemic effectiveness of herbal powders is significantly reduced when administered in hard gelatin capsules instead of dissolved forms likely due to poor dissolution.

Cinnamon a spice derived from the Lauraceae family especially Cinnamomum zeylanicum and C. burmannii has been utilized in traditional medicine for over four thousand years. The primary bioactive component in its essential oil is trans cinnamaldehyde which constitutes sixty to ninety percent of the oil. It is highly effective in inhibiting the enzymes responsible for carbohydrate breakdown alpha amylase and alpha glucosidase. Cinnamaldehyde is highly potent but exhibits limited effectiveness when taken orally only about twenty percent and rapidly converts into cinnamic acid metabolites.

Transdermal drug delivery systems TDDS offer a non invasive controlled release approach to overcome these pharmacokinetic challenges. Cinnamaldehyde is a suitable candidate for TDDS due to its low molecular weight approximately 132.16 Daltons which facilitates passive diffusion through the skin. This thesis adopts an in silico guided approach to improve these formulations utilizing multi target molecular docking against alpha amylase alpha glucosidase and MMP 8 a target linked with diabetic vascular complications. This computational framework along with RP-HPLC validation and ProTox 3.0 safety assessments, guarantees the creation of a standardized, stable, and biologically safe herbal-polymer synergistic patch.

1.1: Importance of Postprandial Hyperglycemia in Diabetes

Elevated blood sugar levels after meals can be very harmful. They often cause endothelial dysfunction, which is worse than the effects of chronic high blood sugar. Post-meal blood sugar spikes significantly influence overall daily hyperglycemia and are better predictors of cardiovascular problems than fasting blood sugar levels. Controlling these spikes is complex. It involves considerations of carbohydrate intake, how quickly the stomach empties, and the rates of insulin secretion.

1.2: Cinnamon Extract as an Anti-Hyperglycemic Agent

Cinnamon, from the genus Cinnamon, has been used in traditional medicine for over 4,000 years to manage blood sugar. Its main antihyperglycemic action is blocking enzymes that break down carbohydrates, particularly pancreatic alpha-amylase and intestinal alpha-glucosidase. By slowing the conversion of complex carbohydrates into simple sugars, cinnamon extracts effectively reduce postprandial spikes. Moreover, cinnamon contains bioactive compounds, such as trans-cinnamaldehyde, which mimic insulin and boost glucose uptake through the GLUT4 pathway.

1.3: Transdermal Drug Delivery Systems for Herbal/Antidiabetic Therapy

The traditional oral form of cinnamon faces challenges. Its low oral absorption (around 20%) and short lifespan (about 1.7 hours) occur because it quickly converts to cinnamic acid in the liver. Additionally, taking it by mouth can irritate the stomach and cause changes in blood levels. Transdermal drug delivery systems (TDDS), especially matrix-type patches, provide a better alternative by avoiding the digestive system and liver. TDDS allows for controlled, steady release of the active ingredients, ensuring constant blood levels and improving patient compliance thanks to a painless, non-invasive method.




Fig1. Transdermal Drug Delivery System

1.4: Physicochemical Characterization of Herbal Transdermal Patches

Creating effective transdermal systems involves thorough testing of the polymer matrix for consistency and durability.

1. Physical Parameters: Testing includes checking weight uniformity thickness and folding endurance to ensure the patch can endure stress without breaking.
2. Chemical Stability: The surface pH should remain between 6.8 and 7.1 to limit skin irritation.
3. Moisture Dynamics: Lower moisture content is preferred for film stability and to prevent microbial growth.
4. Release Kinetics: Optimized formulations aim for a linear steady release to keep therapeutic levels constant for 24 hours.

1.5: FTIR RP-HPLC Docking and Safety Prediction in Phytopharmaceutical Research

Modern development of phytopharmaceuticals requires using analytical and computational tools.

- **FTIR Spectroscopy:** This method is vital for compatibility studies. It confirms that cinnamon extracts and polymer matrices (like HPMC or PVP) do not interact harmfully. Stable characteristic peaks for functional groups such as carbonyl and amine groups show that the bioactive is properly integrated into the polymer.

• RP HPLC Validation: Reversed phase HPLC is vital for standardized herbal evaluation. It offers the sensitivity required to quantify active constituents like cinnamaldehyde in complex patch formulations. Validation checks include linearity recovery rates 98 to 102 percent and precision ensuring uniformity and adherence with regulations.

• Multi target Molecular Docking: In silico simulations estimate how cinnamon constituents attach to key enzymes. In addition to alpha glucosidase and alpha amylase docking against MMP 8 is significant. This enzyme is associated with degradation of the extracellular matrix and vascular complications in diabetes.

• Safety Assessment: Tools such as ProTox II and ADMET screening for instance pkCSM allow researchers to predict the toxicity maximum tolerated dose and specific organ risks of herbal compounds before clinical trials.




Fig 2 Transdermal Patch release mechanism

2.0: Patent Landscape

The patent landscape shows a shift from diagnostic markers to better therapeutic delivery systems.

Table 1 Patent Related Comparison

Patent Number	Title	Year	Country/Jurisdiction	Key Invention	Relevance to Thesis
EP 2 325 652 B1	Method of Diagnosing Postprandial Hyperglycemia	2013	Europe	Diagnosis using hemoglobin with glycated lysine residues (GHbLys)	Establishes the clinical necessity for managing short-term PPHG spikes.

				as a sensitive short-term marker.	
US 10076128 B2	Method for Preventing or Reducing Hyperglycemia	2018	USA	Standardized cinnamon extract (Alpha-401) with defined polyphenol content and temperature-controlled extraction.	Provides parameters for standardized cinnamon extract integrity and bioactivity.
WO 2020/148741 A2	Therapeutic Transdermal Bioreactor or Trap Patch	2020	International	Microneedle-based patch containing functional molecules to convert or trap glucose.	Supports the shift toward advanced transdermal systems for managing insulin spikes.
EP 4 074 256 A1	Detection of Prolonged Postprandial Hyperglycemia	2022	Europe	Computer-aided method to predict postprandial peaks and adjust insulin delivery dynamically.	Highlights the translational value of predictive modeling in hyperglycemia management.
US 2025/0213635 A1	Method of Reducing Postprandial Glucose from a Meal	2025	USA	Nutritional solutions using plant extracts (mulberry) fiber and fatty acids to blunt glycemic response.	Demonstrates the ongoing innovation in multi-component herbal glycemic management.

3.0: Ongoing and Recent Research

Recent studies highlight the issues with traditional oral dosing and the need for standardized herbal analysis.

Table 2 ongoing Research outcomes

Author/Year	Study Type	Main Findings	Model/System Used	Relevance to Thesis
Moreira et al. (2024)	Crossover Clinical Trial	Antihyperglycemic effects of cinnamon powder are significantly weakened when administered in hard	T2DM Patients (Human)	Justifies the need for alternative delivery (transdermal) that avoids the limitations of gastric dissolution/bioavailability.

		gelatin capsules compared to dissolved powder.		
McMillan (2021)	In Silico & In Vitro	High-throughput virtual screening of 1070 herbal compounds; positive correlation found between Glide docking scores and enzyme inhibition constants .	alpha-amylase and alpha-glucosidase (Porcine/Yeast)	Validates the "In Silico-Guided" approach for selecting potent antidiabetic ligands.
Bernardo et al. (2022)	Randomized Controlled Trial	Aqueous cinnamon extract did not significantly lower glucose during an OGTT in T2DM patients but showed strong antioxidant capacity.	T2DM Patients (Human)	Highlights the species-specific and solvent-dependent variations in cinnamon efficacy.

4.0: Research Gap Analysis and Justification

The literature emphasizes a notable gap in research regarding the standardization and high resolution validation of herbal TDDS. While trials such as those by Moreira et al. (2024) demonstrate that encapsulation decreases cinnamon's effectiveness current pharmacological studies mainly concentrate on synthetic drugs or simple oral powders. There is a shortage of stability testing HPLC methods specifically designed for herbal transdermal patches to monitor the integrity of various bioactive components over time. In addition although the inhibition of alpha amylase and alpha glucosidase has been widely examined the role of MMP 8 as a target for diabetes related vascular health is insufficiently explored in the context of herbal patch delivery. This thesis aims to address these gaps by developing a standardized HPLC validated multi target guided transdermal system that enhances the delivery of cinnamon’s active components.

5.0: Conclusion

Postprandial hyperglycemia is a major therapeutic target for preventing the progression of type 2 diabetes and associated vascular complications. Cinnamon extract is a potent natural inhibitor of carbohydrate

digestion but its efficiency is restricted due to issues in oral delivery. The development of cinnamon loaded transdermal patches offers a promising strategy to maintain consistent blood levels while bypassing liver metabolism. By integrating FTIR compatibility studies validated RP HPLC analysis and multi target simulations against MMP 8 alpha glucosidase and alpha amylase this research establishes a strong foundation for standardized herbal therapies. Predictive safety evaluations further ensure that these innovative transdermal systems are both effective and safe for long term clinical use.

6.0: References

1. **Beejmohun V. et al. (2014).** Acute effect of Ceylon cinnamon extract on postprandial glycemia: alpha-amylase inhibition starch tolerance test in rats and randomized crossover clinical trial in healthy volunteers. *BMC Complementary and Alternative Medicine* 14:351.
2. **Shihabudeen H. M. S. et al. (2011).** Cinnamon extract inhibits alpha-glucosidase activity and dampens postprandial glucose excursion in diabetic rats. *Nutrition & Metabolism* 8:46.
3. **Moreira F. D. et al. (2024).** Suppression of the postprandial hyperglycemia in patients with type 2 diabetes by a raw medicinal herb powder is weakened when consumed in ordinary hard gelatin capsules: A randomized crossover clinical trial. *PLoS ONE* 19(10): e0311501.
4. **McMillan J. (2021).** In silico docking and ADMET studies on clinical targets for type 2 diabetes correlated to in vitro inhibition of pancreatic alpha-amylase and alpha-glucosidase by rutin caffeic acid p-coumaric acid and vanillin. *MSc Thesis* University of Pretoria.
5. **Rachid A. P. et al. (2022).** Effect of Aqueous Cinnamon Extract on the Postprandial Glycemia Levels in Patients with Type 2 Diabetes Mellitus: A Randomized Controlled Trial. *Nutrients* 14 1576.
6. **Bhujbal S. S. et al. (2011).** A novel herbal formulation in the management of diabetes [Transdermal patches of Momordica charantia]. *International Journal of Pharma Investigation* 1(4): 222-226.
7. **EP 2 325 652 B1 (2011).** Method of Diagnosing Postprandial Hyperglycemia. *European Patent Office*.
8. **EP 4 074 256 A1 (2022).** Computer-aided methods for detecting prolonged postprandial hyperglycemia and predicting postprandial glycemic peaks. *European Patent Office*.
9. **US 10076128 B2 (2018).** Method for Preventing or Reducing Hyperglycemia [Standardized Cinnamon Extract Alpha-401]. *United States Patent*.

10. **US 9700590 B2 (2017).** Methods and materials for reducing or eliminating risk factors associated with Syndrome X [Water-soluble cinnamon extract]. *United States Patent*.
11. **WO 2020/148741 A2 (2020).** Therapeutic transdermal bioreactor or trap patch for diabetes and other disorders. *World Intellectual Property Organization*.
12. **WO 2008/000063 A1 (2008).** Herbal product comprising cinnamon and bitter melon for treating diabetes. *World Intellectual Property Organization*.
13. **US 2011/0071765 A1 (2011).** Device and method for alleviating postprandial hyperglycemia [skin-securable insulin delivery]. *United States Patent Application*.
14. **US 2011/0275605 A1 (2011).** Postprandial hyperglycemia-improving agent [cycloartenol derivatives]. *United States Patent Application*.
15. **US 2025/0213635 A1 (2025).** Method of reducing postprandial glucose from a meal [mulberry leaf extract composition]. *United States Patent Application*.
16. **Review trial.docx.** In Silico-Guided Development and Physicochemical Characterization of Cinnamon Extract-Loaded Transdermal Patches. *Thesis/Chapter Draft Excerpts*.