E-ISSN: 3139-0668
Volume: I, Issue: II

https://kalingauniversity.ac.in/International-Journal-of-Research-and-Studies

INTERNATIONAL JOURNAL OF RESEARCH AND STUDIES

Month/Year : April-June, 2026 | DOI: https://doi.org/10.5281/zenodo.23595059

RECEIVED 16.04.2026

ACCEPTED 18.04.2026

PUBLISHED 30-06-2026

CHALLENGES AND EMERGING TREATMENT STRATEGIES FOR PEDIATRIC ASTHMA

Harshita Chanap - Research Scholar | harshitachanap25@gmail.com

Department of Pharmacy, Kalinga University, Raipur, Chhattisgarh

ABSTRACT:

Pediatric asthma is a long-term respiratory condition that lead to inflammation in the airways, Millions of the children affectated by the pediatric asthma in the worldwide. This illness can interfere with daily activities like school attendance, health and overall well-being. Challenges include under diagnosis, misinterpretation of symptoms, improper use of inhalers, poor adherence to treatment strategies, and continuous disclosure to environmental irritants such as pollution, allergens, tobacco and smoke. The complexity of this pediatric asthma arises from a combination of genetic, immune, and environmental factors like pollution and allergens. Conventional methods like inhaled corticosteroids, bronchodilators continue to play a crucial role, but many children still experience symptoms. Emerging therapies offer promise for improved control, including medication that trigger specific inflammatory pathways in asthma. Personalized medicine, which involves identifying the specific asthma subtype in each child, allows for tailored treatment approaches. Additional tools such as smart inhalers, digital monitoring systems, allergen immunotherapy, and enhanced drug delivery methods support better medication adherence, reduce flare-ups, and improve disease monitoring. This overview outlines the key obstacles in managing pediatric asthma and explores how new treatment options can address these challenges. It emphasizes the importance of combining medical advances with family education, environmental control, and preventive care to reduce asthma's impact and enhance the lifestyles that affected childrens.

Keywords: *Pediatric Asthma, Asthma Management, Airway Inflammation, Biologic Therapy, Environmental Triggers.*

Introduction

Asthma is a common long-term respiratory condition in children, posing significant public health challenges due to its rising incidence and enduring impact on physical, mental, and social well-being. Pediatric asthma is marked by ongoing airway inflammation, increased bronchial sensitivity, and fluctuating airflow obstruction. Common symptoms include wheezing, coughing, chest tightness, and shortness of breath, which often recur and are triggered by environmental factors, infections, allergens, or physical exertion, making management complex as children develop [1].

Globally, asthma affects millions, with prevalence rates approaching 10% in numerous areas. It is a primary cause of school absenteeism, hospitalizations, and diminished quality of life among pediatric populations. Beyond health issues, asthma contributes to considerable socio-economic costs, including frequent emergency visits, heightened healthcare expenditures, and progressive lung function impairment. In developing regions, challenges such as pollution, limited healthcare availability, and low caregiver awareness exacerbate these effects [2].

The pathophysiology of pediatric asthma involves multifaceted interactions between genetic susceptibility and environmental exposures, resulting in airway inflammation, edema, bronchoconstriction, and structural remodeling [3]. Various immune components, including eosinophils, mast cells, cytokines, and T-helper cells, play roles in these processes. Differences in these mechanisms give rise to distinct asthma phenotypes, each with unique clinical presentations and therapeutic responses [4].

Despite advancements in knowledge and treatment options, many children still suffer from poorly controlled symptoms and frequent exacerbations. This discrepancy underscores the complexity of managing pediatric asthma [5]. Diagnosing asthma in young children is particularly difficult due to challenges in performing reliable lung function tests like spirometry. Overlapping symptoms with other respiratory illnesses further complicate diagnosis. Early childhood asthma often manifests episodically with infection-related symptoms rather than persistent inflammation, making it harder to differentiate from transient wheezing [1].

Asthma severity ranges from mild intermittent to severe, treatment-resistant forms. Although severe asthma affects a smaller proportion of children, it accounts for a significant share of healthcare utilization and morbidity. This variability necessitates tailored treatment strategies, which may be hindered by limited resources and expertise [6].

Medication adherence presents a significant barrier, with many children inconsistently using inhaled corticosteroids due to concerns about side effects, improper inhaler technique, lack of supervision, and socio-economic factors [3]. Excessive reliance on short-acting bronchodilators for symptom relief without addressing underlying inflammation can lead to poorer outcomes [5].

Environmental and lifestyle influences further complicate asthma management. Exposure to allergens such as dust mites, pollen, and pet dander, alongside pollutants like tobacco smoke and urban air contaminants, can provoke asthma attacks. Factors such as climate change and urbanization have increased pollutant exposure, worsening symptoms. Indoor contributors like mold and inadequate ventilation also sustain airway inflammation.

Socio-economic inequalities significantly affect asthma outcomes. Children from low-income households face obstacles including restricted healthcare access, insufficient health education, and living conditions that elevate exposure to triggers. Cultural beliefs and limited asthma awareness among caregivers also impede effective management and adherence [7].

Pediatric asthma care is further limited by a scarcity of child-specific research. Many treatment recommendations are adapted from adult studies and may not fully account for children's unique physiological and developmental characteristics, including airway dimensions, immune responses, drug metabolism, and behavior [5].

International bodies such as GINA have established evidence-based guidelines promoting stepwise therapy, consistent monitoring, and patient education. Inhaled corticosteroids remain the primary treatment, supplemented by bronchodilators and other controller medications tailored to disease severity. Despite these protocols, many children continue to experience inadequate control, emphasizing the need for innovative and individualized therapeutic approaches.

Literature Review

Asthma is a chronic inflammatory disorder of the airways characterized by variable airflow obstruction and airway hyperresponsiveness. In children, it stands as a major non-infectious condition contributing significantly to global health burdens. Pediatric asthma differs from adult asthma in its pathophysiology, triggers, and therapeutic responses, posing unique challenges in clinical management [8].

1. Epidemiology and Global Burden

The prevalence of childhood asthma ranges between 5% and 15%, varying according to geographic location and environmental influences. Globally, asthma affects over 260 million individuals, resulting in approximately 400,000 deaths each year, with the majority occurring in low- and middle-income countries. In India, the incidence of pediatric asthma has almost doubled over the last twenty years, largely due to factors such as rapid urbanization, increased environmental pollution, and shifts in lifestyle patterns [9].

2. Etiology and Pathophysiology

Pediatric asthma develops from a multifaceted interaction between genetic predisposition and environmental factors. Viral infections in infancy, notably respiratory syncytial virus and rhinovirus, have been implicated in disrupting immune system regulation, thereby elevating the risk of childhood asthma onset. Genetic polymorphisms in cytokines such as IL-4, IL-13, and genes like ORMDL3 are associated with airway inflammation and increased airway responsiveness. Environmental exposures such as household smoke, dust mites, pet allergens, industrial pollutants, and poor indoor air quality contribute significantly to disease development. Chronic exposure leads to airway remodeling, characterized by subepithelial fibrosis, excessive mucus secretion, and smooth muscle hypertrophy, which together cause persistent airflow limitation despite treatment [10].

3. Clinical Presentation and Diagnosis

Children with asthma typically exhibit recurrent episodes of wheezing, coughing, chest tightness, and breathing difficulties, with symptom severity and frequency varying widely. Diagnosing asthma in young children is particularly complex due to symptom overlap with other respiratory illnesses like bronchiolitis and viral-induced wheezing.

NHLB(2020) guideline for accurate diagnosis emphasize

- The clinical examination of causes and symptoms
- Asthma or atopy in family members
- Tests for objective lung function, including peak expiratory flow or spirometry for older children
- Analysis of signs for eosinophilic inflammation, such as fractional exhaled nitric oxide (FeNO) [11]

4. Traditional Methods of Therapy (GINA 2024)

- It is recommended to take a slow and steady approach: The fundamental component for handling chronic care is inhaled corticosteroids (ICS).
- SABA, or short-acting beta-agonists, are used to treat acute symptoms.
- Leukotriene receptor antagonists (LTRAs): supplemental treatment (e.g., montelukast)
- For mid-to-high severity asthma, long-acting beta-agonists (LABAs) are used in conjunction with ICS.
- Oral corticosteroids should only be used in cases of severe exacerbations.
- Many children have uncontrolled asthma despite adherence, which highlights the need for innovative treatments [8].

Current Treatment Methods Used in Children

International clinical guidelines, like those created by GINA (the Global Initiative for Asthma), suggest a step-by-step treatment approach for pediatric asthma. This strategy's main goal is to control asthma while reducing the chance of exacerbations and adverse drug reactions. Inhaled corticosteroids (ICS), which function as strong anti-inflammatory drugs that lessen bronchial hyperresponsiveness, reduce airway inflammation, and prevent asthma flare-ups in children, are the mainstay of treatment [8].

ICS are frequently used in conjunction with bronchodilators in clinical practice, including long-duration the beta (2)-agonists (LABA) for long-term management of persistent manifestations and short-acting the beta (2)-agonists (SABA) to provide quick symptom relief [12].

When first-line treatments for asthma are insufficient to control symptoms, an additional pharmacological therapy are used in addition to inhaled corticosteroids and bronchodilators. Among these are leukotriene receptor antagonists (LTRAs), like montelukast are work by obstructing leukotriene-mediated inflammatory pathways to lessen airway inflammation. To improve bronchodilation during acute exacerbations, anticholinergic medications, especially ipratropium bromide, may also be used as supplemental therapy. Additionally, for quickly reduce airway inflammation and restore normal lung function during severe asthma attacks, systemic corticosteroids are often prescribed for brief periods of time [13]. These drugs' main objectives are to maintain long-term symptom control, avoid recurrent flare-ups, and enable kids to continue engaging in regular bodily exercise, growth, and general well-being.

For the patients who suffering from acute or treatment-defying respiratory issues, biologic therapies now represent a significant treatment option in recent years. Monoclonal antibodies are specifically target inflammatory mediators involved in the pathophysiology of asthma are used in these treatments. For instance, for pediatric cases of moderate-to-critical allergic bronchial disease, omalizumab is an immunoglobulin E (IgE) inhibitor, has been demonstrated to lessen allergic inflammation and the frequency of asthma flare-ups. In a similar vein, dupilumab suppresses the signaling routes for IL-4 and IL-13 linked to type-2 inflammation by blocking the interleukin-4 receptor (IL-4R), whereas mepolizumab targets interleukin-5 (IL-5) to reduce eosinophilic inflammation. Benralizumab, another biologic drug, depletes eosinophils, which are important in severe asthma, by targeting the IL-5 receptor [14]. These biologics considerably enhance respiratory performance, lessen chronic flare-ups, plus lessen the requirement for systemic corticosteroids in kids with severe asthma phenotypes, according to clinical research.

Challenges in Pediatric Asthma Management

Effectively managing pediatric asthma over the long term involves navigating multiple clinical, social, environmental, and therapeutic challenges. Despite established guidelines from organizations such as the World Health Organization and the Global Initiative for Asthma, many children still experience uncontrolled symptoms, frequent exacerbations, and diminished quality of life [8]. These difficulties arise from limitations within healthcare systems, diagnostic complexities, treatment delivery issues, and socio-environmental influences.

A key challenge is obtaining an accurate diagnosis, particularly in infants and preschool-aged children. Since objective lung function tests like spirometry are often impractical for this age group, clinicians rely on symptom patterns that may overlap with other respiratory conditions such as viral bronchiolitis or recurrent infections [15]. This reliance can lead to both overdiagnosis and delayed diagnosis, negatively impacting treatment effectiveness and disease progression [16].

The diverse nature of childhood asthma, encompassing various phenotypes such as allergic (eosinophilic), non-allergic, exercise-induced, and virus-induced asthma, adds further complexity. Standard treatments may not be equally effective across these phenotypes, as each responds differently to therapy [17]. The absence of dependable biomarkers for routine clinical use restricts the application of precision medicine approaches that tailor treatments to specific inflammatory pathways [18].

Treatment adherence remains a significant barrier to asthma control. Pediatric patients depend heavily on caregivers for medication administration, and factors such as complicated treatment regimens, fears about steroid side effects, limited disease awareness, and incorrect inhaler technique contribute to poor adherence [19]. Research shows that many children misuse inhaler devices, resulting in inadequate medication delivery and suboptimal disease management [20].

Environmental and lifestyle factors further complicate asthma management. Exposure to tobacco smoke, household irritants like dust mites, animal allergens, and climate-related triggers increase the frequency and severity of asthma attacks [21]. Children living in rapidly urbanizing areas, such as Mumbai, are especially vulnerable to high levels of particulate matter and vehicle emissions, which worsen respiratory symptoms and airway inflammation [22].

Access to advanced therapies, including biologic agents like anti-IgE and anti-interleukin monoclonal antibodies, is limited. While these treatments show promise for severe pediatric asthma, their high cost, need for specialized healthcare infrastructure, and limited availability in resource-poor settings restrict their widespread use [23]. This disparity results in varied treatment outcomes between developed and developing regions.

Children with poorly controlled asthma often face emotional challenges such as anxiety and sleep disturbances. The condition can limit participation in play and social activities. Frequent hospital visits and school absences may adversely affect academic performance and social development. Caregivers also experience stress, financial burdens, and difficulties balancing work and care giving responsibilities [24][25].

Emerging Treatment Strategies for Pediatric Asthma

Recent advancements have improved pediatric asthma care considerably. New therapies targeting specific disease mechanisms, enhanced drug delivery systems, and personalized care plans are contributing to better outcomes. While inhaled corticosteroids and bronchodilators remain foundational, novel approaches aim to address underlying causes, improve adherence, and maintain asthma control. Global health organizations continue to update guidelines to integrate these innovations [8].

One notable advancement is the introduction of biologic therapies that target precise immune pathways. Medications such as omalizumab, mepolizumab, and dupilumab have shown effectiveness in

children with severe allergic or eosinophilic asthma [26]. These agents reduce airway inflammation, decrease exacerbations, improve lung function, and reduce the need for systemic steroids. Although their use in pediatric populations is still expanding, biologics represent a significant step toward personalized asthma management.

Personalized care is increasingly emphasized, tailoring treatment to each child's specific asthma phenotype. Progress in genomics, proteomics, and biomarker research facilitates better identification of inflammatory drivers, enabling clinicians to select the most suitable therapies. Biomarkers such as exhaled nitric oxide, blood eosinophil counts, and allergen-specific IgE are increasingly used to guide treatment decisions and monitor responses [18].

Technological innovations are transforming asthma management. Smart inhalers equipped with sensors can monitor medication use, provide reminders, offer guidance on inhaler technique, and share data with healthcare providers via smartphone apps [27]. This technology supports improved adherence, especially among adolescents, and encourages active disease management.

Immunotherapy is another promising option for children with allergic asthma. By gradually exposing the immune system to allergens through subcutaneous injections or sublingual tablets/drops, immunotherapy can reduce allergic symptoms, decrease medication requirements, and potentially alter disease progression [28].

Improvements in drug delivery devices have enhanced treatment effectiveness. Breath-actuated inhalers and spacer devices improve medication deposition in the lungs and reduce systemic side effects [29]. Portable nebulizers have become more user-friendly, benefiting younger children who find standard inhalers challenging.

Environmental control and trigger avoidance are increasingly recognized as essential components of asthma care. Strategies such as air purifiers, allergen-proof bedding, smoking cessation support, and pollution reduction are particularly important in highly polluted urban centers like Delhi to lessen the asthma burden [30]. Early detection and prevention strategies are under active investigation. Identifying children at high risk and initiating early interventions may slow disease progression and improve long-term outcomes. Potential preventive measures include nutritional optimization, vitamin D supplementation, and microbiome modulation [31].

Finally, multidisciplinary team-based care involving physicians, nurses, therapists, and families enhances asthma management. Coordinated efforts improve education, treatment adherence, and monitoring. School-based programs and community awareness initiatives also contribute to reducing hospitalizations and improving quality of life [32].

Prevention of Pediatric Asthma

Preventing asthma in children is crucial for maintaining healthy lungs and reducing both the occurrence and severity of asthma attacks. Limiting exposure to environmental triggers such as air pollution, cigarette smoke, dust, and pet dander is essential. Supporting healthy growth and development from an early age also plays a vital role. Health organizations like the World Health Organization (WHO) recommend these early interventions to protect children and prevent asthma from developing or worsening [30].

One effective prevention strategy is minimizing exposure to asthma triggers in the home and environment. For example, exposure to tobacco smoke during pregnancy or early childhood significantly increases the risk of developing asthma. Common indoor allergens such as dust mites, mold, pet hair, and cockroach droppings can irritate sensitive airways and provoke asthma attacks. Maintaining good indoor air quality by keeping the air fresh, using allergen-proof bedding covers, controlling humidity, and avoiding tobacco smoke can substantially reduce asthma symptoms in children [8].

Early care of infants is another important factor. Breastfeeding for the first six months helps protect babies from lung infections and allergies, including asthma. Breast milk contains components that strengthen the immune system and lower the likelihood of developing allergies. Additionally, adequate maternal nutrition during pregnancy, including sufficient vitamins and antioxidants, can reduce the child's risk of asthma or allergies later in life [33].

Vaccinations and controlling infections also contribute to asthma prevention. Respiratory infections, especially viral infections like RSV, often cause wheezing and may lead to asthma in young children. Routine vaccinations and good hygiene practices, such as frequent handwashing, help reduce the risk of these infections and protect lung health, thereby decreasing asthma flare-ups triggered by infections [15].

Reducing outdoor air pollution is another critical preventive measure. Pollutants such as fine particulate matter (PM2.5), nitrogen dioxide (NO_2), and ozone in urban environments are linked to increased asthma incidence and severity in children. Long-term exposure to polluted air raises the risk of developing

asthma and worsens respiratory problems. Implementing policies to reduce emissions from vehicles and factories and improving city air quality can help lower the number of children affected by asthma [21].

Emerging prevention methods include allergen-specific immunotherapy (AIT) and personalized early interventions. AIT gradually exposes children to small amounts of allergens to help their immune systems build tolerance, reducing allergy and asthma symptoms over time. Advances in genetic testing and biomarker identification may allow healthcare providers to detect children at high risk for asthma early and begin preventive treatments before the disease progresses, potentially halting its development [28].

Conclusion

Millions of children worldwide are affected by paediatric asthma, significantly impacting their health, school attendance, psychological well-being, and use of healthcare services. According to the World Health Organization (WHO), asthma is one of the most common chronic non-communicable diseases among young people globally, with its prevalence rising notably in developing countries due to rapid urbanization, environmental pollution, and lifestyle changes. This growing burden underscores the urgent need for comprehensive, equitable, and sustainable management strategies tailored specifically for children.

One of the main challenges in managing paediatric asthma is its heterogeneity. The Global Initiative for Asthma (GINA, 2024) emphasizes that childhood asthma comprises various phenotypes and endotypes, each characterized by distinct inflammatory pathways and clinical features rather than a single uniform disease. This complexity is especially pronounced in preschool children, where objective lung function tests are limited, complicating diagnosis and treatment. Advances in understanding the differences between Type 2 (T2) and non-Type 2 airway inflammations have shifted treatment towards phenotype-specific approaches rather than generic therapies.

Poor adherence to inhaled corticosteroids (ICS) remains a persistent obstacle. Studies indicate that incorrect inhaler use, gaps in caregiver education, concerns about steroid side effects, and socioeconomic factors contribute to suboptimal asthma control [12]. To reduce exacerbations, updated GINA guidelines highlight the importance of ICS-containing regimens, either as regular low-dose therapy or as needed (PRN) ICS-formoterol. This reflects a shift toward early anti-inflammatory treatment to prevent long-term airway remodeling and improve outcomes.

Social and environmental factors further complicate asthma management in children. Exposure to indoor allergens, tobacco smoke, ambient air pollution, and climate change-related influences significantly increase the frequency and severity of exacerbations. Research published in The Lancet Respiratory Medicine links air pollution exposure to poorer respiratory health in urban youth. Addressing these determinants requires multidisciplinary strategies beyond medication, including community education, policy interventions, and public health efforts.

The treatment landscape for severe paediatric asthma has evolved considerably with new therapeutic options. Biologic agents targeting specific inflammatory pathways have shown promising results in reducing exacerbations and improving lung function. For example, omalizumab effectively treats allergic asthma by neutralizing IgE and reducing hypersensitivity reactions. Similarly, mepolizumab and dupilumab offer hope for children with severe eosinophilic or type 2-high asthma by targeting the IL-4/IL-13 pathway and eosinophilic inflammation, respectively. These biologics mark a significant step forward in precision medicine for paediatric asthma.

In addition to biologics, digital health technologies are transforming asthma monitoring and adherence. Tools such as telemedicine platforms, smart inhalers, and mobile health applications have improved medication compliance and symptom tracking. The COVID-19 pandemic accelerated telehealth adoption, demonstrating that remote consultations can effectively support asthma care without compromising quality. These innovations are particularly valuable in rural or resource-limited settings where specialist access is scarce.

REFERENCE

1. Miculinic A, et al. Modern difficulties in childhood asthma. Children (Basel), 2024.
2. Elsaid A, Abdelwahab A. Pediatric asthma care: an Egyptian perspective. J Pediatr Neonatal Care, 2024.
3. Jones H, Lawton A, Gupta A. Pediatric asthma flare-ups—difficulties and prospects. Indian Journal of Pediatrics, 2022.
4. Pharmacotherapeutic challenges in pediatric asthma management. Pharmaceutics, 2022.
5. Szefler SJ, Chipps B. Issues in treating asthma among youth and adolescents. Ann Allergy Asthma Immunol, 2018.
6. Bozzetto S, et al. Acute asthma in childhood: diagnostic and therapeutic hurdles. Curr Opin Pulm Med, 2015.
7. BMC Pediatrics study on barriers in childhood asthma care, 2025.
8. GINA (Global Initiative for Asthma). (2024). Worldwide Plan for Asthma Control and Prophylaxis.
9. Aggarwal, A. N., et al. (2022). Incidence and determinants of bronchial disease in Indian children: A multicenter population-based study. Indian Pediatrics, 59(5), 401–408.
10. Ober, C., & Yao, T. C. (2011). The genetics of asthma and allergic disease: A 21st-century perspective. Immunological Reviews, 242(1), 10–30.
11. National Heart, Lung, and Blood Institute (NHLBI). (2020). 2020 Updated Perspectives on Asthma Protocols. JACI, 146(6), 1217–1270.
12. Reddel HK, et al. Global Strategy for Asthma Report. European Respiratory Journal, 2022.
13. National Asthma Education and Prevention Program. Expert Panel Report: Protocols for Assessing and Managing Asthma.
14. Bacharier LB, et al. Biological treatments for acute pediatric bronchial disease. Journal of Allergy and Clinical Immunology, 2021.
15. Martinez FD. Early-life origins of long-term obstructive lung ailment. N Engl J Med, 2016;375:871–878.
16. Bush A, Fleming L. Is asthma overdiagnosed? Arch Dis Child. 2016;101:688–689.

17. Wenzel, S. (2012). Asthma phenotypes and personalized treatment. NEJM.
18. Agache, I., & Akdis, C. (2019). Precision medicine in asthma. Allergy.
19. Bender BG. Overcoming medication neglect in youth respiratory care. Allergy Clin Immunol Pract, 2016;4(5):849–855.
20. Price D et al. Inhalation device proficiency: Frequent mistakes and medical results. Respir Med, 2013;107:37–46.
21. Asher MI et al. Universal patterns in youth bronchial conditions. Lancet, 2020;396:1225–1236.
22. HEI (Health Effects Institute). State of Global Air Report. 2022.
23. Bush A, Saglani S. Controlling acute childhood respiratory disease. Lancet Respir Med, 2017;5(1):73–85.
24. Moonie S et al. The correlation between missed classes and respiratory severity. J Sch Health, 2008;78(3):140–148.
25. Pawankar R et al. Hypersensitivity ailments and bronchial health: a widespread societal issue. WAO Journal, 2013;6:1–17.
26. McGregor MC et al. The function of biological therapies in bronchial care. Lancet Respir Med, 2019;7:699–711.
27. Chan, A. H., et al. (2020). Digital inhalers enhance adherence and patient-reported outcomes. Pediatric Pulmonology.
28. Dhami S et al. Antigen-specific desensitization for allergic bronchial issues. Allergy, 2017;72:1825–1848.
29. Usmani OS. Selecting the appropriate inhalation tool for bronchial conditions. Respir Med, 2019;152:64–71.
30. WHO (World Health Organization). International tracking and management of long-term lung ailments. WHO Press, 2023.
31. Martineau AR et al. Cholecalciferol augmentation to avoid bronchial flare-ups. Lancet Respir Med, 2017;5:881–890.
32. Global Asthma Network. The Universal Respiratory Status Update, 2022.

33. Lodge CJ, et al. Nursing infants and bronchial risk: A comprehensive meta-analysis. Acta Paediatrica, 2015.

34. Busse W, et al. (2011). Anti-IgE therapy in youth with hypersensitive bronchial disease. NEJM.