Project 1 FYS3150

Anna Stray Rongve

August 30, 2019

Abstract

Introduction

Theory and technicalites

Conclusion and perspectives

Project 1 a)

For i=0 and i=n the boundary conditions gives us v(0)=v(1)=0. For i=1

$$-\frac{v_2 + v_0 - 2v_1}{h^2} = f_1$$

For i=2

$$-\frac{v_3 + v_1 - 2v_2}{h^2} = f_2$$

For i = n - 1

$$-\frac{v_n + v_{n-2} - 2v_{n-1}}{h^2} = f_{n-1}$$

If you multiply both sides by h^2

$$-v_2 + v_0 - 2v_1 = h^2 \cdot f_1$$

$$-v_3 + v_1 - 2v_2 = h^2 \cdot f_2$$

$$-v_n + v_{n-2} - 2v_{n-1} = h^2 \cdot f_{n-1}$$

Which you can rewrite as a linear set of equations where

$$A = \begin{bmatrix} 2 & -1 & 0 & & & 0 \\ -1 & 2 & -1 & 0 & & & \\ 0 & -1 & 2 & -1 & 0 & & \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & & & -1 & 2 & -1 \\ 0 & & & 0 & -1 & 2 \end{bmatrix}$$

$$\hat{v} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ \vdots \\ v_{n-1} \end{bmatrix}$$

and
$$\tilde{b_i} = \begin{bmatrix} \tilde{b}_1 \\ \tilde{b}_2 \\ \tilde{b}_3 \\ \vdots \\ \tilde{b}_{n-1} \end{bmatrix}$$

Where $\tilde{b_i} = h^2 \cdot f_i$

Project 1 b)

Appendix

Bibliography