

1 Theory

1.1 The problem

1.2 2×2 lattice, analytical expressions

To get started we will find the analytical expression for the partition function and the corresponding expectation values for the energy E , the mean absolute value of the magnetic moment $|M|$ (which we will refer to as magnetization), the specific heat C_V and the susceptibility χ as function of T using periodic boundary conditions. These calculations will serve as benchmarks for our next steps.

Partition function

The partition function in the canonical ensemble is defined as:

$$Z = \sum_{i=1}^N e^{-\beta E_i}$$

Where $\beta = \frac{1}{k_B T}$ and E_i is the energy of the system in the microstate i and N is the respective microstate.

We therefore have to find E_i which is defined as:

$$E_i = -J \sum_{\langle kl \rangle}^N s_k s_l$$

Where $\langle kl \rangle$ indicates that we sum only over the nearest neighbors and J is a constant for the bonding strenght. For our two dimensional system the equation reads:

$$E_{i,2D} = -J \sum_i^N \sum_j^N (s_{i,j} s_{i,j+1} + s_{i,j} s_{i+1,j})$$

Four our two-spin-state system with two dimensions we get the following table:

Number of spins up	Degeneracy	Energy	Magnetization
4	1	-8J	4
3	4	0	2
2	4	0	0
2	2	8J	0
1	4	0	-2
0	1	-8J	-4

Table 1: Number of spins up, degeneracy, energy and magnetization of the two-dimensional benchmark scenario.

Where the magnetization is found by subtracting the number of spin downs from the number of spin up.

1.3 Ising model

The Ising model is applied for the study of phase transitions at finite temperatures for magnetic systems. Energy is expressed as:

$$E = -J \sum_{\langle kl \rangle}^N s_k s_l \quad s_k = \pm 1 \quad (1)$$

N is the number of spins and J is a constant expressing the interaction between neighboring spins. The sum is over the nearest neighbours only, indicated by $\langle kl \rangle$ in the above equation. For $J < 0$ it is energetically favorable for neighboring spins to align. Leading to, at low temperatures, T, spontaneous magnetisation.

A probability distribution is needed in order to calculate the mean energy $\langle E \rangle$ and magnetization $\langle M \rangle$ at a given temperature. The distribution is given by:

$$P_i(\beta) = \frac{1}{Z} \exp(-\beta E_i) \quad (2)$$

where M is all the microstates, P_i is the probability of having the system in a state/configuration i.