

Problem 1

%%

clear, close all

%% Finding values a,b,c using system of linear equations

x = [-4, 0, 4];

y = [30, 2, 6];

A = [x.^2, x', ones(3, 1)];

d = y';

coefficients = A \ d;

a = coefficients(1);

b = coefficients(2);

c = coefficients(3);

fprintf('Coefficients [a, b, c]: %f, %f, %f\n', a, b, c);

%%

d = [30;2;6];

%% Project Matrix

P = A * inv(A' * A) * A';

disp("The projection matrix P:");

disp(P)

```
%% Predicted
```

```
predicted = P * d;
```

```
disp("The predicted values are:");
```

```
disp(predicted)
```

```
%% Error of vector e
```

```
e = d - predicted;
```

```
disp("The error vector e is:");
```

```
disp(e)
```

```
%% Newton's Method
```

```
% Initial iteration
```

```
A = 1;
```

```
B = 2;
```

```
C = 3;
```

```
% taylor series
```

```
f = @(A, B, C) sum((y - A*(x - B).*(x - C)).^2);
```

```
f_A = @(A, B, C) -2*sum((x - B).*(x - C).*(y - A*(x - B).*(x - C)));
```

```
f_B = @(A, B, C) 2*A*sum((x - B).*(y - A*(x - B).*(x - C))) + 2*A*sum((x - C).*(y - A*(x - B).*(x - C)));
```

```
f_C = @(A, B, C) 2*A*sum((x - B).*(y - A*(x - B).*(x - C))) + 2*A*sum((x - B).*(y - A*(x - B).*(x - C)));
```

```
tol = 1e-6;
```

```
iter = 100000;
```

```
%newton's method
```

```

for i = 1:iter
    df = [f_A(A, B, C), f_B(A, B, C), f_C(A, B, C)];
    delta = df \ -f(A, B, C);
    A = A + delta(1);
    B = B + delta(2);
    C = C + delta(3);
    if max(abs(delta)) < tol
        break;
    end
end

```

```

fprintf('Estimated values:\n');
fprintf('A = %f\n', A);
fprintf('B = %f\n', B);
fprintf('C = %f\n', C);

```

%% Plotting

% Importing initial values

```

A = 1;
B = 2;
C = 3;

```

```

con_B = [];
con_C = [];

```

% Newton's method

```

for i = 1:iter
    df = [f_A(A, B, C) f_B(A, B, C) f_C(A, B, C)]; %Jacobian
    delta = df \ -f(A, B, C); %solve
    A = A + delta(1);
    B = B + delta(2);
    C = C + delta(3);

```

```

con_B(end+1) = B;
con_C(end+1) = C;

if max(abs(delta)) < tol
    break;
end
end

% Plotting convergence in the B,C plane
figure(1)
plot(con_B, con_C, '-', 'LineWidth', 3);
xlabel('B');
ylabel('C');
title('Convergence of Newton's Method in B,C Plane');
grid on;
set(gca, 'FontSize', 20)

%%

```

What happens if you choose $A = 0$ as the initial iterate? Why can you rule out the value of $A = 0$ as a possible value?

- I think you cannot have $A=0$. My code gives me errors when doing the calculations. Yes, we can rule out $A=0$.

Problem 2

```

%%

clear, close all

%% Download ONI Index

```

```
clear, close all
```

```
%%
```

```
url = 'https://psl.noaa.gov/data/correlation/oni.data';
```

```
ONI_table = readtable(url, 'HeaderLines', 0, 'ReadVariableNames', true, 'FileType', 'text');
```

```
ONI_table = ONI_table(1:74, :);
```

```
%%
```

```
index = 1;
```

```
year = [];
```

```
%%
```

```
for i = 1:74
```

```
    %loop per time
```

```
    year_num = ONI_table.x1950(i);
```

```
    year = [year, repmat(year_num, 1, 12)];
```

```
    for b = 2: 13
```

```
        ONI(index) = ONI_table{i, [b]};
```

```
        month(index) = b-1;
```

```
        dayssince(index) = daysact(datetime(year_num,b-1,1));
```

```
index = index + 1;
```

```
end
```

```
end
```

```
%%
```

```
day = ones(888,1);
```

```
dayssince = dayssince';
```

```
month = month';
```

```
year = year';
```

```
ONI = ONI';
```

```
time = table(day, month, year, dayssince, 'VariableNames', {'day', 'month', 'year', 'Days Since  
Beginning'});
```

```
clear year month dayssince day url year_num ONI_table i b index
```

```
%%
```

```
% $d_i = \mu + (t_i - t)m.$ 
```

```
time2 = 1:888;
```

```
time2 = time2';
```

```
t = ONI; % ONI data
```

```
%%
```

```
trend_model = @(params, t) params(1) + (t - mean(t)) * params(2);
```

```
% Initial guess for parameters
```

```
initial_guess = [0, 0]; % [mu, m]
```

```
% Fit the trend model using lsqcurvefit
```

```
param = lsqcurvefit(@(params, t) trend_model(params, t), initial_guess, time2, t);
```

```
% Extract parameters
```

```
mu = param(1);
```

```
m = param(2);
```

```
% Calculate the trend line
```

```
projected_trend = trend_model(param, time2);
```

```
%%
```

```
time3 = time("Days Since Beginning");
```

```
figure(1)
```

```
plot(time3, t, 'k', 'LineWidth', 2);
```

```
hold on
```

```
plot(time3, projected_trend, 'r', 'LineWidth', 2);
```

```
xlim([datenum('January 1, 1950') datenum('dec 25, 2020')])
```

```
datetick('x', 'keeplimits')
```

```
set(gca, 'FontSize', 16);
```

```
xlabel('Time');
```

```
ylabel('ONI Index');
```

```
title('ONI Data with Projected Trend Model');
```

```

legend('ONI Data', 'Projected Trend Model');
grid on;
hold off;

%%

% Calculate residuals
est = ONI - projected_trend; %actual data minus trend

std_est = std(est);

%% Printing

% Output
disp(['Estimated value of mu: ' num2str(mu)]);
disp(['Estimated value of m: ' num2str(m)]);
disp(['Standard deviation of the components of the e vector: ' num2str(std_est)]);

%%

```

Problem 3

```

%%

clear, close all

%% drafting

% 16 different combination of co2 and h2o

%d = [d1; d2; d3; d4; d5; d6; d7; d8; d9; d10; d11; d12; d13; d14; d15; d16]; %data

```


% A matrix with all possible combinations

A = ones(16, 1);

CO2_labels = [0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1]; % CO2

H2O_labels = [0, 0, 1, 1, 0, 0, 1, 1, 0, 0, 1, 1, 0, 0, 1, 1]; % H2O

%model - $d_i = \mu + \beta_1 \Delta \text{CO2}_i + \beta_2 \Delta \text{H2O}_i + \beta_3 \Delta \text{CO2}_i \Delta \text{H2O}_i + e_i$ (error unknown)

A = [A, CO2_labels.', H2O_labels.', CO2_labels.' * H2O_labels.'];

%B = [mu; B1; B2; B3];

% Calculate $A^T A$

ATA = A.' * A;

% Calculate $A^T d$

%ATd = A.' * d;

%d= AB

%%

A =

1	0	0	0
1	1	0	0
1	0	1	0
1	1	1	1
1	0	0	0
1	1	0	0
1	0	1	0
1	1	1	1
1	0	0	0
1	1	0	0
1	0	1	0
1	1	1	1
1	0	0	0

1	1	0	0
1	0	1	0
1	1	1	1

ATA =

16	8	8	4
8	8	4	4
8	4	8	4
4	4	4	4