

You must justify your answers to receive full credit.

1. Let $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ be a basis of \mathbb{R}^3 , and suppose $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear transformation with

$$T(\mathbf{v}_1) = \mathbf{v}_2, \quad T(\mathbf{v}_2) = \mathbf{v}_1, \quad T(\mathbf{v}_3) = \mathbf{v}_1 + \mathbf{v}_3.$$

- a) Show that $\mathbf{v}_1 + \mathbf{v}_2$ is an eigenvector of T , and find its corresponding eigenvalue.
- b) Find the matrix for T relative to \mathcal{B} .

Now let

$$\mathbf{v}_1 = \begin{bmatrix} 0 \\ 5 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 0 \\ -2 \\ 0 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} -1 \\ -5 \\ 2 \end{bmatrix}.$$

- c) Find the change-of-coordinates matrix from the standard basis in \mathbb{R}^3 to \mathcal{B} .
- d) Find the standard matrix for T .

2. Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ and $\mathcal{F} = \{\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3\}$ be bases for a vector space V , and suppose

$$\mathbf{f}_1 = 2\mathbf{b}_1 - \mathbf{b}_2 + \mathbf{b}_3, \quad \mathbf{f}_2 = 3\mathbf{b}_2 + \mathbf{b}_3, \quad \mathbf{f}_3 = -3\mathbf{b}_1 + 2\mathbf{b}_3$$

- a) What is the dimension of V ?
- b) Find the change-of-coordinates matrix from \mathcal{F} to \mathcal{B} .
- c) Now suppose $V = \mathbb{P}_2$, the set of polynomials of degree at most 2, and \mathcal{B} is the standard

basis $\{1, t, t^2\}$. If $[\mathbf{p}]_{\mathcal{F}} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$, find \mathbf{p} .

3. (to be announced later)

4. (to be announced later)

5. Let A be a 2×2 matrix.

- a) Explain why there is a polynomial p of degree at most 4 (i.e. $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$) such that $p(A) = 0$. (Hint: think about linear independence of the set $\{I, A, A^2, \dots\}$ in the vector space of 2×2 matrices.)
- b) **Optional:** Show that, if $a_0 \neq 0$, then A is invertible and A^{-1} is a polynomial in A .

6. State whether each of the following statements is (always) true or (sometimes) false. If it is true, give a brief justification; if it is false, give a counterexample with an explanation.
- a) If V is a 6-dimensional vector space, then any set of 6 vectors in V is a basis for V .
 - b) If A is 4×7 matrix and $\text{rank} A = 4$, then $\text{Col} A = \mathbb{R}^4$.
 - c) If A is 4×7 matrix and $\text{rank} A = 4$, then $\text{Nul} A = \mathbb{R}^3$.
 - d) The sum of the dimensions of the row space and the null space of A equals the number of rows in A .
 - e) If λ is an eigenvalue of A , then $\lambda + 2$ is an eigenvalue of $A + 2I$.
 - f) Let $V = \mathbb{P}_3$ be the set of polynomials of degree at most 3. Then \mathbb{R}^3 and \mathbb{P}_3 are isomorphic vector spaces.

— END —