

In this week's notes, we are interested in finding the *local maxima* and *local minima* of a multivariate function f , i.e. the points (a_1, \dots, a_n) such that

$$\begin{aligned} f(a_1, \dots, a_n) &\geq f(x_1, \dots, x_n) \\ f(a_1, \dots, a_n) &\leq f(x_1, \dots, x_n) \end{aligned} \quad \text{for all } (x_1, \dots, x_n) \text{ close to } (a_1, \dots, a_n).$$


If $\nabla f(a_1, \dots, a_n) \neq \mathbf{0}$, then f is increasing in the direction $\nabla f(a_1, \dots, a_n)$ and decreasing in the direction $-\nabla f(a_1, \dots, a_n)$, so (a_1, \dots, a_n) cannot be a local maximum or minimum. So a local maximum or minimum must be a critical point.

Definition: A point (a_1, \dots, a_n) is a *critical point* of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ if $\nabla f(a_1, \dots, a_n) = \mathbf{0}$, i.e. if all its partial derivatives are 0.


But not every critical point is a local maximum or minimum as we will see.

§13.1: Classifying Critical Points


Recall that a critical point of a single-variable function f is where the derivative f' is zero. A standard way to determine whether it is a local maximum, a local minimum, or neither, is the **second derivative test**:



if $f''(a) > 0$ then a is a local minimum




if $f''(a) < 0$ then a is a local maximum




if $f''(a) = 0$ then we need to investigate further


The reason is clear from considering the change in the slope of the graph, but because graphs of multivariate functions are hard to visualise, we give a different justification on the next page.



if $f''(a) > 0$ then a is a local minimum




if $f''(a) < 0$ then a is a local maximum




if $f''(a) = 0$ then we need to investigate further

The second-order Taylor polynomial of f at a is


$$f(a + h) \approx f(a) + f'(a)h + \frac{f''(a)}{2!}h^2$$



if $f''(a) > 0$ then a is a local minimum



if $f''(a) < 0$ then a is a local maximum



if $f''(a) = 0$ then we need to investigate further

The second-order Taylor polynomial of f at a is

is 0 if a is a critical point

$$f(a+h) \approx f(a) + \underbrace{f'(a)}_{\text{circled in blue}}h + \frac{f''(a)}{2!} \underbrace{h^2}_{\text{circled in red}} \text{ is positive if } h \neq 0, \text{ i.e. } x \neq a$$

$$= f(a) + \frac{f''(a)}{2!}h^2 \quad \begin{cases} > f(a) & \text{if } f''(a) > 0 \text{ and } h \neq 0 \\ < f(a) & \text{if } f''(a) < 0 \text{ and } h \neq 0 \end{cases}$$

Here is a simplified example of how to use second order Taylor polynomials to classify critical points of multivariate functions.

Example: Find and classify the critical points of $f(x, y) = y^2 - x^3 + x$.

Now we develop a multivariate second derivative test by copying the previous example's argument in general.

The second-order Taylor polynomial of f about (a, b) is

$$f(a+h, b+k) \approx f(a, b) + f_x(a, b)h + f_y(a, b)k + \frac{f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2}{2!},$$

Now we develop a multivariate second derivative test by copying the previous example's argument in general.

The second-order Taylor polynomial of f about (a, b) is

$$f(a+h, b+k) \approx f(a, b) + f_x(a, b)h + f_y(a, b)k + \frac{f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2}{2!},$$

is 0 if (a, b) is a critical point

we need the “sign” of the numerator

Now we develop a multivariate second derivative test by copying the previous example's argument in general.

The second-order Taylor polynomial of f about (a, b) is

$$f(a+h, b+k) \approx f(a, b) + \underbrace{f_x(a, b)h + f_y(a, b)k}_{\text{is 0 if } (a, b) \text{ is a critical point}} + \frac{f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2}{2!},$$

is 0 if (a, b) is a critical point

we need the “sign” of the numerator

Definition: A function $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ is a *quadratic form* if it is homogeneous of degree two i.e. a linear combination of $x_i x_j$. A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

A quadratic form can be at most one of the three types. But it is possible to be none of the three types, e.g. $Q(h, k) = h^2$. (see later)

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A \left(h + \frac{B}{A}k \right)^2 + \frac{AC - B^2}{A}k^2$$

So $Q(x)$ is positive definite

if A and $\frac{AC - B^2}{A}$ are both positive, i.e. $A > 0$ and $AC - B^2 > 0$;

$Q(x)$ is negative definite

if A and $\frac{AC - B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$;

$Q(x)$ is indefinite if A and $\frac{AC - B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$, and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$; \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A \left(h + \frac{B}{A}k \right)^2 + \frac{AC - B^2}{A}k^2$$

$$\det \begin{pmatrix} A & B \\ B & C \end{pmatrix}$$

So $Q(x)$ is positive definite

if A and $\frac{AC - B^2}{A}$ are both positive, i.e. $A > 0$ and $\boxed{AC - B^2} > 0$;

$Q(x)$ is negative definite

if A and $\frac{AC - B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$;

$Q(x)$ is indefinite if A and $\frac{AC - B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

To phrase this in a way that will extend to functions of more than 2 variables, we consider a matrix containing the second-order partial derivatives:

Definition: The *Hessian matrix* $\mathcal{H}(\mathbf{a})$ of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at a point \mathbf{a} in \mathbb{R}^n is the $n \times n$ matrix with $\frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{a})$ in row i and column j .

Example: For a 2-variable function f , the Hessian matrix is $\mathcal{H}(a, b) = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix}$.

Let $D_i(\mathbf{a})$ denote the determinant of the $i \times i$ matrix containing only the first i rows and the first i columns of $\mathcal{H}(\mathbf{a})$.

Theorem: Second Derivative Test for 2-variable functions: Let (a, b) be a critical point of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$.

If $D_1(a, b), D_2(a, b) > 0$, then $\mathcal{H}(a, b)$ is positive definite and (a, b) is a **minimum**.

If $D_1(a, b), D_2(a, b) < 0$, then $\mathcal{H}(a, b)$ is negative definite and (a, b) is a **maximum**.

If $D_2(a, b) \neq 0$ and the above conditions do not hold, then $\mathcal{H}(a, b)$ is indefinite and (a, b) is **not a minimum or maximum**.

If $D_2(a, b) = 0$, then the second derivative test is inconclusive, and we need more

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A \left(h + \frac{B}{A}k \right)^2 + \frac{AC - B^2}{A}k^2$$

So $Q(x)$ is positive definite

if A and $\frac{AC - B^2}{A}$ are both positive, i.e. $A > 0$ and $AC - B^2 > 0$;

$Q(x)$ is negative definite

if A and $\frac{AC - B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$;

$Q(x)$ is indefinite if A and $\frac{AC - B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$, and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$; \implies not maximum nor minimum

Let's start with the 2-variable case: any 2-variable quadratic form has the form $Ah^2 + 2Bhk + Ck^2$. (We are interested in $f_{xx}(a, b)h^2 + 2f_{xy}(a, b)hk + f_{yy}(a, b)k^2$.)

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A \left(h + \frac{B}{A}k \right)^2 + \frac{AC - B^2}{A}k^2$$

$$\det \begin{pmatrix} A & B \\ B & C \end{pmatrix}$$

So $Q(x)$ is positive definite

if A and $\frac{AC-B^2}{A}$ are both positive, i.e. $A > 0$ and $\boxed{AC - B^2} > 0$;

$Q(x)$ is negative definite

if A and $\frac{AC-B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$;

$Q(x)$ is indefinite if A and $\frac{AC-B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

An n -variable quadratic form $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ can be encoded in a $n \times n$ matrix \mathcal{H} that is symmetric (i.e. $\mathcal{H}_{ij} = \mathcal{H}_{ji}$): let $Q(\mathbf{x}) = \mathcal{H}_{11}x_1^2 + \dots$. Let D_i be the determinant of the

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

An n -variable quadratic form $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ can be encoded in a $n \times n$ matrix \mathcal{H} that is symmetric (i.e. $\mathcal{H}_{ij} = \mathcal{H}_{ji}$): let $Q(\mathbf{x}) = \mathcal{H}_{11}x_1^2 + \dots$. Let D_i be the determinant of the

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A\left(h + \frac{B}{A}k\right)^2 + \frac{AC - B^2}{A}k^2$$

So $Q(x)$ is positive definite

if A and $\frac{AC - B^2}{A}$ are both positive, i.e. $A > 0$ and $AC - B^2 > 0$

$Q(x)$ is negative definite

if A and $\frac{AC - B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$

$Q(x)$ is indefinite if A and $\frac{AC - B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

Definition: A quadratic form Q is:

- *positive definite* if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies maximum
- *negative definite* if $Q(\mathbf{x}) < 0$ for all $\mathbf{x} \neq \mathbf{0}$; \implies minimum
- *indefinite* if $Q(\mathbf{x}) > 0$ for some $\mathbf{x} \neq \mathbf{0}$,
and $Q(\mathbf{x}) < 0$ for some other $\mathbf{x} \neq \mathbf{0}$. \implies not maximum nor minimum

An n -variable quadratic form $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ can be encoded in a $n \times n$ matrix \mathcal{H} that is symmetric (i.e. $\mathcal{H}_{ij} = \mathcal{H}_{ji}$): let $Q(\mathbf{x}) = \mathcal{H}_{11}x_1^2 + \dots$. Let D_i be the determinant of the

In the previous example where $B = 0$, we can quickly tell the definiteness from the signs of A and C . In the general case, we will have to complete the square:

$$Ah^2 + 2Bhk + Ck^2 = A \left(h + \frac{B}{A}k \right)^2 + \frac{AC - B^2}{A}k^2 \quad \det \begin{pmatrix} A & B \\ B & C \end{pmatrix}$$

So $Q(x)$ is positive definite

if A and $\frac{AC - B^2}{A}$ are both positive, i.e. $A > 0$ and $AC - B^2 > 0$

$Q(x)$ is negative definite

if A and $\frac{AC - B^2}{A}$ are both negative, i.e. $A < 0$ and $AC - B^2 > 0$

$Q(x)$ is indefinite if A and $\frac{AC - B^2}{A}$ have different signs, i.e. $A \neq 0$ and $AC - B^2 < 0$.

Example: Find and classify the critical points of $f(x, y) = xy + y^2e^x$.

§13.1: Classifying Critical Points

Recall that a critical point of a single-variable function f is where the derivative f' is zero. It is the x -value of a point where the graph has a horizontal tangent.

§13.1: Classifying Critical Points

Recall that a critical point of a single-variable function f is where the derivative f' is zero. It is the x -value of a point where the graph has a horizontal tangent.

We make the same definition for functions of more variables.

Definition: A point (a_1, \dots, a_n) is a *critical point* of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ if $\nabla f(a_1, \dots, a_n) = \mathbf{0}$, i.e. if all its partial derivatives are 0.

For a 2-variable function, a critical point is the (x, y) coordinates of a point where the tangent plane to the graph is horizontal.

