

Remember from last Monday:


Fact: A linear system has either

- exactly one solution
- infinitely many solutions
- no solutions

EXAMPLE Two equations in two variables:

$$x_1 + x_2 = 10$$


$$-x_1 + x_2 = 0$$



one unique solution

$$x_1 - 2x_2 = -3$$


$$2x_1 - 4x_2 = 8$$



no solution

$$x_1 + x_2 = 3$$

$$-2x_1 - 2x_2 = -6$$



infinitely many solutions

Thinking geometrically can be helpful. (§1.3-1.7)

§1.3: Vector Equations

A **column vector** is a matrix with only one column.

Until Chapter 4, we will say “vector” to mean “column vector”.

§1.3: Vector Equations

A **column vector** is a matrix with only one column.

Until Chapter 4, we will say “vector” to mean “column vector”.

A vector \mathbf{u} is in \mathbb{R}^n if it has n rows, i.e. $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$

Example: $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ are vectors in \mathbb{R}^2 .

§1.3: Vector Equations


A **column vector** is a matrix with only one column.

Until Chapter 4, we will say “vector” to mean “column vector”.

A vector \mathbf{u} is in \mathbb{R}^n if it has n rows, i.e. $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$

Example: $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ are vectors in \mathbb{R}^2 .

Vectors in \mathbb{R}^2 and \mathbb{R}^3 have a geometric meaning: think of $\begin{bmatrix} x \\ y \end{bmatrix}$ as the point (x, y) in the plane.



There are two operations we can do on vectors:

addition: if $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$, then $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ \vdots \\ u_n + v_n \end{bmatrix}$.

There are two operations we can do on vectors:

addition: if $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$, then $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ \vdots \\ u_n + v_n \end{bmatrix}$.

scalar multiplication: if $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and c is a number (a **scalar**), then $c\mathbf{u} = \begin{bmatrix} cu_1 \\ cu_2 \\ \vdots \\ cu_n \end{bmatrix}$.

These satisfy the usual rules for arithmetic of numbers, e.g.

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}, \quad c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}, \quad 0\mathbf{u} = \mathbf{0} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Definition: Given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ in \mathbb{R}^n and scalars c_1, c_2, \dots, c_p , the vector

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is a *linear combination* of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ with *weights* c_1, c_2, \dots, c_p .

Definition: Given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ in \mathbb{R}^n and scalars c_1, c_2, \dots, c_p , the vector

$$c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_p \mathbf{v}_p$$

is a *linear combination* of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ with *weights* c_1, c_2, \dots, c_p .

Example: $\mathbf{u} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$. Some linear combinations of \mathbf{u} and \mathbf{v} are:

$$3\mathbf{u} + 2\mathbf{v} = \begin{bmatrix} 7 \\ 11 \end{bmatrix}.$$

$$\frac{1}{3}\mathbf{u} = \begin{bmatrix} 1/3 \\ 1 \end{bmatrix}.$$

$$\mathbf{u} - 3\mathbf{v} = \begin{bmatrix} -3 \\ 0 \end{bmatrix}.$$

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Definition: Given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ in \mathbb{R}^n and scalars c_1, c_2, \dots, c_p , the vector

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is a *linear combination* of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ with *weights* c_1, c_2, \dots, c_p .

Example: $\mathbf{u} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$. Some linear combinations of \mathbf{u} and \mathbf{v} are:

$$3\mathbf{u} + 2\mathbf{v} = \begin{bmatrix} 7 \\ 11 \end{bmatrix}.$$

$$\frac{1}{3}\mathbf{u} + 0\mathbf{v} = \begin{bmatrix} 1/3 \\ 1 \end{bmatrix}.$$

$$\mathbf{u} - 3\mathbf{v} = \begin{bmatrix} -3 \\ 0 \end{bmatrix}.$$

$$\mathbf{0} = 0\mathbf{u} + 0\mathbf{v} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Geometric interpretation of linear combinations:



Definition: Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ are in \mathbb{R}^n . The *span* of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$, written

$$\text{Span} \{ \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p \},$$


is the set of all linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$.

In other words, $\text{Span} \{ \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p \}$ is the set of all vectors which can be written as $x_1 \mathbf{v}_1 + x_2 \mathbf{v}_2 + \dots + x_p \mathbf{v}_p$ for any choice of scalars x_1, x_2, \dots, x_p .



Example: Span of one vector in \mathbb{R}^3

- $\text{Span}\{\mathbf{0}\} = \{\mathbf{0}\}$, because $c\mathbf{0} = \mathbf{0}$ for all scalars c .
- If \mathbf{u} is not the zero vector, then $\text{Span}\{\mathbf{u}\}$ is a line through the origin in the direction \mathbf{u} .

We can also say “ $\{\mathbf{u}\}$ spans a line through the origin”.



Example: Span of two vectors in \mathbb{R}^3



The vector equation

$$x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + \cdots + x_p \mathbf{a}_p = \mathbf{b}$$

has the **same solution set** as the linear system whose augmented matrix is

$$\left[\begin{array}{cccc|c} | & | & | & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_p & \mathbf{b} \\ | & | & | & | & | \end{array} \right] .$$

In particular, \mathbf{b} is a linear combination of $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ (i.e. \mathbf{b} is in $\text{Span} \{ \mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p \}$) if and only if there is a solution to the linear system with augmented matrix

$$\left[\begin{array}{cccc|c} | & | & | & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_p & \mathbf{b} \\ | & | & | & | & | \end{array} \right] .$$

§1.4: The Matrix Equation $A\mathbf{x} = \mathbf{b}$

We can think of the weights x_1, x_2, \dots, x_p as a vector.

§1.4: The Matrix Equation $A\mathbf{x} = \mathbf{b}$

We can think of the weights x_1, x_2, \dots, x_p as a vector.

The **product** of an $m \times p$ matrix A and a vector \mathbf{x} in \mathbb{R}^p is **the linear combination** of the columns of A using the entries of \mathbf{x} as weights:

$$A\mathbf{x} = \begin{bmatrix} | & | & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_p \\ | & | & | & | \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_p \end{bmatrix} = x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + \cdots + x_p \mathbf{a}_p.$$

§1.4: The Matrix Equation $A\mathbf{x} = \mathbf{b}$

We can think of the weights x_1, x_2, \dots, x_p as a vector.

The **product** of an $m \times p$ matrix A and a vector \mathbf{x} in \mathbb{R}^p is **the linear combination** of the columns of A using the entries of \mathbf{x} as weights:

$$A\mathbf{x} = \begin{bmatrix} | & | & & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_p \\ | & | & & | \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_p \end{bmatrix} = x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + \cdots + x_p \mathbf{a}_p.$$

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = -2 \begin{bmatrix} 4 \\ 2 \\ 14 \end{bmatrix} + 2 \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = -2 \begin{bmatrix} 4 \\ 2 \\ 14 \end{bmatrix} + 2 \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

There is another way to compute $A\mathbf{x}$, one row of A at a time:

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4(-2) + 3(2) \\ \\ \end{bmatrix} = \begin{bmatrix} -2 \\ \\ \end{bmatrix}.$$

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = -2 \begin{bmatrix} 4 \\ 2 \\ 14 \end{bmatrix} + 2 \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

There is another way to compute $A\mathbf{x}$, one row of A at a time:

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4(-2) + 3(2) \\ 2(-2) + 6(2) \\ 14(-2) + 10(2) \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = -2 \begin{bmatrix} 4 \\ 2 \\ 14 \end{bmatrix} + 2 \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

There is another way to compute $A\mathbf{x}$, one row of A at a time:

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4(-2) + 3(2) \\ 2(-2) + 6(2) \\ 14(-2) + 10(2) \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = -2 \begin{bmatrix} 4 \\ 2 \\ 14 \end{bmatrix} + 2 \begin{bmatrix} 3 \\ 6 \\ 10 \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

There is another way to compute $A\mathbf{x}$, one row of A at a time:

Example:
$$\begin{bmatrix} 4 & 3 \\ 2 & 6 \\ 14 & 10 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} 4(-2) + 3(2) \\ 2(-2) + 6(2) \\ 14(-2) + 10(2) \end{bmatrix} = \begin{bmatrix} -2 \\ 8 \\ -8 \end{bmatrix}.$$

Warning: The product $A\mathbf{x}$ is only defined if the number of columns of A equals the number of rows of x . The number of rows of $A\mathbf{x}$ is the number of rows of A .

It is easy to check that $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$ and $A(c\mathbf{u}) = cA\mathbf{u}$.

We have three ways of viewing the same problem:

1. The system of linear equations with augmented matrix $[A|\mathbf{b}]$,
2. The vector equation $x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_p\mathbf{a}_p = \mathbf{b}$,
3. The matrix equation $A\mathbf{x} = \mathbf{b}$.

We have three ways of viewing the same problem:

1. The system of linear equations with augmented matrix $[A|\mathbf{b}]$,
2. The vector equation $x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_p\mathbf{a}_p = \mathbf{b}$,
3. The matrix equation $A\mathbf{x} = \mathbf{b}$.

So these three things are the same:

1. The system of linear equations with augmented matrix $[A|\mathbf{b}]$ has a solution,
2. \mathbf{b} is a linear combination of the columns of A (or \mathbf{b} is in the span of the columns of A),
3. The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution.

(The three problems have the same solution set.)

We have three ways of viewing the same problem:

1. The system of linear equations with augmented matrix $[A|\mathbf{b}]$,
2. The vector equation $x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_p\mathbf{a}_p = \mathbf{b}$,
3. The matrix equation $A\mathbf{x} = \mathbf{b}$.

So these three things are the same:

1. The system of linear equations with augmented matrix $[A|\mathbf{b}]$ has a solution,
2. \mathbf{b} is a linear combination of the columns of A (or \mathbf{b} is in the span of the columns of A),
3. The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution.

(The three problems have the same solution set.)

Another way of saying this: The span of the columns of A is the set of vectors \mathbf{b} for which $A\mathbf{x} = \mathbf{b}$ has a solution.

Theorem 4: Existence of solutions to linear systems: The following statements are logically equivalent (i.e. for any particular matrix A , they are all true or all false):

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- b. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
- c.
- d.

Theorem 4: Existence of solutions to linear systems: The following statements are logically equivalent (i.e. for any particular matrix A , they are all true or all false):

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- b. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
- c. The columns of A span \mathbb{R}^m (i.e. $\text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p\} = \mathbb{R}^m$).
- d.

Theorem 4: Existence of solutions to linear systems: For an $m \times n$ matrix A , the following statements are logically equivalent (i.e. for any particular matrix A , they are all true or all false):

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- b. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
- c. The columns of A span \mathbb{R}^m .
- d. $\text{rref}(A)$ has a pivot in every row.

Theorem 4: Existence of solutions to linear systems: For an $m \times n$ matrix A , the following statements are logically equivalent (i.e. for any particular matrix A , they are all true or all false):

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- b. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
- c. The columns of A span \mathbb{R}^m .
- d. $\text{rref}(A)$ has a pivot in every row.

Warning: the theorem says nothing about the **uniqueness** of the solution.

Theorem 4: Existence of solutions to linear systems: For an $m \times n$ matrix A , the following statements are logically equivalent (i.e. for any particular matrix A , they are all true or all false):

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- b. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
- c. The columns of A span \mathbb{R}^m .
- d. $\text{rref}(A)$ has a pivot in every row.

Warning: the theorem says nothing about the **uniqueness** of the solution.

Proof: (outline): By previous discussion, (a), (b) and (c) are logically equivalent. So, to finish the proof, we only need to show that (a) and (d) are logically equivalent, i.e. we need to show that,

- if (d) is true, then (a) is true;
- if (d) is false, then (a) is false.

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued)

Suppose (d) is true.

So (a) is true.

Suppose (d) is false.

So (a) is false

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued)

Suppose (d) is true. Then, for every \mathbf{b} in \mathbb{R}^m , the augmented matrix $[A|\mathbf{b}]$ row-reduces to $[\text{rref}(A)|\mathbf{d}]$ for some \mathbf{d} in \mathbb{R}^m . This does not have a row of the form $[0 \dots 0|*]$, so, by Theorem 2, $A\mathbf{x} = \mathbf{b}$ is consistent. So (a) is true.

Suppose (d) is false.

So (a) is false

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued)

Suppose (d) is true. Then, for every \mathbf{b} in \mathbb{R}^m , the augmented matrix $[A|\mathbf{b}]$ row-reduces to $[\text{rref}(A)|\mathbf{d}]$ for some \mathbf{d} in \mathbb{R}^m . This does not have a row of the form $[0 \dots 0 | *]$, so, by Theorem 2, $A\mathbf{x} = \mathbf{b}$ is consistent. So (a) is true.

Suppose (d) is false. We want to find a **counterexample** to (a): i.e. we want to find a vector \mathbf{b} in \mathbb{R}^m such that $A\mathbf{x} = \mathbf{b}$ has no solution.

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued) Suppose (d) is false. We want to find a **counterexample** to (a): i.e. we want to find a vector \mathbf{b} in \mathbb{R}^m such that $A\mathbf{x} = \mathbf{b}$ has no solution.

$\text{rref}(A)$ does not have a pivot in every row, so its last row is $[0 \dots 0]$.

Example:

$$\begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 + 2R_1} \begin{bmatrix} 1 & -3 \\ 0 & 0 \end{bmatrix}$$

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued) Suppose (d) is false. We want to find a **counterexample** to (a): i.e. we want to find a vector \mathbf{b} in \mathbb{R}^m such that $A\mathbf{x} = \mathbf{b}$ has no solution.

$\text{rref}(A)$ does not have a pivot in every row, so its last row is $[0 \dots 0]$.

Let $\mathbf{d} = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$. Then the linear system with augmented matrix $[\text{rref}(A) | \mathbf{d}]$ is inconsistent.

Example:

$$\begin{bmatrix} 1 & -3 \\ -2 & 6 \end{bmatrix} \xrightarrow{R_2 \rightarrow R_2 + 2R_1} \left[\begin{array}{cc|c} 1 & -3 & 1 \\ 0 & 0 & 1 \end{array} \right]$$

- a. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
- d. $\text{rref}(A)$ has a pivot in every row.

Proof: (continued) Suppose (d) is false. We want to find a **counterexample** to (a): i.e. we want to find a vector \mathbf{b} in \mathbb{R}^m such that $A\mathbf{x} = \mathbf{b}$ has no solution.

$\text{rref}(A)$ does not have a pivot in every row, so its last row is $[0 \dots 0]$.

Let $\mathbf{d} = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$. Then the linear system with augmented matrix $[\text{rref}(A)|\mathbf{d}]$ is inconsistent.

Now we apply the row operations in reverse to get an equivalent linear system $[A|\mathbf{b}]$ that is inconsistent.

Example:

$$\left[\begin{array}{cc|c} 1 & -3 & -1 \\ -2 & 6 & -1 \end{array} \right] \xrightarrow[\textcolor{blue}{R_2 \rightarrow R_2 - 2R_1}]{\textcolor{blue}{R_2 \rightarrow R_2 + 2R_1}} \left[\begin{array}{cc|c} 1 & -3 & \textcolor{blue}{1} \\ 0 & 0 & \textcolor{blue}{1} \end{array} \right]$$

§1.5: Solution Sets of Linear Systems

Goal: use vector notation to give geometric descriptions of solution sets to compare the solution sets of $A\mathbf{x} = \mathbf{b}$ and of $A\mathbf{x} = \mathbf{0}$.

Definition: A linear system is *homogeneous* if the right hand side is the zero vector, i.e.

$$A\mathbf{x} = \mathbf{0}.$$

§1.5: Solution Sets of Linear Systems

Goal: use vector notation to give geometric descriptions of solution sets to compare the solution sets of $A\mathbf{x} = \mathbf{b}$ and of $A\mathbf{x} = \mathbf{0}$.

Definition: A linear system is *homogeneous* if the right hand side is the zero vector, i.e.

$$A\mathbf{x} = \mathbf{0}.$$

When we row-reduce $[A|\mathbf{0}]$, the right hand side stays $\mathbf{0}$, so the reduced echelon form does not have a row of the form $[0 \dots 0 | *]$. So a homogeneous system is *always consistent*.

§1.5: Solution Sets of Linear Systems

Goal: use vector notation to give geometric descriptions of solution sets to compare the solution sets of $A\mathbf{x} = \mathbf{b}$ and of $A\mathbf{x} = \mathbf{0}$.

Definition: A linear system is *homogeneous* if the right hand side is the zero vector, i.e.

$$A\mathbf{x} = \mathbf{0}.$$

When we row-reduce $[A|\mathbf{0}]$, the right hand side stays $\mathbf{0}$, so the reduced echelon form does not have a row of the form $[0 \dots 0|*]$. So a homogeneous system is *always consistent*.

In fact, $\mathbf{x} = \mathbf{0}$ is always a solution, because $A\mathbf{0} = \mathbf{0}$. The solution $\mathbf{x} = \mathbf{0}$ called the *trivial solution*.

A *non-trivial solution* \mathbf{x} is a solution where at least one x_i is non-zero.

In our first example:

- The solution set of $A\mathbf{x} = \mathbf{0}$ is a line through the origin parallel to \mathbf{v} .
- The solution set of $A\mathbf{x} = \mathbf{b}$ is a line through \mathbf{p} parallel to \mathbf{v} .

In our second example:

- The solution set of $A\mathbf{x} = \mathbf{0}$ is a plane through the origin parallel to \mathbf{u} and \mathbf{v} .
- The solution set of $A\mathbf{x} = \mathbf{b}$ is a plane through \mathbf{p} parallel to \mathbf{u} and \mathbf{v} .

In both cases: to get the solution set of $A\mathbf{x} = \mathbf{b}$, start with the solution set of $A\mathbf{x} = \mathbf{0}$ and **translate** it by \mathbf{p} .

\mathbf{p} is called a **particular solution** (one solution out of many).

In our first example:

- The solution set of $A\mathbf{x} = \mathbf{0}$ is a line through the origin parallel to \mathbf{v} .
- The solution set of $A\mathbf{x} = \mathbf{b}$ is a line through \mathbf{p} parallel to \mathbf{v} .

In our second example:

- The solution set of $A\mathbf{x} = \mathbf{0}$ is a plane through the origin parallel to \mathbf{u} and \mathbf{v} .
- The solution set of $A\mathbf{x} = \mathbf{b}$ is a plane through \mathbf{p} parallel to \mathbf{u} and \mathbf{v} .


In both cases: to get the solution set of $A\mathbf{x} = \mathbf{b}$, start with the solution set of $A\mathbf{x} = \mathbf{0}$ and **translate** it by \mathbf{p} .

\mathbf{p} is called a **particular solution** (one solution out of many).

In general:

Theorem 6: Solutions and homogeneous equations: Suppose \mathbf{p} is a solution to $A\mathbf{x} = \mathbf{b}$. Then the solution set to $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$, where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

Theorem 6: Solutions and homogeneous equations: Suppose \mathbf{p} is a solution to $A\mathbf{x} = \mathbf{b}$. Then the solution set to $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$, where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.



Parallel solution sets of $A\mathbf{x} = \mathbf{b}$ and $A\mathbf{x} = \mathbf{0}$.

Theorem 6: Solutions and homogeneous equations: Suppose \mathbf{p} is a solution to $A\mathbf{x} = \mathbf{b}$. Then the solution set to $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$, where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

Proof: (outline)

We show that $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ is a solution:

$$\begin{aligned} & A(\mathbf{p} + \mathbf{v}_h) \\ &= A\mathbf{p} + A\mathbf{v}_h \\ &= \mathbf{b} + \mathbf{0} \\ &= \mathbf{b}. \end{aligned}$$

We also need to show that all solutions are of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ - see q25 in Section 1.5 of the textbook.

Question:

Suppose A is a matrix with $\text{rref}(A) = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Find the solution set to $A\mathbf{x} = A \begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$.

Question:

Suppose A is a matrix with $\text{rref}(A) = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Find the solution set to $A\mathbf{x} = A \begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$.

Answer:

$\text{rref}(A) \rightarrow$ the solution set to $A\mathbf{x} = \mathbf{0}$ is $\text{Span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}$ (see earlier today).

$\begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$ is a particular solution to $A\mathbf{x} = A \begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$.

So the solution set to $A\mathbf{x} = A \begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$ is $\begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix} + s \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$, where s can take any value.