

§1.1: Systems of Linear Equations

Example of a linear equation: $y = 5x + 2$

Can be rearranged to give: $(-5)x + (1)y = 2$

§1.1: Systems of Linear Equations

Example of a linear equation: $y = 5x + 2$

Can be rearranged to give: $(-5)x + (1)y = 2$

Another example: $3(x_1 + 2x_2) + 1 = x_1 + 1$

Rearrangement: $(2)x_1 + (6)x_2 = 0$

Another example: $x_2 = \sqrt{2}(\sqrt{6} - x_1) + x_3$

Rearrangement: $\sqrt{2}x_1 + (1)x_2 + (-1)x_3 = 2\sqrt{3}$

§1.1: Systems of Linear Equations

Example of a linear equation: $y = 5x + 2$

Can be rearranged to give: $(-5)x + (1)y = 2$

Another example: $3(x_1 + 2x_2) + 1 = x_1 + 1$

Rearrangement: $(2)x_1 + (6)x_2 = 0$

Another example: $x_2 = \sqrt{2}(\sqrt{6} - x_1) + x_3$

Rearrangement: $\sqrt{2}x_1 + (1)x_2 + (-1)x_3 = 2\sqrt{3}$

Not linear: $x_2 = 2\sqrt{x_1}$

$$xy + x = e^5$$

§1.1: Systems of Linear Equations

Example of a linear equation: $y = 5x + 2$

Can be rearranged to give: $(-5)x + (1)y = 2$

Another example: $3(x_1 + 2x_2) + 1 = x_1 + 1$

Rearrangement: $(2)x_1 + (6)x_2 = 0$

Another example: $x_2 = \sqrt{2}(\sqrt{6} - x_1) + x_3$

Rearrangement: $\sqrt{2}x_1 + (1)x_2 + (-1)x_3 = 2\sqrt{3}$

Not linear: $x_2 = 2\sqrt{x_1}$

$$xy + x = e^5$$

In general, a **linear equation** is an equation of the form

$$a_1x_1 + a_2x_2 + \dots a_nx_n = b.$$

x_1, x_2, \dots, x_n are the **variables**. a_1, a_2, \dots, a_n are the **coefficients**.

In general, a **linear equation** is an equation of the form

$$a_1x_1 + a_2x_2 + \dots a_nx_n = b.$$

x_1, x_2, \dots, x_n are the **variables**. a_1, a_2, \dots, a_n are the **coefficients**.

Definition: A **system of linear equations** (or a **linear system**) is a collection of linear equations involving the same set of variables.

Example:

$$\begin{array}{rclcl} x & +y & & = & 3 \\ 3x & & +2z & = & -2 \end{array}$$

This is a system of **2 equations** in **3 variables**, x, y, z .

In general, a **linear equation** is an equation of the form

$$a_1x_1 + a_2x_2 + \dots a_nx_n = b.$$

x_1, x_2, \dots, x_n are the **variables**. a_1, a_2, \dots, a_n are the **coefficients**.

Definition: A **system of linear equations** (or a **linear system**) is a collection of linear equations involving the same set of variables.

Example:

$$\begin{array}{rrcr} x & +y & +0z & = 3 \\ 3x & +0y & +2z & = -2 \end{array}$$

This is a system of **2 equations** in **3 variables**, x, y, z .

In general, a **linear equation** is an equation of the form

$$a_1x_1 + a_2x_2 + \dots a_nx_n = b.$$

x_1, x_2, \dots, x_n are the **variables**. a_1, a_2, \dots, a_n are the **coefficients**.

Definition: A **system of linear equations** (or a **linear system**) is a collection of linear equations involving the same set of variables.

Example:	$\begin{array}{rcl} x & +y & = 3 \\ 3x & & +2z = -2 \end{array}$	A solution is: $(x, y, z) = (2, 1, -4).$
----------	--	---

This is a system of **2 equations** in **3 variables**, x, y, z .

Definition: A **solution** of a linear system is a list (s_1, s_2, \dots, s_n) of numbers that makes each equation a true statement when the values s_1, s_2, \dots, s_n are substituted for x_1, x_2, \dots, x_n respectively.

Definition: The **solution set** of a linear system is the set of all possible solutions.

Definition: A linear system is *consistent* if it has a solution,
and *inconsistent* if it does not have a solution.

Fact: A linear system has either

- | | |
|-----------------------------|--------------|
| • exactly one solution | consistent |
| • infinitely many solutions | consistent |
| • no solutions | inconsistent |


Definition: A linear system is *consistent* if it has a solution,
and *inconsistent* if it does not have a solution.

Fact: A linear system has either

- exactly one solution consistent
- infinitely many solutions consistent
- no solutions inconsistent


EXAMPLE Two equations in two variables:

$$\begin{aligned}x_1 + x_2 &= 10 \\ -x_1 + x_2 &= 0\end{aligned}$$




one unique solution
consistent

$$\begin{aligned}x_1 - 2x_2 &= -3 \\ 2x_1 - 4x_2 &= 8\end{aligned}$$



no solution
inconsistent

$$\begin{aligned}x_1 + x_2 &= 3 \\ -2x_1 - 2x_2 &= -6\end{aligned}$$




infinitely many solutions
consistent


$$ax + by + cz = d, \text{ or } z = a'x + b'y + d'$$

EXAMPLE: Three equations in three variables. Each equation determines a plane in 3-space.


i) The planes intersect in one point. (*one solution*)



ii) The planes intersect in one line. (*infinitely many solutions*)



iii) There is no point in common to all three planes. (*no solution*)




Which of these cases are consistent?


$$ax + by + cz = d, \text{ or } z = a'x + b'y + d'$$

EXAMPLE: Three equations in three variables. Each equation determines a plane in 3-space.


i) The planes intersect in one point. (*one solution*)



ii) The planes intersect in one line. (*infinitely many solutions*)



iii) There is no point in common to all three planes. (*no solution*)



Which of these cases are consistent?

consistent

consistent

inconsistent

How to solve a linear system? **Example:**

$$x_1 - 2x_2 = -1$$

$$-x_1 + 3x_2 = 3$$

How to solve a linear system? **Example:**

$$\begin{array}{rcl} x_1 - 2x_2 & = & -1 \\ -x_1 + 3x_2 & = & 3 \end{array} \quad \rightarrow \quad \begin{array}{rcl} x_1 - 2x_2 & = & -1 \\ & x_2 & = & 2 \end{array} \quad \rightarrow \quad \begin{array}{rcl} x_1 & = & 3 \\ & x_2 & = & 2 \end{array}$$

How to solve a linear system? **Example:**

$$\begin{array}{lcl} R_1 & x_1 - 2x_2 & = -1 \\ R_2 & -x_1 + 3x_2 & = 3 \end{array} \quad \xrightarrow{R_2 + R_1} \quad \begin{array}{lcl} & x_1 - 2x_2 & = -1 \\ & x_2 & = 2 \end{array} \quad \xrightarrow{R_1 + 2R_2} \quad \begin{array}{lcl} & x_1 & = 3 \\ & x_2 & = 2 \end{array}$$

How to solve a linear system? **Example:**

$$R_1 \quad x_1 - 2x_2 = -1$$




$$R_2 \quad -x_1 + 3x_2 = 3$$

$$\rightarrow \quad x_1 - 2x_2 = -1$$

$$R_2 + R_1 \rightarrow \quad x_2 = 2$$




$$R_1 \rightarrow R_1 + 2R_2 \rightarrow \quad x_1 = 3$$

$$x_2 = 2$$



How to solve a linear system? **Example:**

$$\begin{array}{lcl} R_1 & x_1 - 2x_2 & = -1 \\ R_2 & -x_1 + 3x_2 & = 3 \end{array} \quad \rightarrow \quad \begin{array}{lcl} & x_1 - 2x_2 & = -1 \\ R_2 + R_1 & & x_2 = 2 \end{array} \quad \begin{array}{lcl} R_1 \rightarrow +2R_2 & & x_1 = 3 \\ & & x_2 = 2 \end{array}$$



Definition: Two linear systems are *equivalent* if they have the same solution set.

General strategy for solving a linear system: replace one system with an equivalent system that is easier to solve.

Simplify the writing by using **matrix notation**:

$$\begin{array}{lcl}
 R_1 & x_1 - 2x_2 = -1 & \\
 R_2 & -x_1 + 3x_2 = 3 &
 \end{array}
 \xrightarrow{R_2 + R_1}
 \begin{array}{lcl}
 & x_1 - 2x_2 = -1 & \\
 & x_2 = 2 &
 \end{array}
 \xrightarrow{R_1 + 2R_2}
 \begin{array}{lcl}
 & x_1 = 3 & \\
 & x_2 = 2 &
 \end{array}$$

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

coefficient of x_1
coefficient of x_2
right hand side

Simplify the writing by using **matrix notation**:

$$\begin{array}{l} R_1 \quad x_1 - 2x_2 = -1 \\ R_2 \quad -x_1 + 3x_2 = 3 \end{array} \quad \xrightarrow{R_2 + R_1} \quad \begin{array}{l} x_1 - 2x_2 = -1 \\ x_2 = 2 \end{array} \quad \xrightarrow{R_1 + 2R_2} \quad \begin{array}{l} x_1 = 3 \\ x_2 = 2 \end{array}$$

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

coefficient of x_1
coefficient of x_2
right hand side

The **augmented matrix** of a linear system contains the right hand side:

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right]$$

The **coefficient matrix** of a linear system is the left hand side only:

$$\left[\begin{array}{cc} 1 & -2 \\ -1 & 3 \end{array} \right]$$

$$\begin{array}{rcl}
 R_1 & x_1 - 2x_2 & = -1 \\
 R_2 & -x_1 + 3x_2 & = 3
 \end{array}
 \quad \xrightarrow{\quad} \quad
 \begin{array}{rcl}
 & x_1 - 2x_2 & = -1 \\
 R_2 + R_1 & \rightarrow & x_2 = 2
 \end{array}
 \quad \xrightarrow{R_1 \rightarrow R_1 + 2R_2} \quad
 \begin{array}{rcl}
 & x_1 & = 3 \\
 & x_2 & = 2
 \end{array}$$

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

Elementary row operations:

1. **Replacement**: add a multiple of one row to another row. $R_i \rightarrow R_i + cR_j$
2. **Interchange**: interchange two rows. $R_i \rightarrow R_j, R_j \rightarrow R_i$
3. **Scaling**: multiply all entries in a row by a nonzero constant. $R_i \rightarrow cR_i, c \neq 0$

$$\begin{array}{rcl}
 R_1 & x_1 - 2x_2 & = -1 \\
 R_2 & -x_1 + 3x_2 & = 3
 \end{array}
 \xrightarrow{R_2 + R_1}
 \begin{array}{rcl}
 & x_1 - 2x_2 & = -1 \\
 & x_2 & = 2
 \end{array}
 \xrightarrow{R_1 + 2R_2}
 \begin{array}{rcl}
 & x_1 & = 3 \\
 & x_2 & = 2
 \end{array}$$

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

Elementary row operations:

1. **Replacement**: add a multiple of one row to another row. $R_i \rightarrow R_i + cR_j$
2. **Interchange**: interchange two rows. $R_i \rightarrow R_j, R_j \rightarrow R_i$
3. **Scaling**: multiply all entries in a row by a nonzero constant. $R_i \rightarrow cR_i, c \neq 0$

Definition: Two matrices are *row equivalent* if one can be transformed into the other by a sequence of elementary row operations.

$$\begin{array}{rcl}
 R_1 & x_1 - 2x_2 & = -1 \\
 R_2 & -x_1 + 3x_2 & = 3
 \end{array}
 \xrightarrow{R_2 + R_1}
 \begin{array}{rcl}
 & x_1 - 2x_2 & = -1 \\
 & x_2 & = 2
 \end{array}
 \xrightarrow{R_1 + 2R_2}
 \begin{array}{rcl}
 & x_1 & = 3 \\
 & x_2 & = 2
 \end{array}$$

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & -2 & -1 \\ 0 & 1 & 2 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

Elementary row operations:

1. **Replacement**: add a multiple of one row to another row. $R_i \rightarrow R_i + cR_j$
2. **Interchange**: interchange two rows. $R_i \rightarrow R_j, R_j \rightarrow R_i$
3. **Scaling**: multiply all entries in a row by a nonzero constant. $R_i \rightarrow cR_i, c \neq 0$

Definition: Two matrices are *row equivalent* if one can be transformed into the other by a sequence of elementary row operations.

Fact: If the augmented matrices of two linear systems are row equivalent, then the two systems have the same solution set, i.e. they are equivalent linear systems.

Warning: Do not do multiple elementary row operations at the same time, **except** adding multiples of **the same** row to several rows.

$$\begin{aligned}x_1 - 2x_2 &= 1 \\ -x_1 + 3x_2 &= 3\end{aligned}$$

$$x_2 = 2$$

$$x_2 = 2$$

These are NOT
equivalent systems.

$$\left[\begin{array}{cc|c} 1 & -2 & -1 \\ -1 & 3 & 3 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 0 & 1 & 2 \\ 0 & 1 & 2 \end{array} \right] \begin{array}{l} \leftarrow R_1 + R_2 \\ \leftarrow R_2 + R_1 \end{array}$$

$$\begin{aligned}x_1 - 2x_2 &= -3 \\ x_2 &= 16 \\ x_3 &= 3\end{aligned} \quad \left[\begin{array}{ccc|c} 1 & -2 & 0 & -3 \\ 0 & 1 & 0 & 16 \\ 0 & 0 & 1 & 3 \end{array} \right] \begin{array}{l} \leftarrow R_1 - R_3 \\ \leftarrow R_2 + 4R_3 \end{array}$$

$$\begin{aligned}x_1 &= 29 \\ x_2 &= 16 \\ x_3 &= 3\end{aligned} \quad \left[\begin{array}{ccc|c} 1 & 0 & 0 & 29 \\ 0 & 1 & 0 & 16 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

Two fundamental questions:

1. **Existence** of solutions: is the system consistent?
2. **Uniqueness** of solutions: if a solution exists, is it the only one?

Two fundamental questions:

1. **Existence** of solutions: is the system consistent?
2. **Uniqueness** of solutions: if a solution exists, is it the only one?

Answering this requires less work than finding the solution.

Example:

$$\begin{array}{rrcr} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ -4x_1 & + & 5x_2 & + & 9x_3 & = & -9 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ -4 & 5 & 9 & -9 \end{array} \right]$$

$$\begin{array}{rrcr} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ & & - & 3x_2 & + & 13x_3 & = & -9 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & -3 & 13 & -9 \end{array} \right]$$

$$\begin{array}{rrcr} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & - & 3x_2 & + & 13x_3 & = & -9 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & -3 & 13 & -9 \end{array} \right]$$

$$\begin{array}{rrcr} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array} \quad \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right]$$

We can stop here:
back-substitution shows
that we can find a unique
solution.

echelon

$$\begin{bmatrix} 0 & \blacksquare & * & * & * & * & * & * & * & * & * \\ 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * \end{bmatrix}$$

reduced echelon

$$\begin{bmatrix} 0 & 1 & * & 0 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 1 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & * & * \end{bmatrix}$$

Are these matrices in echelon form, reduced echelon form, or neither?

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

echelon

$$\begin{bmatrix} 0 & \blacksquare & * & * & * & * & * & * & * & * & * \\ 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * \end{bmatrix}$$

reduced echelon

$$\begin{bmatrix} 0 & 1 & * & 0 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 1 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & * & * \end{bmatrix}$$

Are these matrices in echelon form, reduced echelon form, or neither?

$$\begin{bmatrix} 1 & 0 & 0 \\ \textcolor{red}{1} & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

neither

$$\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

reduced
echelon

$$\begin{bmatrix} 1 & \textcolor{red}{1} & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

echelon

$$\begin{bmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

reduced
echelon

$$\begin{array}{rclcrcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ -4x_1 & + & 5x_2 & + & 9x_3 & = & -9 \end{array}$$

$$\begin{array}{rclcrcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & 2x_2 & - & 8x_3 & = & 8 \\ & & -3x_2 & + & 13x_3 & = & -9 \end{array}$$

$$\begin{array}{rclcrcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & -3x_2 & + & 13x_3 & = & -9 \end{array}$$

$$\begin{array}{rclcl} x_1 & - & 2x_2 & + & x_3 & = & 0 \\ & & x_2 & - & 4x_3 & = & 4 \\ & & & & x_3 & = & 3 \end{array}$$

$$\begin{array}{rcl} x_1 & - & 2x_2 & = & -3 \\ & & x_2 & = & 16 \\ & & & x_3 & = & 3 \end{array}$$

$$\begin{array}{rcl} x_1 & & = 29 \\ & x_2 & = 16 \\ & & x_3 = 3 \end{array}$$

Theorem: Any matrix A is row-equivalent to exactly one reduced echelon matrix, which is called its **reduced echelon form** and written $\text{rref}(A)$.

General strategy for solving a linear system: apply row operations to its augmented matrix to obtain its rref .

General strategy for determining existence/uniqueness of solutions: apply row operations to its augmented matrix to obtain an **echelon form**, i.e. a row-equivalent echelon matrix.

These processes of row operations (to get to echelon or reduced echelon form) are called **row reduction**.

Theorem: Any matrix A is row-equivalent to exactly one reduced echelon matrix, which is called its **reduced echelon form** and written $\text{rref}(A)$.

General strategy for solving a linear system: apply row operations to its augmented matrix to obtain its rref.

General strategy for determining existence/uniqueness of solutions: apply row operations to its augmented matrix to obtain an **echelon form**, i.e. a row-equivalent echelon matrix.

Warning: an echelon form is not unique. Its entries depend on the row operations we used. But its pattern of ■ and * is unique.

These processes of row operations (to get to echelon or reduced echelon form) are called **row reduction**.

Row reduction:

augmented matrix of linear
system



echelon
form



existence and
uniqueness of
solutions




reduced
echelon
form



solution set

Row reduction:



The rest of this section:

- The row reduction algorithm
- Getting the solution, existence/uniqueness from the (reduced) echelon form

Important terms in the row reduction algorithm:

- **pivot position**: the position of a leading entry in a row-equivalent echelon matrix.
- **pivot**: a nonzero entry of the matrix that is used in a pivot position to create zeroes below it.
- **pivot column**: a column containing a pivot position.

The black squares are the pivot positions.

$$\begin{bmatrix} 0 & \blacksquare & * & * & * & * & * & * & * & * & * \\ 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * \end{bmatrix}$$

Check your answer: www.wolframalpha.com



rref{{0 , 3 , -6 , 6 , 4 , -5},{3 , -7 , 8 , -5 , 8 , 9},{1 , -3 , 4 , -3 , 2 , 5}}

☆

≡



Web Apps Examples Random

Input:

row reduce

$$\begin{pmatrix} 0 & 3 & -6 & 6 & 4 & -5 \\ 3 & -7 & 8 & -5 & 8 & 9 \\ 1 & -3 & 4 & -3 & 2 & 5 \end{pmatrix}$$

Result:

Step-by-step solution

$$\begin{pmatrix} 1 & 0 & -2 & 3 & 0 & -24 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{pmatrix}$$

Getting the solution set from the reduced echelon form:

A **basic variable** is a variable corresponding to a pivot column.

All other variables are **free variables**.

Example:

$$\left[\begin{array}{ccccc|c} 1 & 0 & -2 & 3 & 0 & -24 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{array} \right]$$

$$\begin{aligned} x_1 - 2x_3 + 3x_4 &= -24 \\ x_2 - 2x_3 + 2x_4 &= -7 \\ x_5 &= 4 \end{aligned}$$

basic variables: x_1, x_2, x_5 , free variables: x_3, x_4 .

Getting the solution set from the reduced echelon form:

A **basic variable** is a variable corresponding to a pivot column.

All other variables are **free variables**.

Example:

$$\left[\begin{array}{ccccc|c} 1 & 0 & -2 & 3 & 0 & -24 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{array} \right]$$

$$\begin{aligned} x_1 - 2x_3 + 3x_4 &= -24 \\ x_2 - 2x_3 + 2x_4 &= -7 \\ x_5 &= 4 \end{aligned}$$

basic variables: x_1, x_2, x_5 , free variables: x_3, x_4 .

The free variables can take any value. These values then uniquely determine the basic variables.

Example:

$$x_1 = -24 + 2x_3 - 3x_4$$

$$x_2 = -7 + 2x_3 - 2x_4$$

$$x_3 = x_3$$

$$x_4 = x_4$$

$$x_5 = 4$$

Getting the solution set from the reduced echelon form:

A **basic variable** is a variable corresponding to a pivot column.

All other variables are **free variables**.

Example:

$$\left[\begin{array}{ccccc|c} 1 & 0 & -2 & 3 & 0 & -24 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{array} \right]$$

$$\begin{aligned} x_1 - 2x_3 + 3x_4 &= -24 \\ x_2 - 2x_3 + 2x_4 &= -7 \\ x_5 &= 4 \end{aligned}$$

basic variables: x_1, x_2, x_5 , free variables: x_3, x_4 .

The free variables can take any value. These values then uniquely determine the basic variables.

Example:

$$\begin{aligned} x_1 &= -24 + 2x_3 - 3x_4 \\ x_2 &= -7 + 2x_3 - 2x_4 \\ x_3 &= x_3 \\ x_4 &= x_4 \\ x_5 &= 4 \end{aligned}$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} -24 + 2s - 3t \\ -7 + 2s - 2t \\ s \\ t \\ 4 \end{pmatrix}$$

Getting the solution set from the reduced echelon form:

Another example: reduced echelon form is $\left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & 15 \end{array} \right]$

Getting the solution set from the reduced echelon form:

Another example: reduced echelon form is $\left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & 15 \end{array} \right]$

$$0x_1 + 0x_2 + 0x_3 = 15 \longrightarrow$$

the system is inconsistent

Getting the solution set from the reduced echelon form:

Another example: reduced echelon form is $\left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & 15 \end{array} \right]$

$0x_1 + 0x_2 + 0x_3 = 15 \longrightarrow$
the system is inconsistent

Theorem 2: Existence and Uniqueness:

A linear system is consistent if and only if an echelon form of its augmented matrix has **no** row of the form $[0 \dots 0 | b]$ with $b \neq 0$.

If a linear system is consistent, then:

- it has a unique solution if there are no free variables;
- it has infinitely many solutions if there are free variables.