

A Hopf-algebraic lift of the down-up-chain on partitions to permutations

Amy Pang, LaCIM

based on the preprint [arXiv:1508.01570](https://arxiv.org/abs/1508.01570)

slides available at tinyurl.com/HopfLift

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1.

2.

3.

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1. restrict to \mathfrak{S}_{n-1}
- 2.
- 3.

$$\text{Res} \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}$$

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
- 3.

$$\begin{aligned}
 \text{Ind} \circ \text{Res} \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \right) &= \text{Ind} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) + \text{Ind} \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array} \right) \\
 &= \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} + \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & & & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array}
 \end{aligned}$$

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
3. choose an irreducible with probability $\frac{\dim(\text{irreducible})}{n \dim(\text{original})}$

$$\begin{aligned}
 \text{Ind} \circ \text{Res} \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \right) &= \text{Ind} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right) + \text{Ind} \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array} \right) \\
 &= \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \\ \hline \end{array} \frac{5}{25} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \frac{5}{25} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \frac{5}{25} + \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & & & \\ \hline \end{array} \frac{4}{25} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array} \frac{6}{25}
 \end{aligned}$$

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
3. choose an irreducible with probability $\frac{\dim(\text{irreducible})}{n \dim(\text{original})}$

Fulman, 2004: Stationary distribution is Plancherel: $\frac{(\dim \lambda)^2}{n!}$

Fulman's down-up chain on partitions

A chain on irreducible representations of \mathfrak{S}_n

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
3. choose an irreducible with probability $\frac{\dim(\text{irreducible})}{n \dim(\text{original})}$

Fulman, 2004: Stationary distribution is Plancherel: $\frac{(\dim \lambda)^2}{n!}$

Question: Is there a chain X_t on permutations, with uniform stationary distribution, so that

$\text{shape}(\text{RSK}(X_t)) = \text{down-up on partitions?}$

lift

lumping

My weekly film list

Sunday afternoon:



My weekly film list

Sunday afternoon:



My weekly film list

Sunday afternoon:




Monday morning:



My weekly film list

Sunday afternoon:





Monday morning:



My weekly film list

Sunday afternoon:





Monday morning:



My weekly film list

Sunday afternoon:



Monday morning:




My weekly film list

Sunday afternoon:



Monday morning:



My weekly film list

Sunday afternoon:




Monday morning:




My weekly film list

Sunday afternoon:



Monday morning:



Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.

1.

2.

3.

$(4, 1, 3, 5, 2)$

Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.

1. remove last letter
- 2.
- 3.

$(4, 1, 3, 5, 2)$

remove

$(4, 1, 3, 5)$

Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.

1. remove last letter
2. standardise
- 3.

$(4, 1, 3, 5, 2)$


remove

$(4, 1, 3, 5) \xrightarrow{\text{std}} (3, 1, 2, 4)$

Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.


1. remove last letter
2. standardise
3. insert n at random position



Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.


1. remove last letter
2. standardise
3. insert n at random position



Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.


1. remove last letter
2. standardise
3. insert n at random position



Lift: bottom-to-random-with-standardisation

Theorem (2015): Taking RSK-shape of the following chain gives the down-up chain on partitions, under an initial condition.

1. remove last letter
2. standardise
3. insert n at random position



Proof

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

Proof

symmetric
functions
partitions

Poirier-Reutenauer
standard tableaux

Malvenuto-
Reutenauer
permutations

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym} \xhookrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

Proof

symmetric
functions
partitions

Poirier-Reutenauer
standard tableaux

Malvenuto-
Reutenauer
permutations

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$



down-up on
partitions



down-up on
standard
tableaux




bottom-to-
random-with-
standardisation

Proof

symmetric
functions
partitions

Poirier-Reutenauer
standard tableaux

Malvenuto-
Reutenauer
permutations



... and a new lumping theorem for Markov chains from combinatorial Hopf algebras

Markov chains from combinatorial Hopf algebras

- graded Hopf algebra: $\mathcal{H} = \bigoplus \mathcal{H}_n$
- basis of \mathcal{H}_n is \mathcal{B}_n
- product mult : $\mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$
- “partial” coproduct $\Delta_{n-1,1} : \mathcal{H} \rightarrow \mathcal{H}_{n-1} \otimes \mathcal{H}_1$

Markov chains from combinatorial Hopf algebras

- graded Hopf algebra: $\mathcal{H} = \bigoplus \mathcal{H}_n$
- basis of \mathcal{H}_n is \mathcal{B}_n
- product $\text{mult} : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$
- “partial” coproduct $\Delta_{n-1,1} : \mathcal{H} \rightarrow \mathcal{H}_{n-1} \otimes \mathcal{H}_1$

For $x, y \in \mathcal{B}_n$:

$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x)$

Markov chains from combinatorial Hopf algebras

For $x, y \in \mathcal{B}_n$:

$$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x)$$

FQSym, **F** basis \rightsquigarrow bottom-to-random-with-standardisation:

$$\Delta_{4,1}((4, 1, 3, 5, 2)) = (3, 1, 2, 4) \otimes (1)$$

Markov chains from combinatorial Hopf algebras

For $x, y \in \mathcal{B}_n$:

$$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x)$$

FQSym, **F** basis \rightsquigarrow bottom-to-random-with-standardisation:

$$\begin{aligned} \text{mult} \circ \Delta_{4,1}((4, 1, 3, 5, 2)) &= \text{mult}((3, 1, 2, 4) \otimes (1)) \\ &= (5, 3, 1, 2, 4) + (3, 5, 1, 2, 4) \\ &\quad + (3, 1, 5, 2, 4) + (3, 1, 2, 5, 4) \\ &\quad + (3, 1, 2, 4, 5) \end{aligned}$$

Markov chains from combinatorial Hopf algebras

For $x, y \in \mathcal{B}_n$:

$$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x)$$

FQSym, **F** basis \rightsquigarrow bottom-to-random-with-standardisation:

$$\begin{aligned} \frac{1}{5} \text{mult} \circ \Delta_{4,1}((4, 1, 3, 5, 2)) &= \frac{1}{5} \text{mult}((3, 1, 2, 4) \otimes (1)) \\ &= \frac{1}{5} (5, 3, 1, 2, 4) + \frac{1}{5} (3, 5, 1, 2, 4) \\ &\quad + \frac{1}{5} (3, 1, 5, 2, 4) + \frac{1}{5} (3, 1, 2, 5, 4) \\ &\quad + \frac{1}{5} (3, 1, 2, 4, 5) \end{aligned}$$

Markov chains from combinatorial Hopf algebras

For $x, y \in \mathcal{B}_n$:

(Doob h -transform)

$$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x) \frac{\dim y}{\dim x}$$

Λ , Schur basis \rightsquigarrow down-up on partitions:

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
3. choose an irreducible with probability $\frac{\dim(\text{irreducible})}{n \dim(\text{original})}$

Markov chains from combinatorial Hopf algebras

For $x, y \in \mathcal{B}_n$:

(Doob h -transform)

$$\text{Prob}(x \rightarrow y) = \text{coefficient of } y \text{ in } \frac{1}{n} \text{mult} \circ \Delta_{n-1,1}(x) \frac{\dim y}{\dim x}$$

$$\dim(x) = \text{sum of coefficients in } \Delta_{1,\dots,1}(x)$$

Λ , Schur basis \rightsquigarrow down-up on partitions:

1. restrict to \mathfrak{S}_{n-1}
2. induce to \mathfrak{S}_n
3. choose an irreducible with probability $\frac{\dim(\text{irreducible})}{n \dim(\text{original})}$

Strong lumping theorem

$$\Lambda \leftarrow \text{FSym}$$

$$\mathcal{S}_{\text{shape}}(T) \leftarrow \mathbf{S}_T$$

Strong lumping theorem

$$\Lambda \longleftarrow \text{FSym}$$

$$\mathcal{S}_{\text{shape}(T)} \longleftarrow \mathbf{S}_T$$

In general:

$$\bar{\mathcal{B}} \longleftarrow \mathcal{B} : \text{lump}$$

Theorem (2014): If a map on the bases $\text{lump} : \mathcal{B} \rightarrow \bar{\mathcal{B}}$ extends linearly to a surjective Hopf morphism, then

$$\text{lump}(\text{chain on } \mathcal{B}_n) = \text{chain on } \bar{\mathcal{B}}_n.$$

Weak lumping theorem

$$\mathbf{FSym} \not\leftarrow \mathbf{FQSym}$$

Weak lumping theorem

$$\mathbf{FSym} \hookrightarrow \mathbf{FQSym}$$

$$\mathbf{S}_T \longrightarrow \sum_{\text{RSK}(\sigma)=T} \mathbf{F}_\sigma$$

Weak lumping theorem

$$\mathbf{FSym} \hookrightarrow \mathbf{FQSym}$$

$$\mathbf{S}_T \longrightarrow \sum_{\text{RSK}(\sigma)=T} \mathbf{F}_\sigma$$

In general:

$$\mathcal{B} \longleftarrow \tilde{\mathcal{B}} : \text{lump}$$

$$\text{lump}^* : x \longrightarrow \sum_{\text{lump}(\tilde{x})=x} \tilde{x}$$

Weak lumping theorem

$$\mathbf{FSym} \hookrightarrow \mathbf{FQSym}$$

$$\mathbf{S}_T \longrightarrow \sum_{\text{RSK}(\sigma)=T} \mathbf{F}_\sigma$$

In general:

$$\begin{aligned} \mathcal{B} &\longleftarrow \tilde{\mathcal{B}} : \text{lump} \\ \text{lump}^* : x &\longrightarrow \sum_{\text{lump}(\tilde{x})=x} \tilde{x} \end{aligned}$$

Theorem (2015): If this “preimage sum map” extends linearly to a Hopf morphism, then

$$\text{lump}(\text{chain on } \tilde{\mathcal{B}}_n) = \text{chain on } \mathcal{B}_n,$$

Weak lumping theorem

$$\mathbf{FSym} \hookrightarrow \mathbf{FQSym}$$

$$\mathbf{S}_T \longrightarrow \sum_{\text{RSK}(\sigma)=T} \mathbf{F}_\sigma$$

In general:

$$\begin{array}{ccc} \mathcal{B} & \longleftarrow & \tilde{\mathcal{B}} : \text{lump} \\ \text{lump}^* : x & \longrightarrow & \sum_{\text{lump}(\tilde{x})=x} \tilde{x} \end{array}$$

Theorem (2015): If this “preimage sum map” extends linearly to a Hopf morphism, then

$$\text{lump}(\text{chain on } \tilde{\mathcal{B}}_n) = \text{chain on } \mathcal{B}_n,$$

provided the starting distribution X_0 on $\tilde{\mathcal{B}}_n$ satisfies

$$\frac{X_0(\tilde{x})}{\dim(\tilde{x})} = \frac{X_0(\tilde{y})}{\dim(\tilde{y})} \text{ whenever } \text{lump}(\tilde{x}) = \text{lump}(\tilde{y}).$$

In general

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym}$$

$$\mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

In general

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym}$$

$$\mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

$$\begin{array}{ccc} \Lambda & \xleftarrow{\text{shape}} & \mathbf{FSym} \\ \downarrow \frac{1}{n} \text{mult} \circ \Delta_{n-1,1} & & \downarrow \frac{1}{n} \text{mult} \circ \Delta_{n-1,1} \\ \Lambda & \xleftarrow{\text{shape}} & \mathbf{FSym} \end{array}$$

$$\begin{array}{ccc} \mathbf{FSym} & \xrightarrow{\text{RSK}^*} & \mathbf{FQSym} \\ \downarrow \frac{1}{n} \text{mult} \circ \Delta_{n-1,1} & & \downarrow \frac{1}{n} \text{mult} \circ \Delta_{n-1,1} \\ \mathbf{FSym} & \xrightarrow{\text{RSK}^*} & \mathbf{FQSym} \end{array}$$

In general

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym}$$

$$\mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

$$\begin{array}{ccc} \bar{V} & \xleftarrow{\text{lump}} & V \\ \bar{\mathbf{T}} \downarrow & & \downarrow \mathbf{T} \\ \bar{V} & \xleftarrow{\text{lump}} & V \end{array}$$

$$\begin{array}{ccc} V & \xrightarrow{\text{lump}^*} & \tilde{V} \\ \mathbf{T} \downarrow & & \downarrow \tilde{\mathbf{T}} \\ V & \xrightarrow{\text{lump}^*} & \tilde{V} \end{array}$$

$\bar{\mathbf{T}}$ chain = lump (\mathbf{T} chain) \mathbf{T} chain = lump ($\tilde{\mathbf{T}}$ chain),
with condition on starting distribution

In general

$$\Lambda \xleftarrow{\text{shape}} \mathbf{FSym}$$

$$\mathbf{FSym} \xrightarrow{\text{RSK}^*} \mathbf{FQSym}$$

$$\begin{array}{ccc} \bar{V} & \xleftarrow{\text{lump}} & V \\ \bar{\mathbf{T}} \downarrow & & \downarrow \mathbf{T} \\ \bar{V} & \xleftarrow{\text{lump}} & V \end{array}$$

$$\begin{array}{ccc} V & \xrightarrow{\text{lump}^*} & \tilde{V} \\ \mathbf{T} \downarrow & & \downarrow \tilde{\mathbf{T}} \\ V & \xrightarrow{\text{lump}^*} & \tilde{V} \end{array}$$

$\bar{\mathbf{T}}$ chain = lump (\mathbf{T} chain) \mathbf{T} chain = lump ($\tilde{\mathbf{T}}$ chain),
with condition on starting distribution

Question: a simpler application without Hopf algebras?

A final question


Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.

$(5, 2, 4, 1, 3)$

A final question

Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.


$(5, 2, 4, 1, 3)$



$(5, 2, 4, 1)$


A final question

Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.




A final question

Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.




A final question

Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.




A final question

Fulman (2004): The bottom-to-random shuffle is a lift of the down-up chain on partitions, if starting at the identity.



Question: is there a Hopf proof? $\Lambda \leftarrow \dots \hookrightarrow$ shuffle algebra?



Thank you!