

Deformations of Comonoids-in-Species, Coalgebras, Hopf Algebras and their Quasisymmetric Invariants

C.Y. Amy Pang

Department of Mathematics,
Hong Kong Baptist University

Deformations of Quasisymmetric Invariants

Wanted: people who work with quasisymmetric invariants
e.g. chromatic symmetric functions for graphs,
Billera-Jia-Reiner invariant for matroids,
Ehrenborg rank-generating function for posets, ...

Deformations of Quasisymmetric Invariants

Wanted: people who work with quasisymmetric invariants
e.g. chromatic symmetric functions for graphs,
Billera-Jia-Reiner invariant for matroids,
Ehrenborg rank-generating function for posets, ...

- Idea:**
- Quasisymmetric invariants are the image of the Aguiar-Bergeron-Sottile morphism: $\text{coalgebra} \rightarrow QSym$.
 - There is a general framework to q -deform the coalgebra $\rightarrow q$ -deformation of quasisymmetric invariants.
 - I would like applications for the deformed invariants.

Deformations of Quasisymmetric Invariants

Wanted: people who work with quasisymmetric invariants
e.g. chromatic symmetric functions for graphs,
Billera-Jia-Reiner invariant for matroids,
Ehrenborg rank-generating function for posets, ...

- Idea:**
- Quasisymmetric invariants are the image of the Aguiar-Bergeron-Sottile morphism: $\text{coalgebra} \rightarrow QSym$.
 - There is a general framework to q -deform the coalgebra $\rightarrow q$ -deformation of quasisymmetric invariants.
 - I would like applications for the deformed invariants.


Currently, each coalgebra deformation is calculated case-by-case (if the coalgebra is noncocommutative).

Dream: Find a uniform deformation formula.

Example: q -chromatic quasisymmetric function (q CQSF)

how it arises from a coalgebra of graphs


Consider graphs on vertices
labelled $\{1, 2, \dots, n\}$,



Example: q -chromatic quasisymmetric function (q CQSF)

how it arises from a coalgebra of graphs

Consider graphs on vertices
labelled $\{1, 2, \dots, n\}$,




Coalgebra structure:

$$\Delta_{\{2,4\},\{1,3,5\}}(x) = \bullet 1 \quad \bullet 2 \otimes \begin{array}{c} \text{graph with vertices } 1, 2, 3 \\ \text{edges } (1,2), (2,3), (1,3) \end{array}$$

Example: q -chromatic quasisymmetric function (q CQSF)

how it arises from a coalgebra of graphs

Consider graphs on vertices labelled $\{1, 2, \dots, n\}$,



Coalgebra structure:

$$\Delta_{\{2,4\},\{1,3,5\}}(x) = \bullet_1 \bullet_2 \otimes \begin{array}{ccc} & \text{---} & \\ \bullet_1 & & \bullet_2 & \text{---} & \bullet_3 \\ & \text{---} & \end{array}$$


$$\Delta_{\{1,2,4\},\{3,5\}}(x) = \bullet_1 \bullet_2 \bullet_3 \otimes \begin{array}{cc} \bullet_1 & \bullet_2 \\ \uparrow & \uparrow \\ & \end{array}$$

both are disconnected graphs
 \therefore contributes $x_1^3 x_2^2$ to CQSF

Example: q -chromatic quasisymmetric function (q CQSF)

how it arises from a coalgebra of graphs

Consider graphs on vertices
labelled $\{1, 2, \dots, n\}$,



Coalgebra structure:


Define $\gamma_{S,T}(x)$ = number of edges $s - t$ in x with $s < t$.
Add $q^{\gamma_{S,T}(x)}$ as a coefficient in $\Delta_{S,T}(x)$.

$$\Delta_{\{2,4\},\{1,3,5\}}(x) = q \bullet 1 \quad \bullet 2 \otimes$$

Example: q -chromatic quasisymmetric function (q CQSF)

how it arises from a coalgebra of graphs

Consider graphs on vertices labelled $\{1, 2, \dots, n\}$,



Coalgebra structure:

Define $\gamma_{S,T}(x)$ = number of edges $s - t$ in x with $s < t$.
 Add $q^{\gamma_{S,T}}(x)$ as a coefficient in $\Delta_{S,T}(x)$.

$$\Delta_{\{2,4\},\{1,3,5\}}(x) = q \bullet 1 \quad \bullet 2 \otimes \begin{array}{ccc} & & \\ \bullet 1 & \text{---} & \bullet 2 \\ & \text{---} & \bullet 3 \end{array}$$

$$\Delta_{\{1,2,4\},\{3,5\}}(x) = q^3 \bullet 1 \quad \bullet 2 \quad \bullet 3 \otimes \begin{array}{cc} & \\ \bullet 1 & \bullet 2 \end{array}$$

both are disconnected graphs
 \therefore contributes $q^3 x_1^3 x_2^2$ to CQSF