

Enumerative geometry and quantum integrable systems

Anton M. Zeitlin

Louisiana State University, Department of Mathematics

Louisiana State University

Baton Rouge

December, 2017



We will talk about the relationship between two seemingly independent areas of mathematics:

- ▶ Integrable systems based on quantum groups

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- ▶ Enumerative geometry: quantum equivariant K-theory

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

[Outline](#)

[Quantum Integrability](#)

[Nekrasov-Shatashvili ideas](#)

[Quantum K-theory](#)

[Many-body systems](#)

[Further directions](#)

We will talk about the relationship between two seemingly independent areas of mathematics:

- ▶ **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- ▶ **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- ▶ **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- ▶ **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

We will talk about the relationship between two seemingly independent areas of mathematics:

- **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

[Outline](#)

[Quantum Integrability](#)

[Nekrasov-Shatashvili ideas](#)

[Quantum K-theory](#)

[Many-body systems](#)

[Further directions](#)

We will talk about the relationship between two seemingly independent areas of mathematics:

- ▶ **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- ▶ **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

[Outline](#)

[Quantum Integrability](#)

[Nekrasov-Shatashvili ideas](#)

[Quantum K-theory](#)

[Many-body systems](#)

[Further directions](#)

We will talk about the relationship between two seemingly independent areas of mathematics:

- ▶ **Integrable systems based on quantum groups**

Exactly solvable models of statistical physics: spin chains, vertex models

1930s: Hans Bethe: **Bethe ansatz** solution of Heisenberg model

1960-70s: R.J. Baxter, C.N. Young: **Yang-Baxter equation**, **Baxter operator**

1980s: Development of "QISM" by Leningrad school leading to the discovery of **quantum groups** by Drinfeld and Jimbo

Since 1990s: textbook subject and an established area of mathematics and physics.

- ▶ **Enumerative geometry: quantum equivariant K-theory**

Generalization of **quantum cohomology** in the early 2000s by A. Givental, Y.P. Lee and collaborators. Recently big progress in this direction by A. Okounkov and his school.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Path to this relationship:

- First hints: work of Nekrasov and Shatashvili on 3-dimensional gauge theories, now known as **Gauge-Bethe correspondence**:

N. Nekrasov, S. Shatashvili, *Supersymmetric vacua and Bethe ansatz*, arXiv:0901.4744

N. Nekrasov, S. Shatashvili, *Quantum integrability and supersymmetric vacua*, arXiv:0901.4748

- Subsequent work in **geometric representation theory**:

A. Braverman, D. Maulik, A. Okounkov, *Quantum cohomology of the Springer resolution*, Adv. Math. 227 (2011) 421-458

D. Maulik, A. Okounkov, *Quantum Groups and Quantum Cohomology*, arXiv:1211.1287

A. Okounkov, *Lectures on K-theoretic computations in enumerative geometry*, arXiv: arXiv:1512.07363

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Path to this relationship:

- First hints: work of Nekrasov and Shatashvili on 3-dimensional gauge theories, now known as **Gauge-Bethe correspondence**:

N. Nekrasov, S. Shatashvili, *Supersymmetric vacua and Bethe ansatz*, arXiv:0901.4744

N. Nekrasov, S. Shatashvili, *Quantum integrability and supersymmetric vacua*, arXiv:0901.4748

- Subsequent work in **geometric representation theory**:

A. Braverman, D. Maulik, A. Okounkov, *Quantum cohomology of the Springer resolution*, Adv. Math. 227 (2011) 421-458

D. Maulik, A. Okounkov, *Quantum Groups and Quantum Cohomology*, arXiv:1211.1287

A. Okounkov, *Lectures on K-theoretic computations in enumerative geometry*, arXiv: arXiv:1512.07363

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Path to this relationship:

- First hints: work of Nekrasov and Shatashvili on 3-dimensional gauge theories, now known as **Gauge-Bethe correspondence**:

N. Nekrasov, S. Shatashvili, *Supersymmetric vacua and Bethe ansatz*, arXiv:0901.4744

N. Nekrasov, S. Shatashvili, *Quantum integrability and supersymmetric vacua*, arXiv:0901.4748

- Subsequent work in **geometric representation theory**:

A. Braverman, D. Maulik, A. Okounkov, *Quantum cohomology of the Springer resolution*, Adv. Math. 227 (2011) 421-458

D. Maulik, A. Okounkov, *Quantum Groups and Quantum Cohomology*, arXiv:1211.1287

A. Okounkov, *Lectures on K-theoretic computations in enumerative geometry*, arXiv: arXiv:1512.07363

In fact, this is a part of an ambitious program:

Understanding (enumerative) geometry of **symplectic resolutions**:

"Lie algebras of XXI century" (A. Okounkov' 2012)

Important examples: Springer resolution, Hilbert scheme of points in the plane, Hypertoric varieties,...

A large class of symplectic resolutions is provided by Nakajima quiver varieties (simplest subclass: $T^*Gr(k, n)$)

In this talk our main example will be $T^*Gr(k, n)$ and more generally (partial) flag varieties.

In fact, this is a part of an ambitious program:

Understanding (enumerative) geometry of **symplectic resolutions**:

"Lie algebras of XXI century" (A. Okounkov' 2012)

Important examples: Springer resolution, Hilbert scheme of points in the plane, Hypertoric varieties,...

A large class of symplectic resolutions is provided by Nakajima quiver varieties (simplest subclass: $T^*Gr(k, n)$)

In this talk our main example will be $T^*Gr(k, n)$ and more generally (partial) flag varieties.

In fact, this is a part of an ambitious program:

Understanding (enumerative) geometry of **symplectic resolutions**:

"Lie algebras of XXI century" (A. Okounkov' 2012)

Important examples: **Springer resolution**, **Hilbert scheme of points in the plane**, **Hypertoric varieties**,...

A large class of symplectic resolutions is provided by **Nakajima quiver varieties** (simplest subclass: $T^*Gr(k, n)$)

In this talk our main example will be $T^*Gr(k, n)$ and more generally (partial) flag varieties.

In fact, this is a part of an ambitious program:

Understanding (enumerative) geometry of **symplectic resolutions**:

"Lie algebras of XXI century" (A. Okounkov' 2012)

Important examples: **Springer resolution**, **Hilbert scheme of points in the plane**, **Hypertoric varieties**,...

A large class of symplectic resolutions is provided by **Nakajima quiver varieties** (simplest subclass: $T^*Gr(k, n)$)

In this talk our main example will be $T^*Gr(k, n)$ and more generally (partial) flag varieties.

In fact, this is a part of an ambitious program:

Understanding (enumerative) geometry of **symplectic resolutions**:

"Lie algebras of XXI century" (A. Okounkov' 2012)

Important examples: **Springer resolution**, **Hilbert scheme of points in the plane**, **Hypertoric varieties**,...

A large class of symplectic resolutions is provided by **Nakajima quiver varieties** (simplest subclass: $T^*Gr(k, n)$)

In this talk our main example will be $T^*Gr(k, n)$ and more generally (partial) **flag varieties**.

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Quantum groups, quantum integrability and quantum difference
equations

Nekrasov-Shatashvili ideas

Quantum K-theory (via Okounkov) and integrability

Many-Body systems: back to Givental's ideas

Further directions

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Based on:

- ▶ Petr P. Pushkar, Andrey Smirnov, A.Z., *Baxter Q-operator from quantum K-theory*, arXiv:1612.08723
- ▶ Peter Koroteev, Petr P. Pushkar, Andrey Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*, arXiv:1705.10419

Let us consider Lie algebra \mathfrak{g} . The associated loop algebra is $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$.

The following representations form a tensor category of $\hat{\mathfrak{g}}$:

$$V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n),$$

where

- ▶ V_i are representations of \mathfrak{g}
- ▶ a_i are values for *spectral parameter* t

Let us consider Lie algebra \mathfrak{g} . The associated loop algebra is $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$.

The following representations form a tensor category of $\hat{\mathfrak{g}}$:

$$V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n),$$

where

- ▶ V_i are representations of \mathfrak{g}
- ▶ a_i are values for *spectral parameter* t

Quantum group

$$U_{\hbar}(\hat{\mathfrak{g}})$$

is a deformation of $U(\hat{\mathfrak{g}})$, with a **nontrivial intertwiner** $R_{V_1, V_2}(a_1/a_2)$:

$$V_1(a_1) \otimes V_2(a_2)$$



$$V_2(a_2) \otimes V_1(a_1)$$

which is a rational function of a_1, a_2 , satisfying **Yang-Baxter equation**:



The generators of $U_{\hbar}(\hat{\mathfrak{g}})$ emerge as matrix elements of R -matrices (the so-called FRT construction).

[Outline](#)[Quantum Integrability](#)[Nekrasov-Shatashvili ideas](#)[Quantum K-theory](#)[Many-body systems](#)[Further directions](#)

Source of integrability: commuting *transfer matrices*, generating *Baxter algebra* which are weighted traces of

$$\tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} : W(u) \otimes \mathcal{H}_{\text{phys}} \rightarrow W(u) \otimes \mathcal{H}_{\text{phys}}$$

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems


Further directions

Source of integrability: commuting *transfer matrices*, generating *Baxter algebra* which are weighted traces of

$$\tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} : W(u) \otimes \mathcal{H}_{\text{phys}} \rightarrow W(u) \otimes \mathcal{H}_{\text{phys}}$$

over auxiliary $W(u)$ space:

$$T_W(u) = \text{Tr}_{W(u)} \left((Z \otimes 1) \tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} \right)$$



Here $Z \in e^{\mathfrak{h}}$, where $\mathfrak{h} \in \mathfrak{g}$ are diagonal matrices.

Integrability:

$$[T_{W'}(u'), T_W(u)] = 0$$

Special important case:

$W(u)$ can be certain infinite-dimensional representation of Borel subalgebra of $U_h(\hat{\mathfrak{g}})$.

Then the transfer matrix is called *Baxter Q-operator*. Such operators generate all Baxter algebra.

Primary goal for physicists is to diagonalize $T_W(u)$ simultaneously.

Integrability:

$$[T_{w'}(u'), T_w(u)] = 0$$

Special important case:

$W(u)$ can be certain infinite-dimensional representation of Borel subalgebra of $U_h(\hat{\mathfrak{g}})$.

Then the transfer matrix is called *Baxter Q-operator*. Such operators generate all Baxter algebra.

Primary goal for physicists is to diagonalize $T_w(u)$ simultaneously.

Integrability:

$$[T_{w'}(u'), T_w(u)] = 0$$

Special important case:

$W(u)$ can be certain infinite-dimensional representation of Borel subalgebra of $U_h(\hat{\mathfrak{g}})$.

Then the transfer matrix is called *Baxter Q-operator*. Such operators generate all Baxter algebra.

Primary goal for physicists is to diagonalize $T_w(u)$ simultaneously.

Integrability:

$$[T_{w'}(u'), T_w(u)] = 0$$

Special important case:

$W(u)$ can be certain infinite-dimensional representation of Borel subalgebra of $U_h(\hat{\mathfrak{g}})$.

Then the transfer matrix is called *Baxter Q-operator*. Such operators generate all Baxter algebra.

Primary goal for physicists is to diagonalize $T_w(u)$ simultaneously.

$\mathfrak{g} = \mathfrak{sl}(2)$: XXZ chain

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Textbook example (and main example in this talk) is XXZ Heisenberg spin chain:

$$\mathcal{H}_{\text{XXZ}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

States:

$\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow\downarrow \uparrow\uparrow\uparrow$

Here \mathbb{C}^2 stands for 2-dimensional representation of $U_h(\widehat{\mathfrak{sl}}_2)$.

Algebraic method to diagonalize transfer matrices:

Algebraic Bethe ansatz

as a part of Quantum Inverse Scattering Method developed in the 1980s.

$\mathfrak{g} = \mathfrak{sl}(2)$: XXZ chain

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Textbook example (and main example in this talk) is XXZ Heisenberg spin chain:

$$\mathcal{H}_{\text{XXZ}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

States:

↑↑↑↑ ↓ ↑↑↑ ↓ ↑↑↑↑ ↓ ↑↑↑↑ ↓↓ ↑↑↑

Here \mathbb{C}^2 stands for 2-dimensional representation of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

Algebraic method to diagonalize transfer matrices:

Algebraic Bethe ansatz

as a part of Quantum Inverse Scattering Method developed in the 1980s.

Textbook example (and main example in this talk) is XXZ Heisenberg spin chain:

$$\mathcal{H}_{\text{XXZ}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

States:

$$\uparrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow$$

Here \mathbb{C}^2 stands for 2-dimensional representation of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

Algebraic method to diagonalize transfer matrices:

Algebraic Bethe ansatz

as a part of Quantum Inverse Scattering Method developed in the 1980s.

Bethe equations and Q-operator

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

The eigenvalues are generated by symmetric functions of **Bethe roots** $\{x_i\}$:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \cdots k,$$

so that the eigenvalues $\Lambda(u)$ of the Q -operator are the generating function for the elementary symmetric functions of Bethe roots:

$$\Lambda(u) = \prod_{i=1}^k (1 + u \cdot x_i)$$

A real challenge is to do it for general \mathfrak{g} (possibly infinite-dimensional).

The eigenvalues are generated by symmetric functions of **Bethe roots** $\{x_i\}$:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \cdots k,$$

so that the eigenvalues $\Lambda(u)$ of the **Q**-operator are the generating function for the elementary symmetric functions of Bethe roots:

$$\Lambda(u) = \prod_{i=1}^k (1 + u \cdot x_i)$$

A real challenge is to do it for general \mathfrak{g} (possibly infinite-dimensional).

The eigenvalues are generated by symmetric functions of **Bethe roots** $\{x_i\}$:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \cdots k,$$

so that the eigenvalues $\Lambda(u)$ of the **Q**-operator are the generating function for the elementary symmetric functions of Bethe roots:

$$\Lambda(u) = \prod_{i=1}^k (1 + u \cdot x_i)$$

A real challenge is to do it for general **g** (possibly infinite-dimensional).

Modern way of looking at Bethe ansatz: solving q-difference equations for

$$\Psi(z_1, \dots, z_k; a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)[[z_1, \dots, z_k]]$$

known as quantum Knizhnik-Zamolodchikov (aka Frenkel-Reshetikhin) equations:

$$\Psi(qa_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$

+
commuting difference equations in z - variables

Here $\{z_i\}$ are the components of twist variable Z .

The latter series of equations are known as dynamical equations, studied by Etingof, Felder, Tarasov, Varchenko, ...

In $q \rightarrow 1$ limit we arrive to an eigenvalue problem.

[Outline](#)[Quantum Integrability](#)[Nekrasov-Shatashvili ideas](#)[Quantum K-theory](#)[Many-body systems](#)[Further directions](#)

Modern way of looking at Bethe ansatz: solving q-difference equations for

$$\Psi(z_1, \dots, z_k; a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)[[z_1, \dots, z_k]]$$

known as quantum Knizhnik-Zamolodchikov (aka Frenkel-Reshetikhin) equations:

$$\Psi(qa_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$

+

commuting difference equations in z – variables

Here $\{z_i\}$ are the components of twist variable Z .

The latter series of equations are known as dynamical equations, studied by Etingof, Felder, Tarasov, Varchenko, ...

In $q \rightarrow 1$ limit we arrive to an eigenvalue problem.

Modern way of looking at Bethe ansatz: solving q-difference equations for

$$\Psi(z_1, \dots, z_k; a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)[[z_1, \dots, z_k]]$$

known as quantum Knizhnik-Zamolodchikov (aka Frenkel-Reshetikhin) equations:

$$\Psi(qa_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi +$$

commuting difference equations in z – variables

Here $\{z_i\}$ are the components of twist variable Z .

The latter series of equations are known as dynamical equations, studied by Etingof, Felder, Tarasov, Varchenko, ...

In $q \rightarrow 1$ limit we arrive to an eigenvalue problem.

Modern way of looking at Bethe ansatz: solving q-difference equations for

$$\Psi(z_1, \dots, z_k; a_1, \dots, a_n) \in V_1(a_1) \otimes \dots \otimes V_n(a_n)[[z_1, \dots, z_k]]$$

known as quantum Knizhnik-Zamolodchikov (aka Frenkel-Reshetikhin) equations:

$$\Psi(qa_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi +$$

commuting difference equations in z – variables

Here $\{z_i\}$ are the components of twist variable Z .

The latter series of equations are known as dynamical equations, studied by Etingof, Felder, Tarasov, Varchenko, ...

In $q \rightarrow 1$ limit we arrive to an eigenvalue problem.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as Yang-Yang functional and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}}$ are the eigenvalues of qKZ operators

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as Yang-Yang functional and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}}$ are the eigenvalues of qKZ operators

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as Yang-Yang functional and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}}$ are the eigenvalues of qKZ operators

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as **Yang-Yang functional** and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}} \text{ are the eigenvalues of qKZ operators}$$

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as Yang-Yang functional and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}}$ are the eigenvalues of qKZ operators

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Solutions of qKZ equations and Bethe ansatz

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

$$\Psi_\mu = \int F_\mu(\{x_i\}, \{a_i\}) \mathcal{K}(\{x_i\}, \{a_i\}, \{z_i\}, q) \{dx_i\}$$

Here index μ runs through the Hilbert space $\mathcal{H} = V_1(a_1) \otimes \cdots \otimes V_n(a_n)$

These were studied extensively by Tarasov and Varchenko in the case of $\mathfrak{g} = \mathfrak{gl}_n$.

Here F_μ are rational functions, of its variables and \mathcal{K} is a fixed kernel, such that:

$$\mathcal{K} \sim e^{\frac{\mathcal{S}(\{x_i\}, \{a_i\}, \{z_i\})}{\ln(q)} + \dots}$$

where \mathcal{S} is known as Yang-Yang functional and

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \text{ are Bethe equations for Bethe roots } \{x_i\}$$

$e^{a_i \frac{\partial \mathcal{S}}{\partial a_i}}$ are the eigenvalues of qKZ operators

In other words, $\mu \rightarrow F_\mu$ (off-shell Bethe function) is the diagonalization procedure.

Outline

Quantum Integrability

**Nekrasov-Shatashvili
ideas**

Quantum K-theory

Many-body systems

Further directions

Nekrasov and Shatashvili looked in 2009 at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



with gauge group

$$G = U(v_1) \times U(v_2) \times \dots U(v_{\text{rank } \mathfrak{g}}),$$

so that the collection $\{v_i\}$ determines the weights of the corresponding subspace in \mathcal{H} .

In the simplest case of $\mathfrak{g} = \mathfrak{sl}(2)$ we just have one $U(v)$ and

$$\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow, \text{ and } \# \downarrow = v$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Nekrasov and Shatashvili looked in 2009 at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



with gauge group

$$G = U(v_1) \times U(v_2) \times \dots U(v_{\text{rank } \mathfrak{g}}),$$

so that the collection $\{v_i\}$ determines the weights of the corresponding subspace in \mathcal{H} .

In the simplest case of $\mathfrak{g} = \mathfrak{sl}(2)$ we just have one $U(v)$ and

$$\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow, \text{ and } \# \downarrow = v$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Nekrasov and Shatashvili looked in 2009 at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



with gauge group

$$G = U(v_1) \times U(v_2) \times \dots U(v_{\text{rank } \mathfrak{g}}),$$

so that the collection $\{v_i\}$ determines the weights of the corresponding subspace in \mathcal{H} .

In the simplest case of $\mathfrak{g} = \mathfrak{sl}(2)$ we just have one $U(v)$ and

$$\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow, \text{ and } \# \downarrow = v$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Nekrasov and Shatashvili looked in 2009 at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



with gauge group

$$G = U(v_1) \times U(v_2) \times \dots U(v_{\text{rank } \mathfrak{g}}),$$

so that the collection $\{v_i\}$ determines the weights of the corresponding subspace in \mathcal{H} .

In the simplest case of $\mathfrak{g} = \mathfrak{sl}(2)$ we just have one $U(v)$ and

$$\uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow \downarrow \uparrow\uparrow\uparrow\uparrow\uparrow \downarrow\downarrow \uparrow\uparrow\uparrow, \text{ and } \#\downarrow = v$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a *framing* (“*flavor*”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” *quiver* data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a *framing* (“*flavor*”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” *quiver* data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a *framing* (“*flavor*”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” *quiver* data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a *framing* (“*flavor*”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” *quiver* data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank } g})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank } g} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a **framing** (“*flavor*”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.

- ▶ “Bifundamental” *quiver* data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$

The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a **framing** (“**flavor**”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” **quiver** data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$


The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a **framing** (“**flavor**”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” **quiver** data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$



The quiver serves as a “kind of” Dynkin diagram for \mathfrak{g} .

To have enough supersymmetries \oplus duals : $T^*\mathcal{M}$.

Gauge group G : $U(v_1) \times U(v_2) \times \dots U(v_{\text{rank}})$


The set $\{v_i\}$ determines the weight (i.e. number of inverted spins)

Maximal torus: $\{x_{i_1}, \dots, x_{i_{v_i}}\}$ — these are **Bethe roots** variables.

Matter Fields: affine space \mathcal{M}

- ▶ Standard matter fields: $\bigoplus_{i=1}^{\text{rank}} V_i \otimes W_i$, s.t. $\dim(V_i) = v_i$;
 W_i is a **framing** (“**flavor**”) space, where $\mathbb{C}_{a_1}^\times \times \mathbb{C}_{a_2}^\times \times \dots$ act.
- ▶ “Bifundamental” **quiver** data:

$$\bigoplus_{i \rightarrow j} V_i^* \otimes V_j$$



The quiver serves as a “kind of” Dynkin diagram for \mathfrak{g} .

To have enough supersymmetries \oplus duals : $T^*\mathcal{M}$.

Moduli of Higgs vacua \longleftrightarrow Nakajima quiver variety:

$$T^*\mathcal{M} \mathrel{\mathop{////}} G = \mu^{-1}(0) \mathrel{\mathop{//}} G = N$$

where $\mu = 0$ is low energy configuration condition.

Moduli of Higgs vacua \longleftrightarrow Nakajima quiver variety:

$$T^*\mathcal{M} // G = \mu^{-1}(0) // G = N$$

where $\mu = 0$ is low energy configuration condition.

Hilbert space $\mathcal{H} \longleftrightarrow$ Wilson line operators \longleftrightarrow



equivariant K-theory of Nakajima variety.

Known to be a module for the action of a quantum group $U_h(\hat{\mathfrak{g}})$ due to Nakajima.

Moduli of Higgs vacua \longleftrightarrow Nakajima quiver variety:

$$T^*\mathcal{M} // G = \mu^{-1}(0) // G = N$$

where $\mu = 0$ is low energy configuration condition.

Hilbert space $\mathcal{H} \longleftrightarrow$ Wilson line operators \longleftrightarrow



equivariant K-theory of Nakajima variety.

Known to be a module for the action of a quantum group $U_{\hbar}(\hat{\mathfrak{g}})$ due to Nakajima.

Physicists interested in computing SUSY indices for theories $\mathcal{C} \times S^1$

Mathematically those correspond to (very similar to GW curve counting!) weighted equivariant integrals in K-theory of **quasimaps**:

$$\mathcal{C} \xrightarrow{\text{quasimap } f} \text{Nakajima variety } N$$

The weight (Kähler) parameter is $z^{\deg(f)}$, which is exactly twist parameter Z we encountered before.

Physicists interested in computing SUSY indices for theories $\mathcal{C} \times S^1$

Mathematically those correspond to (very similar to GW curve counting!) weighted equivariant integrals in K-theory of **quasimaps**:

$$\mathcal{C} \xrightarrow{\text{quasimap } f} \text{Nakajima variety } \mathcal{N}$$

The weight (Kähler) parameter is $z^{\deg(f)}$, which is exactly twist parameter Z we encountered before.

Physicists interested in computing SUSY indices for theories $\mathcal{C} \times S^1$

Mathematically those correspond to (very similar to GW curve counting!) weighted equivariant integrals in K-theory of **quasimaps**:

$$\mathcal{C} \xrightarrow{\text{quasimap } f} \text{Nakajima variety } \mathcal{N}$$

The weight (Kähler) parameter is $z^{\deg(f)}$, which is exactly twist parameter Z we encountered before.


Physicists interested in computing SUSY indices for theories $\mathcal{C} \times S^1$

Mathematically those correspond to (very similar to GW curve counting!) weighted equivariant integrals in K-theory of **quasimaps**:

$$\mathcal{C} \xrightarrow{\text{quasimap } f} \text{Nakajima variety } \mathcal{N}$$

The weight (Kähler) parameter is $z^{\deg(f)}$, which is exactly twist parameter z we encountered before.

One can think of **quantum K-theory ring**:



Nekrasov and Shatashvili:

Quantum K – theory ring of Nakajima variety =


symmetric polynomials in x_{i_j} / Bethe equations

Nekrasov and Shatashvili:

Quantum K – theory ring of Nakajima variety =
symmetric polynomials in x_{ij} / Bethe equations

Further input by Okounkov:

q – difference equations = qKZ equations + dynamical equations



In the following we will talk about this in the simplest case:

- ▶ Nakajima variety: $N = T^* Gr(k, n)$
- ▶ Quantum Integrable System: $\mathfrak{sl}(2)$ XXZ spin chain.

$$T^*Gr(k, n) = N_{k,n}, \quad \sqcup_k N_{k,n} = N(n).$$

As a Nakajima variety:

$$N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0)_s / GL(V), \text{ where } \mathcal{M} = Hom(V, W)$$

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in K_{GL(V)}(\cdot) = \Lambda(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we introduce a tautological bundle:

$$\tau = T^*\mathcal{M} \times \tau(V) // GL(V)$$

$$T^*Gr(k, n) = N_{k,n}, \quad \sqcup_k N_{k,n} = N(n).$$

As a Nakajima variety:

$$N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0)_s / GL(V), \text{ where } \mathcal{M} = Hom(V, W)$$

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in K_{GL(V)}(\cdot) = \Lambda(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we introduce a tautological bundle:

$$\tau = T^*\mathcal{M} \times \tau(V) // GL(V)$$

$$T^*Gr(k, n) = N_{k,n}, \quad \sqcup_k N_{k,n} = N(n).$$

As a Nakajima variety:

$$N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0)_s / GL(V), \text{ where } \mathcal{M} = Hom(V, W)$$

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in K_{GL(V)}(\cdot) = \Lambda(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we introduce a tautological bundle:

$$\tau = T^*\mathcal{M} \times \tau(V) // GL(V)$$

$$T^*Gr(k, n) = N_{k,n}, \quad \sqcup_k N_{k,n} = N(n).$$

As a Nakajima variety:

$$N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0)_s / GL(V), \text{ where } \mathcal{M} = Hom(V, W)$$

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in K_{GL(V)}(\cdot) = \Lambda(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we introduce a tautological bundle:

$$\tau = T^*\mathcal{M} \times \tau(V) // GL(V)$$

Tori, Fixed points and Bethe roots

Anton Zeitlin

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathfrak{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Denote $\mathcal{A} := \mathbb{Q}(a_1, \dots, a_n, \hbar)$, $R := \mathbb{Z}(a_1, \dots, a_n, \hbar)$, then **localized K-theory** is:

$$K_T(N(n))_{loc} = K_T(N(n)) \otimes_R \mathcal{A} = \sum_{k=0}^n K_T(N_{k,n}) \otimes_R \mathcal{A}$$

is a 2^n -dimensional \mathcal{A} -vector space, spanned by $\mathcal{O}_{\mathfrak{p}}$.

Classical Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Tori, Fixed points and Bethe roots

Anton Zeitlin

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathfrak{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Denote $\mathcal{A} := \mathbb{Q}(a_1, \dots, a_n, \hbar)$, $R := \mathbb{Z}(a_1, \dots, a_n, \hbar)$, then **localized K-theory** is:

$$K_T(N(n))_{loc} = K_T(N(n)) \otimes_R \mathcal{A} = \sum_{k=0}^n K_T(N_{k,n}) \otimes_R \mathcal{A}$$

is a 2^n -dimensional \mathcal{A} -vector space, spanned by $\mathcal{O}_{\mathfrak{p}}$.

Classical Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Denote $\mathcal{A} := \mathbb{Q}(a_1, \dots, a_n, \hbar)$, $R := \mathbb{Z}(a_1, \dots, a_n, \hbar)$, then **localized K-theory** is:

$$K_T(N(n))_{loc} = K_T(N(n)) \otimes_R \mathcal{A} = \sum_{k=0}^n K_T(N_{k,n}) \otimes_R \mathcal{A}$$

is a 2^n -dimensional \mathcal{A} -vector space, spanned by $\mathcal{O}_{\mathbf{p}}$.

Classical Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Tori, Fixed points and Bethe roots

Anton Zeitlin

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Denote $\mathcal{A} := \mathbb{Q}(a_1, \dots, a_n, \hbar)$, $R := \mathbb{Z}(a_1, \dots, a_n, \hbar)$, then **localized K-theory** is:

$$K_T(N(n))_{loc} = K_T(N(n)) \otimes_R \mathcal{A} = \sum_{k=0}^n K_T(N_{k,n}) \otimes_R \mathcal{A}$$

is a 2^n -dimensional \mathcal{A} -vector space, spanned by $\mathcal{O}_{\mathbf{p}}$.

Classical Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Tori, Fixed points and Bethe roots

Anton Zeitlin

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Denote $\mathcal{A} := \mathbb{Q}(a_1, \dots, a_n, \hbar)$, $R := \mathbb{Z}(a_1, \dots, a_n, \hbar)$, then **localized K-theory** is:

$$K_T(N(n))_{loc} = K_T(N(n)) \otimes_R \mathcal{A} = \sum_{k=0}^n K_T(N_{k,n}) \otimes_R \mathcal{A}$$

is a 2^n -dimensional \mathcal{A} -vector space, spanned by $\mathcal{O}_{\mathbf{p}}$.

Classical Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quantum tautological classes and Bethe equations

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

We will use theory of **quasimaps**:

$$\mathcal{C} \dashrightarrow N_{k,n}$$

in order to deform tensor product: $A \circledast B = A \otimes B + \sum_{d=1}^{\infty} A \otimes_d B z^d$.

We will also define quantum tautological classes:

$$\hat{\tau}(z) = \tau + \sum_{d=1}^{\infty} \tau_d z^d \in K_T(N(n))[[z]]$$

Theorem. [P. Pushkar, A. Smirnov, A.Z] The eigenvalues of operators of quantum multiplication by $\hat{\tau}(z)$ are given by the values of the corresponding Laurent polynomials $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \dots k,$$

We will use theory of **quasimaps**:

$$\mathcal{C} \dashrightarrow N_{k,n}$$

in order to deform tensor product: $A \circledast B = A \otimes B + \sum_{d=1}^{\infty} A \otimes_d B z^d$.

We will also define quantum tautological classes:

$$\hat{\tau}(z) = \tau + \sum_{d=1}^{\infty} \tau_d z^d \in K_T(N(n))[[z]]$$

Theorem. [P. Pushkar, A. Smirnov, A.Z] The eigenvalues of operators of quantum multiplication by $\hat{\tau}(z)$ are given by the values of the corresponding Laurent polynomials $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \dots k,$$

Quantum tautological classes and Bethe equations

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

We will use theory of **quasimaps**:

$$\mathcal{C} \dashrightarrow N_{k,n}$$

in order to deform tensor product: $A \circledast B = A \otimes B + \sum_{d=1}^{\infty} A \otimes_d B z^d$.

We will also define quantum tautological classes:

$$\hat{\tau}(z) = \tau + \sum_{d=1}^{\infty} \tau_d z^d \in K_T(N(n))[[z]]$$

Theorem. [P. Pushkar, A. Smirnov, A.Z] The eigenvalues of operators of quantum multiplication by $\hat{\tau}(z)$ are given by the values of the corresponding Laurent polynomials $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \dots k,$$

Quantum tautological classes and Bethe equations

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

We will use theory of **quasimaps**:

$$\mathcal{C} \dashrightarrow N_{k,n}$$

in order to deform tensor product: $A \circledast B = A \otimes B + \sum_{d=1}^{\infty} A \otimes_d B z^d$.

We will also define quantum tautological classes:

$$\hat{\tau}(z) = \tau + \sum_{d=1}^{\infty} \tau_d z^d \in K_T(N(n))[[z]]$$

Theorem. [P. Pushkar, A. Smirnov, A.Z] The eigenvalues of operators of quantum multiplication by $\hat{\tau}(z)$ are given by the values of the corresponding Laurent polynomials $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1 \dots k,$$

K-theoretic realization of quantum groups

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

View $K_T(N(n))$ as $\mathcal{H}_{\text{XXZ}} = \mathbb{C}^2(a_1) \otimes \cdots \otimes \mathbb{C}^2(a_n)$, a tensor product of evaluation representations of $U_h(\widehat{\mathfrak{sl}}(2))$.

Geometric realization of $U_h(\widehat{\mathfrak{sl}}(2))$?

Consider torus $A = \mathbb{C}_a^\times \curvearrowright W$, so that $W = a'W' + a''W''$.

Maulik and Okounkov:

$$K_T(N(n)^A) = K_T(N(n')) \otimes K_T(N(n'')) \xrightarrow{\text{Stab}_\pm} K_T(N(n)),$$

so that

$R(a'/a'') = \text{Stab}_-^{-1} \text{Stab}_+$ is a trigonometric R – matrix

which is a universal R-matrix considered in the corresponding evaluation representations.

K-theoretic realization of quantum groups

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

View $K_T(N(n))$ as $\mathcal{H}_{\text{XZ}} = \mathbb{C}^2(a_1) \otimes \cdots \otimes \mathbb{C}^2(a_n)$, a tensor product of evaluation representations of $U_h(\widehat{\mathfrak{sl}}(2))$.

Geometric realization of $U_h(\widehat{\mathfrak{sl}}(2))$?

Consider torus $A = \mathbb{C}_a^\times \curvearrowright W$, so that $W = a'W' + a''W''$.

Maulik and Okounkov:

$$K_T(N(n)^A) = K_T(N(n')) \otimes K_T(N(n'')) \xrightarrow{\text{Stab}_\pm} K_T(N(n)),$$

so that

$R(a'/a'') = \text{Stab}_-^{-1} \text{Stab}_+$ is a trigonometric R – matrix

which is a universal R-matrix considered in the corresponding evaluation representations.

K-theoretic realization of quantum groups

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

View $K_T(N(n))$ as $\mathcal{H}_{\text{XZ}} = \mathbb{C}^2(a_1) \otimes \cdots \otimes \mathbb{C}^2(a_n)$, a tensor product of evaluation representations of $U_h(\widehat{\mathfrak{sl}}(2))$.

Geometric realization of $U_h(\widehat{\mathfrak{sl}}(2))$?

Consider torus $A = \mathbb{C}_a^\times \curvearrowright W$, so that $W = a'W' + a''W''$.

Maulik and Okounkov:

$$K_T(N(n)^A) = K_T(N(n')) \otimes K_T(N(n'')) \xrightarrow{\text{Stab}_\pm} K_T(N(n)),$$

so that

$R(a'/a'') = \text{Stab}_-^{-1} \text{Stab}_+$ is a trigonometric R – matrix

which is a universal R-matrix considered in the corresponding evaluation representations.

View $K_T(N(n))$ as $\mathcal{H}_{\text{XZ}} = \mathbb{C}^2(a_1) \otimes \cdots \otimes \mathbb{C}^2(a_n)$, a tensor product of evaluation representations of $U_h(\widehat{\mathfrak{sl}}(2))$.

Geometric realization of $U_h(\widehat{\mathfrak{sl}}(2))$?

Consider torus $A = \mathbb{C}_a^\times \curvearrowright W$, so that $W = a'W' + a''W''$.

Maulik and Okounkov:

$$K_T(N(n)^A) = K_T(N(n')) \otimes K_T(N(n'')) \xrightarrow{\text{Stab}_\pm} K_T(N(n)),$$

so that

$R(a'/a'') = \text{Stab}_-^{-1} \text{Stab}_+$ is a trigonometric R – matrix

which is a universal R-matrix considered in the corresponding evaluation representations.

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Universal R-matrix $\in U_{\hbar}(b_-) \otimes U_{\hbar}(b_+)$ is represented by $U_{\hbar}(\hat{\mathfrak{sl}}(2))$ generators, so that

$$T_w(u) = Tr_{w(u)}[(Z \otimes 1)R_{w(u) \otimes \mathcal{H}_{XXZ}}]$$

Universal R-matrix $\in U_{\hbar}(b_-) \otimes U_{\hbar}(b_+)$ is represented by $U_{\hbar}(\hat{\mathfrak{sl}}(2))$ generators, so that

$$T_w(u) = Tr_{w(u)}[(Z \otimes 1)R_{w(u) \otimes \mathcal{H}_{XXZ}}]$$

Prefundamental representations $V_{\pm}(u)$:

(introduced by Bazhanov, Lukyanov, Zamolochikov, see also E. Frenkel and Hernandez)

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Prefundamental representations $V_{\pm}(u)$:

- ▶ $V_{\pm}(u)$ generate the Grothendieck group of evaluation representations.

Universal R-matrix $\in U_{\hbar}(b_-) \otimes U_{\hbar}(b_+)$ is represented by $U_{\hbar}(\hat{\mathfrak{sl}}(2))$ generators, so that

$$T_w(u) = \text{Tr}_{W(u)}[(Z \otimes 1)R_{W(u) \otimes \mathcal{H}_{\text{XYZ}}}]$$

Prefundamental representations $V_{\pm}(u)$:

(introduced by Bazhanov, Lukyanov, Zamolochikov, see also E. Frenkel and Hernandez)

- ▶ $V_{\pm}(u)$ generate the Grothendieck group of evaluation representations.
- ▶ $Q_{+}(u) = T_{V_{+}(u)}$ has eigenvalues $\prod_{i=1}^k (1 + u \cdot x_i)$, where x_i are solutions of the Bethe equations.
- ▶ Same story for $Q_{-}(u) = T_{V_{-}(u)}$, but Bethe equations change:
 $z \rightarrow z^{-1}$
- ▶ Quantum Wronskian relation:

Universal R-matrix $\in U_{\hbar}(b_-) \otimes U_{\hbar}(b_+)$ is represented by $U_{\hbar}(\hat{\mathfrak{sl}}(2))$ generators, so that

$$T_w(u) = Tr_{w(u)}[(Z \otimes 1)R_{w(u) \otimes \mathcal{H}_{XXZ}}]$$

Prefundamental representations $V_{\pm}(u)$:

(introduced by Bazhanov, Lukyanov, Zamolochikov, see also E. Frenkel and Hernandez)

- ▶ $V_{\pm}(u)$ generate the Grothendieck group of evaluation representations.
- ▶ $Q_+(u) = T_{V_+(u)}$ has eigenvalues $\prod_{i=1}^k (1 + u \cdot x_i)$, where x_i are solutions of the Bethe equations.
- ▶ Same story for $Q_-(u) = T_{V_-(u)}$, but Bethe equations change:
 $z \rightarrow z^{-1}$
- ▶ Quantum Wronskian relation:

More details on transfer matrices and Q-operators

Anton Zeitlin

Universal R-matrix $\in U_{\hbar}(b_-) \otimes U_{\hbar}(b_+)$ is represented by $U_{\hbar}(\hat{\mathfrak{sl}}(2))$ generators, so that

$$T_W(u) = \text{Tr}_{W(u)}[(Z \otimes 1)R_{W(u) \otimes \mathcal{H}_{\text{XZX}}}]$$

Prefundamental representations $V_{\pm}(u)$:

(introduced by Bazhanov, Lukyanov, Zamolochikov, see also E. Frenkel and Hernandez)

- ▶ $V_{\pm}(u)$ generate the Grothendieck group of evaluation representations.
- ▶ $Q_+(u) = T_{V_+(u)}$ has eigenvalues $\prod_{i=1}^k (1 + u \cdot x_i)$, where x_i are solutions of the Bethe equations.
- ▶ Same story for $Q_-(u) = T_{V_-(u)}$, but Bethe equations change:
 $z \rightarrow z^{-1}$
- ▶ Quantum Wronskian relation:

$$z K Q_+(\hbar u) Q_-(u) - Q_+(u) Q_-(\hbar u) = (z K - 1) \prod_{i=1}^n (1 + a_i \cdot u)$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q-operator

Anton Zeitlin

Central result of our first paper:

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products::

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1}^\ell,$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_{\hbar}! \prod_{i=1}^m (1 - (-1)^i z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^{\vee}(z)$, where $\tau_u^{\vee} = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^{\vee}$, $\mathcal{V}^{\vee} = \mathcal{W} - \mathcal{V}$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q -operator

Anton Zeitlin

Central result of our first paper:

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e.

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products::

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1}^\ell,$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_{\hbar}! \prod_{i=1}^m (1 - (-1)^n z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^{\vee}(z)$, where $\tau_u^{\vee} = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^{\vee}$, $\mathcal{V}^{\vee} = \mathcal{W} - \mathcal{V}$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q-operator

Anton Zeitlin

Central result of our first paper:

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products::

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1},$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_{\hbar}! \prod_{i=1}^m (1 - (-1)^n z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^{\vee}(z)$, where $\tau_u^{\vee} = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^{\vee}$, $\mathcal{V}^{\vee} = \mathcal{W} - \mathcal{V}$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q-operator

Anton Zeitlin

Central result of our first paper:

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e.

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products::

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1},$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_{\hbar}! \prod_{i=1}^m (1 - (-1)^n z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^{\vee}(z)$, where $\tau_u^{\vee} = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^{\vee}$, $\mathcal{V}^{\vee} = \mathcal{W} - \mathcal{V}$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q-operator

Anton Zeitlin

Central result of our first paper:

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products::

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1},$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_{\hbar}! \prod_{i=1}^m (1 - (-1)^n z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^{\vee}(z)$, where $\tau_u^{\vee} = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^{\vee}$, $\mathcal{V}^{\vee} = \mathcal{W} - \mathcal{V}$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \bigsqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \mathcal{W})$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \bigsqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \mathcal{W})$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \bigsqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \mathcal{W})$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathbb{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(C, \mathcal{M} \oplus \mathcal{M}^* \otimes \mathcal{h})$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$ev_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Quasimap $f: \mathcal{C} \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathcal{C} of rank k .
- ▶ section $f \in H^0(\mathcal{C}, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of quasimaps

$$QM(N_{k,n}) = \text{stable quasimaps to } N_{k,n} / \sim$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Relative quasimaps

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Relative quasimaps

Anton Zeitlin

Resolution, to make evaluation map proper:

$$\begin{array}{ccc} & QM^d(N_{k,n})_{\text{relative } p} & \\ \nearrow & & \searrow \tilde{ev}_p \\ QM^d(N_{k,n})_{\text{nonsing } p} & \xrightarrow{ev_p} & N_{k,n} \end{array}$$

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory


Many-body systems

Further directions

Relative quasimaps

Anton Zeitlin

Resolution, to make evaluation map proper:



That allows the curve to break: emergence of “*accordeons*”:



$$\begin{array}{ccccc}
 p' & \longrightarrow & \mathcal{C}' & \overset{f'}{\dashrightarrow} & N_{k,n} \\
 \downarrow & & \downarrow \pi & & \\
 p & \longrightarrow & \mathcal{C} & &
 \end{array}$$

i) π is a stabilization of (\mathcal{C}', p')

ii) f' : nonsing at p' and nodes of \mathcal{C}'

iii) $\text{Aut}(f')$ is finite

Outline
Quantum Integrability
Nekrasov-Shatashvili ideas
Quantum K-theory
Many-body systems
Further directions

$QM^d(N_{k,n})$ have perfect deformation-obstruction theory:

- If $(\mathcal{V}, \mathcal{W})$ defines quasimap nonsingular at p .

$$T_{(\mathcal{V}, \mathcal{W})}^{\text{vir}} QM^d_{\text{nonsing } p}(N_{k,n}) = H^\bullet(\mathcal{P} \oplus \hbar \mathcal{P}^*) - T_{f(p)} N_{k,n}$$

where $T_{f(p)} N_{k,n}$ is a normalizing term, \mathcal{P} is the polarization bundle on the curve \mathbb{C} :

$$\mathcal{P} = \mathcal{W}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{V}.$$

- Virtual structure sheaf:

$$\hat{\mathcal{O}}_{\text{vir}} = \mathcal{O}_{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2} q^{\deg(\mathcal{P})/2},$$

where $\mathcal{K}_{\text{vir}} = \det^{-1} T^{\text{vir}} QM^d$ is the virtual canonical bundle.

Equivariant parameter $q : \mathbb{C}_q^\times$ acts on \mathbb{C} with stationary points p_1, p_2 so that $T_{p_1} \mathbb{C}$ has character q .

$QM^d(N_{k,n})$ have perfect deformation-obstruction theory:

- If $(\mathcal{V}, \mathcal{W})$ defines quasimap nonsingular at p .

$$T_{(\mathcal{V}, \mathcal{W})}^{\text{vir}} QM^d_{\text{nonsing } p}(N_{k,n}) = H^*(\mathcal{P} \oplus \hbar \mathcal{P}^*) - T_{f(p)} N_{k,n}$$

where $T_{f(p)} N_{k,n}$ is a normalizing term, \mathcal{P} is the polarization bundle on the curve \mathbb{C} :

$$\mathcal{P} = \mathcal{W}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{W}.$$

- Virtual structure sheaf:

$$\hat{\mathcal{O}}_{\text{vir}} = \mathcal{O}_{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2} q^{\deg(\mathcal{P})/2},$$

where $\mathcal{K}_{\text{vir}} = \det^{-1} T^{\text{vir}} QM^d$ is the virtual canonical bundle.

Equivariant parameter $q : \mathbb{C}_q^\times$ acts on \mathbb{C} with stationary points p_1, p_2 so that $T_{p_1} \mathbb{C}$ has character q .

$QM^d(N_{k,n})$ have perfect deformation-obstruction theory:

- ▶ If $(\mathcal{V}, \mathcal{W})$ defines quasimap nonsingular at p .

$$T_{(\mathcal{V}, \mathcal{W})}^{\text{vir}} QM^d_{\text{nonsing } p}(N_{k,n}) = H^\bullet(\mathcal{P} \oplus \hbar \mathcal{P}^*) - T_{f(p)} N_{k,n}$$

where $T_{f(p)} N_{k,n}$ is a normalizing term, \mathcal{P} is the polarization bundle on the curve \mathbb{C} :

$$\mathcal{P} = \mathcal{W}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{W}.$$

- ▶ Virtual structure sheaf:

$$\hat{\mathcal{O}}_{\text{vir}} = \mathcal{O}_{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2} q^{\deg(\mathcal{P})/2},$$

where $\mathcal{K}_{\text{vir}} = \det^{-1} T^{\text{vir}} QM^d$ is the virtual canonical bundle.

Equivariant parameter $q : \mathbb{C}_q^\times$ acts on \mathbb{C} with stationary points p_1, p_2 so that $T_{p_1} \mathbb{C}$ has character q .

$QM^d(N_{k,n})$ have perfect deformation-obstruction theory:

- If $(\mathcal{V}, \mathcal{W})$ defines quasimap nonsingular at p .

$$T_{(\mathcal{V}, \mathcal{W})}^{\text{vir}} QM^d_{\text{nonsing } p}(N_{k,n}) = H^\bullet(\mathcal{P} \oplus \hbar \mathcal{P}^*) - T_{f(p)} N_{k,n}$$

where $T_{f(p)} N_{k,n}$ is a normalizing term, \mathcal{P} is the polarization bundle on the curve \mathbb{C} :

$$\mathcal{P} = \mathcal{W}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{V}.$$

- Virtual structure sheaf:

$$\hat{\mathcal{O}}_{\text{vir}} = \mathcal{O}_{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2} q^{\deg(\mathcal{P})/2},$$

where $\mathcal{K}_{\text{vir}} = \det^{-1} T^{\text{vir}} QM^d$ is the virtual canonical bundle.

Equivariant parameter $q : \mathbb{C}_q^\times$ acts on \mathbb{C} with stationary points p_1, p_2 so that $T_{p_1} \mathbb{C}$ has character q .

- ▶ If $(\mathcal{V}, \mathcal{W})$ defines quasimap nonsingular at p .

where $T_{f(p)}N_{k,n}$ is a normalizing term, \mathcal{P} is the polarization bundle on the curve \mathbb{C} :

$$\mathcal{P} = \mathcal{W}^* \otimes \mathcal{V} - \mathcal{V}^* \otimes \mathcal{W}.$$

- ▶ Virtual structure sheaf:

$$\hat{\mathcal{O}}_{\text{vir}} = \mathcal{O}_{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2} q^{\deg(\mathcal{P})/2},$$

where $\mathcal{K}_{\text{vir}} = \det^{-1} T^{\text{vir}} QM^d$ is the virtual canonical bundle.

Equivariant parameter $q : \mathbb{C}_q^\times$ acts on \mathbb{C} with stationary points p_1, p_2 so that $T_{p_1}\mathbb{C}$ has character q .

Pushforwards and degeneration formula

Anton Zeitlin

How to degenerate curve in a suitable way?

Blowup: $\pi : \mathbf{C} = \text{Bl}_{\mathbb{C},0} \mathbb{C} \times \mathbb{C}_\epsilon \rightarrow \mathbb{C}_\epsilon$, so that $\mathbb{C}_\epsilon = \pi^{-1}(\epsilon) \cong \mathbb{C}$ for $\epsilon \neq 0$

Central fiber is: $\mathbb{C}_0 = \mathbb{C}_{0,1} \cup_c \mathbb{C}_{0,2}$

Avoiding singularities \rightarrow **Degeneration formula**:

$$\chi(QM(\mathbb{C}_\epsilon \rightarrow N_{k,n}), \hat{\mathcal{O}}_{\text{vir}} z^d) = (\mathbf{G}^{-1} \text{ev}_{1,*}(\hat{\mathcal{O}}_{\text{vir}} z^d), \text{ev}_{2,*}(\hat{\mathcal{O}}_{\text{vir}} z^d))$$

Here **pairing** $(\mathcal{F}, \mathcal{G}) := \chi(\mathcal{F} \otimes \mathcal{G} \otimes K^{-1/2})$,

$$\text{ev}_i : QM(\mathbb{C}_{0,i} \rightarrow N_{k,n})_{\text{relative gluing point}} \rightarrow N_{k,n}$$

$$\text{---} = \text{---} \times \text{---} = \text{---} \rangle \mathbf{G}^{-1} (\text{---} .$$

so that \mathbf{G} is a **gluing operator** \longleftrightarrow :

$$\mathbf{G} = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_2, *} \left(QM_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \right) \in K_T(N_{k,n})^{\otimes 2}[[z]]$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Pushforwards and degeneration formula

Anton Zeitlin

How to degenerate curve in a suitable way?

Blowup: $\pi : \mathbf{C} = \mathrm{Bl}_{\mathbf{C},0} \mathbf{C} \times \mathbb{C}_\epsilon \rightarrow \mathbb{C}_\epsilon$, so that $\mathbf{C}_\epsilon = \pi^{-1}(\epsilon) \cong \mathbf{C}$ for $\epsilon \neq 0$

Central fiber is: $\mathbf{C}_0 = \mathbf{C}_{0,1} \cup_c \mathbf{C}_{0,2}$

Avoiding singularities \rightarrow **Degeneration formula**:

$$\chi(QM(\mathbf{C}_\epsilon \rightarrow N_{k,n}), \hat{\mathcal{O}}_{\mathrm{vir}} z^d) = (\mathbf{G}^{-1} \mathrm{ev}_{1,*}(\hat{\mathcal{O}}_{\mathrm{vir}} z^d), \mathrm{ev}_{2,*}(\hat{\mathcal{O}}_{\mathrm{vir}} z^d))$$

Here **pairing** $(\mathcal{F}, \mathcal{G}) := \chi(\mathcal{F} \otimes \mathcal{G} \otimes K^{-1/2})$,

$$\mathrm{ev}_i : QM(\mathbf{C}_{0,i} \rightarrow N_{k,n})_{\text{relative gluing point}} \rightarrow N_{k,n}$$

$$\text{---} = \text{---} \times \text{---} = \text{---} \rangle \mathbf{G}^{-1} (\text{---} .$$

so that \mathbf{G} is a **gluing operator** \longleftrightarrow :

$$\mathbf{G} = \sum_{d=0}^{\infty} z^d \mathrm{ev}_{p_1, p_2, *} \left(QM_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\mathrm{vir}} \right) \in K_T(N_{k,n})^{\otimes 2}[[z]]$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Pushforwards and degeneration formula

Anton Zeitlin

How to degenerate curve in a suitable way?

Blowup: $\pi : \mathbf{C} = \text{Bl}_{c,0} \mathbb{C} \times \mathbb{C}_\epsilon \rightarrow \mathbb{C}_\epsilon$, so that $\mathbb{C}_\epsilon = \pi^{-1}(\epsilon) \cong \mathbb{C}$ for $\epsilon \neq 0$

Central fiber is: $\mathbb{C}_0 = \mathbb{C}_{0,1} \cup_c \mathbb{C}_{0,2}$

Avoiding singularities \rightarrow Degeneration formula:

$$\chi(QM(\mathbb{C}_\epsilon \rightarrow N_{k,n}), \hat{\mathcal{O}}_{\text{vir}} z^d) = (\mathbf{G}^{-1} \text{ev}_{1,*}(\hat{\mathcal{O}}_{\text{vir}} z^d), \text{ev}_{2,*}(\hat{\mathcal{O}}_{\text{vir}} z^d))$$

Here pairing $(\mathcal{F}, \mathcal{G}) := \chi(\mathcal{F} \otimes \mathcal{G} \otimes K^{-1/2})$,

$$\text{ev}_i : QM(\mathbb{C}_{0,i} \rightarrow N_{k,n})_{\text{relative gluing point}} \rightarrow N_{k,n}$$

$$\text{---} = \text{---} \times \text{---} = \text{---} \rangle \mathbf{G}^{-1} (\text{---} .$$

so that \mathbf{G} is a gluing operator \longleftrightarrow :

$$\mathbf{G} = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_2, *} \left(QM_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \right) \in K_T(N_{k,n})^{\otimes 2}[[z]]$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Pushforwards and degeneration formula

Anton Zeitlin

How to degenerate curve in a suitable way?

Blowup: $\pi : \mathbf{C} = \text{Bl}_{\mathbb{C},0} \mathbb{C} \times \mathbb{C}_\epsilon \rightarrow \mathbb{C}_\epsilon$, so that $\mathbb{C}_\epsilon = \pi^{-1}(\epsilon) \cong \mathbb{C}$ for $\epsilon \neq 0$

Central fiber is: $\mathbb{C}_0 = \mathbb{C}_{0,1} \cup_c \mathbb{C}_{0,2}$

Avoiding singularities \rightarrow Degeneration formula:

$$\chi(QM(\mathbb{C}_\epsilon \rightarrow N_{k,n}), \hat{\mathcal{O}}_{\text{vir}} z^d) = (\mathbf{G}^{-1} \text{ev}_{1,*}(\hat{\mathcal{O}}_{\text{vir}} z^d), \text{ev}_{2,*}(\hat{\mathcal{O}}_{\text{vir}} z^d))$$

Here pairing $(\mathcal{F}, \mathcal{G}) := \chi(\mathcal{F} \otimes \mathcal{G} \otimes K^{-1/2})$,

$$\text{ev}_i : QM(\mathbb{C}_{0,i} \rightarrow N_{k,n})_{\text{relative gluing point}} \rightarrow N_{k,n}$$

$$\text{---} = \text{---} \times \text{---} = \text{---} \rangle \mathbf{G}^{-1} (\text{---} .$$

so that \mathbf{G} is a gluing operator \longleftrightarrow :

$$\mathbf{G} = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_2, *} \left(QM_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \right) \in K_T(N_{k,n})^{\otimes 2}[[z]]$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Pushforwards and degeneration formula

Anton Zeitlin

How to degenerate curve in a suitable way?

Blowup: $\pi : \mathbf{C} = \text{Bl}_{\mathbb{C},0} \mathbb{C} \times \mathbb{C}_\epsilon \rightarrow \mathbb{C}_\epsilon$, so that $\mathbb{C}_\epsilon = \pi^{-1}(\epsilon) \cong \mathbb{C}$ for $\epsilon \neq 0$

Central fiber is: $\mathbb{C}_0 = \mathbb{C}_{0,1} \cup_c \mathbb{C}_{0,2}$

Avoiding singularities \rightarrow **Degeneration formula**:

$$\chi(\mathcal{QM}(\mathbb{C}_\epsilon \rightarrow N_{k,n}), \hat{\mathcal{O}}_{\text{vir}} z^d) = (\mathbf{G}^{-1} \text{ev}_{1,*}(\hat{\mathcal{O}}_{\text{vir}} z^d), \text{ev}_{2,*}(\hat{\mathcal{O}}_{\text{vir}} z^d))$$

Here **pairing** $(\mathcal{F}, \mathcal{G}) := \chi(\mathcal{F} \otimes \mathcal{G} \otimes K^{-1/2})$,

$$\text{ev}_i : \mathcal{QM}(\mathbb{C}_{0,i} \rightarrow N_{k,n})_{\text{relative gluing point}} \rightarrow N_{k,n}$$

$$\text{---} = \text{X} = \text{---} \rangle \mathbf{G}^{-1} (\text{---} .$$

so that \mathbf{G} is a **gluing operator** \longleftrightarrow :

$$\mathbf{G} = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_2, *} \left(\mathcal{QM}_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \right) \in K_T(N_{k,n})^{\otimes 2}[[z]]$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quantum multiplication, quantum tautological classes

Anton Zeitlin

We define the **quantum product** by means of the following element in $K_T(N_{k,n})^{\otimes 2}[[z]]$:

$$\mathcal{F} \circledast \cdot = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_3, *} \left(QM^d_{\text{relative } p_1, p_2, p_3}, \text{ev}_{p_2}^* (\mathbf{G}^{-1} \mathcal{F}) \hat{\theta}_{\text{vir}} \right) \mathbf{G}^{-1}$$

represented by

$$\left(\overbrace{\quad}^{\mathbf{G}^{-1} \mathcal{F}} \right) \mathbf{G}^{-1}$$

$QK_T(N_{k,n}) = K_T(N_{k,n})[[z]]$ is a **unital algebra**, so that:

$$\hat{\mathbf{1}}(z) = \mathbf{1} \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\theta}_{\text{vir}} \right)$$

Similarly, one defines **quantum tautological classes**:

$$\hat{\tau}(z) = \tau \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\theta}_{\text{vir}} \tau(\mathcal{V}|_{p_1}) \right)$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quantum multiplication, quantum tautological classes

Anton Zeitlin

We define the **quantum product** by means of the following element in

$K_T(N_{k,n})^{\otimes 2}[[z]]$:

$$\mathcal{F} \circledast \cdot = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_3, *} \left(QM^d_{\text{relative } p_1, p_2, p_3}, \text{ev}_{p_2}^*(\mathbf{G}^{-1} \mathcal{F}) \hat{\mathcal{O}}_{\text{vir}} \right) \mathbf{G}^{-1}$$

represented by

$$\left(\overbrace{\quad}^{\quad} \right)_{\mathbf{G}^{-1} \mathcal{F}} \mathbf{G}^{-1}$$

$QK_T(N_{k,n}) = K_T(N_{k,n})[[z]]$ is a **unital algebra**, so that:

$$\hat{\mathbf{1}}(z) = \mathbf{1} \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\mathcal{O}}_{\text{vir}} \right)$$

Similarly, one defines **quantum tautological classes**:

$$\hat{\tau}(z) = \tau \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\mathcal{O}}_{\text{vir}} \tau(\mathcal{V}|_{p_1}) \right)$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quantum multiplication, quantum tautological classes

Anton Zeitlin

We define the **quantum product** by means of the following element in

$K_T(N_{k,n})^{\otimes 2}[[z]]$:

$$\mathcal{F} \circledast \cdot = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_3, *} \left(QM^d_{\text{relative } p_1, p_2, p_3}, \text{ev}_{p_2}^*(\mathbf{G}^{-1} \mathcal{F}) \hat{\theta}_{\text{vir}} \right) \mathbf{G}^{-1}$$

represented by

$$\left(\overbrace{\quad}^{\mathbf{G}^{-1} \mathcal{F}} \right) \mathbf{G}^{-1}$$

$QK_T(N_{k,n}) = K_T(N_{k,n})[[z]]$ is a **unital algebra**, so that:

$$\hat{\mathbf{1}}(z) = \mathbf{1} \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\theta}_{\text{vir}} \right)$$

Similarly, one defines **quantum tautological classes**:

$$\hat{\tau}(z) = \tau \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\theta}_{\text{vir}} \tau(\mathcal{V}|_{p_1}) \right)$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quantum multiplication, quantum tautological classes

Anton Zeitlin

We define the **quantum product** by means of the following element in

$K_T(N_{k,n})^{\otimes 2}[[z]]$:

$$\mathcal{F} \circledast \cdot = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_3, *} \left(QM^d_{\text{relative } p_1, p_2, p_3}, \text{ev}_{p_2}^*(\mathbf{G}^{-1} \mathcal{F}) \hat{\mathcal{O}}_{\text{vir}} \right) \mathbf{G}^{-1}$$

represented by

$$\left(\overbrace{\quad}^{\quad} \right)_{\mathbf{G}^{-1} \mathcal{F}} \mathbf{G}^{-1}$$

$QK_T(N_{k,n}) = K_T(N_{k,n})[[z]]$ is a **unital algebra**, so that:

$$\hat{\mathbf{1}}(z) = \mathbf{1} \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\mathcal{O}}_{\text{vir}} \right)$$

Similarly, one defines **quantum tautological classes**:

$$\hat{\tau}(z) = \tau \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2, *} \left(QM^d_{\text{relative } p_2}, \hat{\mathcal{O}}_{\text{vir}} \tau(\mathcal{V}|_{p_1}) \right)$$

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Let us talk about $G = T \times \mathbb{C}_q^\times$ -equivariant K-theory.

- ▶ **Vertex**, a class in $K_G(N_{k,n})_{loc}[[z]]$:

$$V^{(\tau)}(z) = \tau \text{---} \circ = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2,*} \left(QM^d_{\text{nonsing } p_2}, \hat{\mathcal{O}}_{\text{vir}} \tau(\gamma|_{p_1}) \right)$$

singular in $q \rightarrow 1$ limit.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Let us talk about $G = T \times \mathbb{C}_q^\times$ -equivariant K-theory.

- ▶ **Vertex**, a class in $K_G(N_{k,n})_{loc}[[z]]$:

$$V^{(\tau)}(z) = \tau \text{---} \circ = \sum_{d=0}^{\infty} z^d \text{ev}_{p_2,*} \left(QM^d_{\text{nonsing } p_2}, \hat{\theta}_{\text{vir}} \tau(\gamma|_{p_1}) \right)$$

singular in $q \rightarrow 1$ limit.

- ▶ **Capped Vertex**, a class in $K_G(N_{k,n})[[z]]$:

$$\hat{V}^{(\tau)}(z) = \tau \bullet \longrightarrow = \sum_{d=0}^{\infty} z^d \operatorname{ev}_{p_2,*} \left(QM^d_{\text{relative } p_2}, \hat{\mathcal{O}}_{\text{vir}} \tau(\gamma|_{p_1}) \right)$$

Therefore, $\lim_{a \rightarrow 1} \hat{V}^{(\tau)}(z) = \hat{\tau}(z)$

Fusion operator is defined as the following class in $K_G^{\otimes 2}(N_{k,n})_{loc}[[z]]$:

$$\Psi(z) = \sum_{d=0}^{\infty} z^d \text{ev}_{p_1, p_2, *} \left(\begin{matrix} QM^d \\ \text{relative } p_1 \\ \text{nonsing } p_2 \end{matrix}, \hat{\theta}_{\text{vir}} \right)$$

q-difference equation

Anton Zeitlin

Fusion relates two types of vertices:

$$\hat{V}^{(\tau)}(z) = \Psi(z) V^{(\tau)}(z)$$
$$\leftarrow \bullet \tau = \leftarrow \circ \circ \bullet \tau$$

Theorem. i) [A. Okounkov] Fusion operator satisfies q-difference equation:

$$\Psi(qz) = M(z) \Psi(z) \mathcal{O}(1)^{-1},$$

where $\mathcal{O}(1)$ is the operator of classical multiplication by the corresponding line bundle and

$$M(z) = \sum_{d=0}^{\infty} z^d \operatorname{ev}_* \left(QM^d_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \det H^\bullet(\mathcal{V} \otimes \pi^*(\mathcal{O}_{p_1})) \right) \mathbf{G}^{-1},$$

where π is a projection from semistable curve $\mathcal{C}' \rightarrow \mathcal{C}$ and \mathcal{O}_{p_1} is a class of point $p_1 \in \mathcal{C}$.

ii) [P. Pushkar, A. Smirnov, A.Z] Under the specialization $q = 1$ the operator $M(z)$ coincides with the operator of quantum multiplication by the quantum line bundle:

$$M(z)|_{q=1} = \widehat{\mathcal{O}(1)}(z).$$

Explicit formula for $M(z)$ was computed by Smirnov and Okounkov.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

q-difference equation

Anton Zeitlin

Fusion relates two types of vertices:

$$\hat{V}^{(\tau)}(z) = \Psi(z) V^{(\tau)}(z)$$
$$\leftarrow \bullet_{\tau} = \leftarrow \circ \circ \circ \bullet_{\tau}$$

Theorem. i) [A. Okounkov] Fusion operator satisfies q-difference equation:

$$\Psi(qz) = M(z) \Psi(z) \mathcal{O}(1)^{-1},$$

where $\mathcal{O}(1)$ is the operator of classical multiplication by the corresponding line bundle and

$$M(z) = \sum_{d=0}^{\infty} z^d \operatorname{ev}_* \left(QM^d_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \det H^*(\mathcal{V} \otimes \pi^*(\mathcal{O}_{p_1})) \right) \mathbf{G}^{-1},$$

where π is a projection from semistable curve $\mathcal{C}' \rightarrow \mathcal{C}$ and \mathcal{O}_{p_1} is a class of point $p_1 \in \mathcal{C}$.

ii) [P. Pushkar, A. Smirnov, A.Z] Under the specialization $q = 1$ the operator $M(z)$ coincides with the operator of quantum multiplication by the quantum line bundle:

$$M(z)|_{q=1} = \widehat{\mathcal{O}(1)}(z).$$

Explicit formula for $M(z)$ was computed by Smirnov and Okounkov.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

q-difference equation

Anton Zeitlin

Fusion relates two types of vertices:

$$\hat{V}^{(\tau)}(z) = \Psi(z) V^{(\tau)}(z)$$

$$\leftarrow \bullet_{\tau} = \leftarrow \circ \circ \circ \bullet_{\tau}$$

Theorem. i) [A. Okounkov] Fusion operator satisfies q-difference equation:

$$\Psi(qz) = M(z) \Psi(z) \mathcal{O}(1)^{-1},$$

where $\mathcal{O}(1)$ is the operator of classical multiplication by the corresponding line bundle and

$$M(z) = \sum_{d=0}^{\infty} z^d \text{ev}_* \left(QM^d_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \det H^{\bullet}(\mathcal{V} \otimes \pi^*(\mathcal{O}_{p_1})) \right) \mathbf{G}^{-1},$$

where π is a projection from semistable curve $\mathcal{C}' \rightarrow \mathcal{C}$ and \mathcal{O}_{p_1} is a class of point $p_1 \in \mathcal{C}$.

ii) [P. Pushkar, A. Smirnov, A.Z] Under the specialization $q = 1$ the operator $M(z)$ coincides with the operator of quantum multiplication by the quantum line bundle:

$$M(z)|_{q=1} = \widehat{\mathcal{O}(1)}(z).$$

Explicit formula for $M(z)$ was computed by Smirnov and Okounkov.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Fusion relates two types of vertices:

$$\hat{V}^{(\tau)}(z) = \Psi(z) V^{(\tau)}(z)$$

$$\leftarrow \bullet_{\tau} = \leftarrow \circ \circ \circ \bullet_{\tau}$$

Theorem. i) [A. Okounkov] Fusion operator satisfies q-difference equation:

$$\Psi(qz) = M(z) \Psi(z) \mathcal{O}(1)^{-1},$$

where $\mathcal{O}(1)$ is the operator of classical multiplication by the corresponding line bundle and

$$M(z) = \sum_{d=0}^{\infty} z^d \text{ev}_* \left(QM^d_{\text{relative } p_1, p_2}, \hat{\mathcal{O}}_{\text{vir}} \det H^{\bullet}(\mathcal{V} \otimes \pi^*(\mathcal{O}_{p_1})) \right) \mathbf{G}^{-1},$$

where π is a projection from semistable curve $\mathcal{C}' \rightarrow \mathcal{C}$ and \mathcal{O}_{p_1} is a class of point $p_1 \in \mathcal{C}$.

ii) [P. Pushkar, A. Smirnov, A.Z] Under the specialization $q = 1$ the operator $M(z)$ coincides with the operator of quantum multiplication by the quantum line bundle:

$$M(z)|_{q=1} = \widehat{\mathcal{O}(1)}(z).$$

Explicit formula for $M(z)$ was computed by Smirnov and Okounkov.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

i) Localization formula implies the following integral formula for the vertex:

$$V_{\mathbf{p}}^{(\tau)}(z) = \frac{1}{2\pi i \alpha_p} \int_{C_p} \prod_{i=1}^k \frac{ds_i}{s_i} e^{-\frac{\ln(z_{\#}) \ln(s_i)}{\ln(q)}} \prod_{i,j=1}^k \frac{\varphi\left(\frac{s_i}{s_j}\right)}{\varphi\left(\frac{q}{h} \frac{s_i}{s_j}\right)} \prod_{i=1}^n \prod_{j=1}^k \frac{\varphi\left(\frac{q}{h} \frac{s_j}{a_j}\right)}{\varphi\left(\frac{s_j}{a_j}\right)} \tau(s_1, \dots, s_k),$$

where $\varphi(x) = \prod_{i=0}^{\infty} (1 - q^i x)$, $z_{\sharp} = (-1)^n \hbar^{n/2} z$, α_p is a normalization parameter.

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

i) Localization formula implies the following integral formula for the vertex:

$$V_p^{(\tau)}(z) = \frac{1}{2\pi i \alpha_p} \int_{C_p} \prod_{i=1}^k \frac{ds_i}{s_i} e^{-\frac{\ln(z_{\#}) \ln(s_i)}{\ln(q)}} \prod_{i,j=1}^k \frac{\varphi\left(\frac{s_i}{s_j}\right)}{\varphi\left(\frac{q}{h} \frac{s_i}{s_j}\right)} \prod_{i=1}^n \prod_{j=1}^k \frac{\varphi\left(\frac{q}{h} \frac{s_j}{a_i}\right)}{\varphi\left(\frac{s_j}{a_i}\right)} \tau(s_1, \dots, s_k),$$

where $\varphi(x) = \prod_{i=0}^{\infty} (1 - q^i x)$, $z_{\#} = (-1)^n h^{n/2} z$, α_p is a normalization parameter.

ii) The eigenvalues $\tau_p(z)$ of $\hat{\tau}(z)$ are labeled by fixed points are given by the following formula:

$$\tau_p(z) = \lim_{q \rightarrow 1} \frac{V_p^{(\tau)}(z)}{V_p^{(1)}(z)} = \tau(x_{i_1}, x_{i_2}, \dots, x_{i_k})$$

where $V_p^{(\tau)}(z)$ are the components of bare vertex in the basis of fixed points and $\{x_{i_r}\}$ are the solutions of Bethe equations.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

The quantum K-theoretic meaning of the Q-operator

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Theorem. [P. Pushkar, A. Smirnov, A.Z.]

- ▶ The quantum multiplication on quantum tautological class corresponding to $\tau_u := \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}$ coincides with Q_+ -operator, i.e

$$\hat{\tau}_u(z) = Q_+(u)$$

- ▶ Explicit universal formulas for quantum products:

$$\widehat{\Lambda^\ell \mathcal{V}}(z) = \Lambda^\ell \mathcal{V} + a_1(z) F_0 \Lambda^{\ell-1} \mathcal{V} E_{-1} + \cdots + a_\ell(z) F_0^\ell E_{-1},$$

$$\text{where } a_m(z) = \frac{(\hbar-1)^m \hbar^{\frac{m(m+1)}{2}} K^m}{(m)_\hbar! \prod_{i=1}^m (1 - (-1)^n z^{-1} \hbar^i K)},$$

where K, F_0, E_{-1} are the generators of $U_\hbar(\widehat{\mathfrak{sl}}_2)$.

- ▶ $Q_-(u) = \widehat{\tau}_u^\vee(z)$, where $\tau_u^\vee = \bigoplus_{m \geq 0} u^m \Lambda^m \mathcal{V}^\vee$, $\mathcal{V}^\vee = \mathcal{W} - \mathcal{V}$.

Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $W_{n-1} \rightarrow V_{n-1}$ and $V_i \rightarrow V_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}\ell_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $W_{n-1} \rightarrow V_{n-1}$ and $V_i \rightarrow V_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}l_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions


Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $W_{n-1} \rightarrow V_{n-1}$ and $V_i \rightarrow V_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}\ell_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions


Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $\mathbf{W}_{n-1} \rightarrow \mathbf{V}_{n-1}$ and $\mathbf{V}_i \rightarrow \mathbf{V}_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}\ell_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions


Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $w_{n-1} \rightarrow v_{n-1}$ and $v_i \rightarrow v_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}\ell_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions


Relation to Many-Body systems: (partial) flags

Anton Zeitlin

Givental and his collaborators (1990s and early 2000s): relation between quantum geometry of flag varieties and many body systems.

Cotangent bundle to partial flag variety is a

Nakajima variety of type A:



Stability condition: maps $W_{n-1} \rightarrow V_{n-1}$ and $V_i \rightarrow V_{i-1}$ are surjective.

The fixed points: chains of subsets $V_1 \subset \dots \subset V_{n-1} \subset W_{n-1}$, where $|V_i| = v_i$, $W_{n-1} = \{a_1, \dots, a_{w_{n-1}}\}$.

Special case when $v_i = i$, $w_{n-1} = n$ is known as complete flag variety, which we denote as $T^*\mathbb{F}\ell_n$.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Quasimaps, quantum K-theory and Baxter operators

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n - 1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n-1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n-1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n-1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n-1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

We have more bundles for quasimaps: \mathcal{V}_i , $i = 1, \dots, n-1$.

Proceeding as before, we obtain

$$Q_i(u) = \sum_{k=0}^{v_i} (-1)^k u^k \widehat{\Lambda^k \mathcal{V}_i}(z)$$

with eigenvalues of the operator $Q_i(u)$ are the following polynomials in u :

$$\Lambda_i(u) = \prod_{k=1}^{v_i} (1 + x_{i,k} \cdot u),$$

so that the coefficients are elementary symmetric functions of the Bethe roots in $x_{i,k}$ for fixed i .

The Bethe equations at hand are the equations for $\mathfrak{gl}(n)$ spin chain.

Q_i and their dual counterparts \tilde{Q}_i satisfy more involved quadratic quantum Wronskian relations.

Relation to Ruijsenaars-Schneider and Toda systems

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili
ideas

Quantum K-theory

Many-body systems

Further directions

Relations between classical **multiparticle systems** and **quantum integrable models** were observed on various levels (Mukhin, Tarasov, Varchenko, Zabrodin, Zotov, ...)

Gaiotto and Koroteev indicated that in the context of Gauge/Bethe correspondence of Nekrasov and Shatashvili.

Quantum K-theory of $T^*\mathbb{F}\ell$ is an algebra of functions on the Lagrangian subvariety in the phase space of Ruijsenaars-Schneider system, corresponding to the critical locus of Yang-Yang functional.

Relations between classical **multiparticle systems** and **quantum integrable models** were observed on various levels (Mukhin, Tarasov, Varchenko, Zabrodin, Zotov, ...)

Gaiotto and Koroteev indicated that in the context of Gauge/Bethe correspondence of Nekrasov and Shatashvili.

Quantum K-theory of $T^*\mathbb{F}\ell$ is an algebra of functions on the Lagrangian subvariety in the phase space of Ruijsenaars-Schneider system, corresponding to the critical locus of Yang-Yang functional.

Relations between classical **multiparticle systems** and **quantum integrable models** were observed on various levels (Mukhin, Tarasov, Varchenko, Zabrodin, Zotov, ...)

Gaiotto and Koroteev indicated that in the context of Gauge/Bethe correspondence of Nekrasov and Shatashvili.

Quantum K-theory of $T^*\mathbb{F}\ell$ is an algebra of functions on the Lagrangian subvariety in the phase space of Ruijsenaars-Schneider system, corresponding to the critical locus of Yang-Yang functional.

Relations between classical **multiparticle systems** and **quantum integrable models** were observed on various levels (Mukhin, Tarasov, Varchenko, Zabrodin, Zotov, ...)

Gaiotto and Koroteev indicated that in the context of Gauge/Bethe correspondence of Nekrasov and Shatashvili.

Quantum K-theory of $T^*\mathbb{F}\ell$ is an algebra of functions on the Lagrangian subvariety in the phase space of Ruijsenaars-Schneider system, corresponding to the critical locus of Yang-Yang functional.

Theorem.[KPSZ] Quantum equivariant K-theory of the cotangent bundle to complete n-flag variety is given by:

$$QK_T(T^*\mathbb{F}\ell_n) = \frac{\mathbb{C}[\zeta_1^{\pm 1}, \dots, \zeta_n^{\pm 1}; a_1^{\pm 1}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}; p_1^{\pm 1}, \dots, p_n^{\pm 1}]}{\{H_r(\zeta_i, p_i, \hbar) = e_r(\alpha_1, \dots, \alpha_n)\}},$$

so that RS Hamiltonians H_r are given by:

$$H_r = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} \prod_{k \in \mathcal{J}} p_k,$$

where $r = 0, 1, \dots, n$, so that $z_i = \frac{\zeta_i}{\zeta_{i+1}}$ for $i = 1, \dots, n-2$, $\alpha_k = \hbar^{n/2} a_k$ and

$$e_r(\alpha_1, \dots, \alpha_n) = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{k \in \mathcal{J}} \alpha_k, \quad p_j = \hbar^{j-\frac{1}{2}} \widehat{N^j V_j}(z) \otimes \widehat{N^{j-1} V^*_{j-1}}(z),$$

In particular,

$$H_1 = \sum_{i=1}^n \prod_{j \neq i} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} p_i, \quad H_n = \prod_{k=1}^n p_k.$$

Anton Zeitlin

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

Theorem.[KPSZ] Quantum equivariant K-theory of the cotangent bundle to complete n-flag variety is given by:

$$QK_T(T^*\mathbb{F}\ell_n) = \frac{\mathbb{C}[\zeta_1^{\pm 1}, \dots, \zeta_n^{\pm 1}; a_1^{\pm 1}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}; p_1^{\pm 1}, \dots, p_n^{\pm 1}]}{\{H_r(\zeta_i, p_i, \hbar) = e_r(\alpha_1, \dots, \alpha_n)\}},$$

so that RS Hamiltonians H_r are given by:

$$H_r = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}| = r}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} \prod_{k \in \mathcal{J}} p_k,$$

In particular,

$$H_1 = \sum_{i=1}^n \prod_{j \neq i}^n \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} p_i, \quad H_n = \prod_{k=1}^n p_k.$$

Theorem.[KPSZ] Quantum equivariant K-theory of the cotangent bundle to complete n -flag variety is given by:

$$QK_T(T^*\mathbb{F}\ell_n) = \frac{\mathbb{C}[\zeta_1^{\pm 1}, \dots, \zeta_n^{\pm 1}; a_1^{\pm 1}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}; p_1^{\pm 1}, \dots, p_n^{\pm 1}]}{\{H_r(\zeta_i, p_i, \hbar) = e_r(\alpha_1, \dots, \alpha_n)\}},$$

so that RS Hamiltonians H_r are given by:

$$H_r = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} \prod_{k \in \mathcal{J}} p_k,$$

where $r = 0, 1, \dots, n$, so that $z_i = \frac{\zeta_i}{\zeta_{i+1}}$ for $i = 1, \dots, n-2$,
 $\alpha_k = \hbar^{n/2} a_k$ and

$$e_r(\alpha_1, \dots, \alpha_n) = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{k \in \mathcal{J}} \alpha_k, \quad p_j = \hbar^{j-\frac{1}{2}} \widehat{N^j V_j}(z) \otimes \widehat{N^{j-1} V_{j-1}^*}(z),$$

In particular,

$$H_1 = \sum_{i=1}^n \prod_{j \neq i} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} p_i, \quad H_n = \prod_{k=1}^n p_k.$$

Theorem.[KPSZ] Quantum equivariant K-theory of the cotangent bundle to complete n -flag variety is given by:

$$QK_T(T^*\mathbb{F}\ell_n) = \frac{\mathbb{C}[\zeta_1^{\pm 1}, \dots, \zeta_n^{\pm 1}; a_1^{\pm 1}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}; p_1^{\pm 1}, \dots, p_n^{\pm 1}]}{\{H_r(\zeta_i, p_i, \hbar) = e_r(\alpha_1, \dots, \alpha_n)\}},$$

so that RS Hamiltonians H_r are given by:

$$H_r = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} \prod_{k \in \mathcal{J}} p_k,$$

where $r = 0, 1, \dots, n$, so that $z_i = \frac{\zeta_i}{\zeta_{i+1}}$ for $i = 1, \dots, n-2$,
 $\alpha_k = \hbar^{n/2} a_k$ and

$$e_r(\alpha_1, \dots, \alpha_n) = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{k \in \mathcal{J}} \alpha_k, \quad p_j = \hbar^{j-\frac{1}{2}} \widehat{N^j V_j}(z) \otimes \widehat{N^{j-1} V^*_{j-1}}(z),$$

In particular,

$$H_1 = \sum_{i=1}^n \prod_{j \neq i} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} p_i, \quad H_n = \prod_{k=1}^n p_k.$$

Theorem.[KPSZ] Quantum equivariant K-theory of the cotangent bundle to complete n -flag variety is given by:

$$QK_T(T^*\mathbb{F}\ell_n) = \frac{\mathbb{C}[\zeta_1^{\pm 1}, \dots, \zeta_n^{\pm 1}; a_1^{\pm 1}, \dots, a_n^{\pm 1}, \hbar^{\pm 1}; p_1^{\pm 1}, \dots, p_n^{\pm 1}]}{\{H_r(\zeta_i, p_i, \hbar) = e_r(\alpha_1, \dots, \alpha_n)\}},$$

so that RS Hamiltonians H_r are given by:

$$H_r = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{\substack{i \in \mathcal{J} \\ j \notin \mathcal{J}}} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} \prod_{k \in \mathcal{J}} p_k,$$

where $r = 0, 1, \dots, n$, so that $z_i = \frac{\zeta_i}{\zeta_{i+1}}$ for $i = 1, \dots, n-2$,
 $\alpha_k = \hbar^{n/2} a_k$ and

$$e_r(\alpha_1, \dots, \alpha_n) = \sum_{\substack{\mathcal{J} \subset \{1, \dots, n\} \\ |\mathcal{J}|=r}} \prod_{k \in \mathcal{J}} \alpha_k, \quad p_j = \hbar^{j-\frac{1}{2}} \widehat{N^j V_j}(z) \otimes \widehat{N^{j-1} V_{j-1}^*}(z),$$

In particular,

$$H_1 = \sum_{i=1}^n \prod_{j \neq i} \frac{\zeta_i \hbar^{-1/2} - \zeta_j \hbar^{1/2}}{\zeta_i - \zeta_j} p_i, \quad H_n = \prod_{k=1}^n p_k.$$

Special limiting procedure $\hbar \rightarrow \infty$ and rescaling Kähler parameters, which correspond to the removal of the fiber contributions in localization theorem leads to the following result [KPSZ]:

$$QK_T(\mathbb{F}\ell_n) = \frac{\mathbb{C}[\mathfrak{z}_1^{\pm 1}, \dots, \mathfrak{z}_n^{\pm 1}; \mathfrak{a}_1^{\pm 1}, \dots, \mathfrak{a}_n^{\pm 1}; \mathfrak{p}_1^{\pm 1}, \dots, \mathfrak{p}_n^{\pm 1}]}{\{H_r^{\text{q-Toda}}(\mathfrak{z}_i, \mathfrak{p}_i) = e_r(\mathfrak{a}_1, \dots, \mathfrak{a}_n)\}},$$

where H_r are given by:

$$H_r^{\text{q-Toda}} = \sum_{\substack{\mathcal{J} = \{i_1 < \dots < i_r\} \\ \mathcal{J} \subset \{1, \dots, n\}}} \prod_{\ell=1}^r \left(1 - \frac{\mathfrak{z}_{i_\ell-1}}{\mathfrak{z}_{i_\ell}}\right)^{1-\delta_{i_\ell-i_{\ell-1}, 1}} \prod_{k \in \mathcal{J}} \mathfrak{p}_k,$$

so, e.g.

$$H_1^{\text{q-Toda}} = \mathfrak{p}_1 + \sum_{i=2}^n \mathfrak{p}_i \left(1 - \frac{\mathfrak{z}_{i-1}}{\mathfrak{z}_i}\right).$$

That limiting procedure

Ruijsenaars – Schneider \rightarrow Toda

was widely discussed in integrable systems community (Etingof; Gerasimov, Lebedev, Oblezin, ...).

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q -difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q -difference Hamiltonians \leftrightarrow q -difference equation for the vertex

Recent work of Zabrodin and Zotov connects q KZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q -operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q -difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q -difference Hamiltonians \leftrightarrow q -difference equation for the vertex

Recent work of Zabrodin and Zotov connects q KZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q -operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q -difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q -difference Hamiltonians \leftrightarrow q -difference equation for the vertex

Recent work of Zabrodin and Zotov connects q KZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q -operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

Outline

Quantum Integrability

Nekrasov-Shatashvili ideas

Quantum K-theory

Many-body systems

Further directions

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q-difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q-difference Hamiltonians \leftrightarrow q-difference equation for the vertex

Recent work of Zabrodin and Zotov connects qKZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q-operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

- Outline
- Quantum Integrability
- Nekrasov-Shatashvili ideas
- Quantum K-theory
- Many-body systems
- Further directions**

q-difference Hamiltonians \leftrightarrow q-difference equation for the vertex

Geometric meaning of Wronskian relations?

- ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ▶ ↺ 🔍 ↻

- Outline
- Quantum Integrability
- Nekrasov-Shatashvili ideas
- Quantum K-theory
- Many-body systems
- Further directions**

q-difference Hamiltonians \leftrightarrow q-difference equation for the vertex

Geometric meaning of Wronskian relations?

- ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ▶ ↺ 🔍 ↻

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q-difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q-difference Hamiltonians \leftrightarrow q-difference equation for the vertex

Recent work of Zabrodin and Zotov connects qKZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q-operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

- ▶ Original Givental's work: J-functions (vertices) as eigenfunctions of q-difference Toda Hamiltonians

What we obtained so far is $q \rightarrow 1$ limit.

q-difference Hamiltonians \leftrightarrow q-difference equation for the vertex

Recent work of Zabrodin and Zotov connects qKZ equations and eigenfunction problem for RS Hamiltonians. Geometric meaning?

- ▶ Geometric meaning of Wronskian relations?
- ▶ New kinds of integrable systems. Simplest example: Hilbert scheme of points on a plane.
Feigin, Jimbo, Miwa, Mukhin have some recent results on Q-operators.
- ▶ Elliptic quantum groups and Elliptic cohomology
Aganagic, Okounkov laid out foundations
Tarasov, Varchenko recently computed explicitly stable envelopes.

Thank you!