

Hidden Homotopy Symmetries of Einstein Field Equations

Anton M. Zeitlin

Louisiana State University, Department of Mathematics

Louisiana State University

Baton Rouge

February 6, 2019

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations



Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Sigma-models and conformal invariance conditions

Beltrami-Courant differential and first order sigma-models

Vertex/Courant algebroids, G_∞ -algebras and quasiclassical limit

Einstein Equations from G_∞ -algebras

Sigma-models for string theory in curved spacetimes:

Let $X : \Sigma \rightarrow M$, where Σ is a compact Riemann surface (worldsheet) and M is a Riemannian manifold (target space).

Action functional of sigma model:

$$S_{so} = \frac{1}{4\pi h} \int_{\Sigma} (G_{\mu\nu}(X) dX^{\mu} \wedge *dX^{\nu} + X^* B)$$

where G is a metric on M , B is a 2-form on M .

Symmetries:

- i) conformal symmetry on the worldsheet,
- ii) diffeomorphism symmetry and $B \rightarrow B + d\lambda$ on target space.

Sigma-models for string theory in curved spacetimes:

Let $X : \Sigma \rightarrow M$, where Σ is a compact Riemann surface (worldsheet) and M is a Riemannian manifold (target space).

Action functional of sigma model:

$$S_{so} = \frac{1}{4\pi h} \int_{\Sigma} (G_{\mu\nu}(X) dX^{\mu} \wedge *dX^{\nu} + X^* B)$$

where G is a metric on M , B is a 2-form on M .

Symmetries:

- i) conformal symmetry on the worldsheet,
- ii) diffeomorphism symmetry and $B \rightarrow B + d\lambda$ on target space.

Sigma-models for string theory in curved spacetimes:

Let $X : \Sigma \rightarrow M$, where Σ is a compact Riemann surface (worldsheet) and M is a Riemannian manifold (target space).

Action functional of sigma model:

$$S_{so} = \frac{1}{4\pi h} \int_{\Sigma} (G_{\mu\nu}(X) dX^{\mu} \wedge *dX^{\nu} + X^* B)$$

where G is a metric on M , B is a 2-form on M .

Symmetries:

- i) conformal symmetry on the worldsheet,
- ii) diffeomorphism symmetry and $B \rightarrow B + d\lambda$ on target space.

Sigma-models for string theory in curved spacetimes:

Let $X : \Sigma \rightarrow M$, where Σ is a compact Riemann surface (worldsheet) and M is a Riemannian manifold (target space).

Action functional of sigma model:

$$S_{so} = \frac{1}{4\pi h} \int_{\Sigma} (G_{\mu\nu}(X) dX^{\mu} \wedge *dX^{\nu} + X^* B)$$

where G is a metric on M , B is a 2-form on M .

Symmetries:

- i) conformal symmetry on the worldsheet,
- ii) diffeomorphism symmetry and $B \rightarrow B + d\lambda$ on target space.

On the quantum level one can add one more term to the action (due to E. Fradkin and A. Tseytlin):

$$S_{so} \rightarrow S_{so}^{\Phi} = S_{so} + \int_{\Sigma} \Phi(X) R^{(2)}(\gamma) \text{vol}_{\Sigma},$$

where function Φ is called *dilaton*, γ is a metric on Σ .

In order to make sense of path integral

$$Z = \int DX e^{-S_{so}^{\Phi}(X, \gamma)}$$

one has to apply renormalization procedure, so that G , B , Φ depend on certain *cutoff* parameter μ , so that in general quantum theory is not conformally invariant.

On the quantum level one can add one more term to the action (due to E. Fradkin and A. Tseytlin):

$$S_{so} \rightarrow S_{so}^{\Phi} = S_{so} + \int_{\Sigma} \Phi(X) R^{(2)}(\gamma) \text{vol}_{\Sigma},$$

where function Φ is called *dilaton*, γ is a metric on Σ .

In order to make sense of path integral

$$Z = \int DX \ e^{-S_{so}^{\Phi}(X, \gamma)}$$

one has to apply renormalization procedure, so that G , B , Φ depend on certain *cutoff* parameter μ , so that in general quantum theory is not conformally invariant.

Outline

Sigma-models and
conformal invariance
conditionsBeltrami-Courant
differentialVertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

Conformal invariance conditions

$$\mu \frac{d}{d\mu} G_{\mu\nu} = \beta_{\mu\nu}^G(G, B, \Phi, h) = 0, \quad \mu \frac{d}{d\mu} B_{\mu\nu} = \beta_{\mu\nu}^B(G, B, \Phi, h) = 0,$$

$$\mu \frac{d}{d\mu} \Phi = \beta^\Phi(G, B, \Phi, h) = 0$$

at the level h^0 turn out to be Einstein Equations with 2-form field B and dilaton Φ :

$$R_{\mu\nu} = \frac{1}{4} H_\mu^{\lambda\rho} H_{\nu\lambda\rho} - 2\nabla_\mu \nabla_\nu \Phi,$$

$$\nabla^\mu H_{\mu\nu\rho} - 2(\nabla^\lambda \Phi) H_{\lambda\nu\rho} = 0,$$

$$4(\nabla_\mu \Phi)^2 - 4\nabla_\mu \nabla^\mu \Phi + R + \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} = 0,$$

where 3-form $H = dB$, and $R_{\mu\nu}$, R are Ricci and scalar curvature correspondingly.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

In the early days of string theory:

Linearized Einstein Equations and their symmetries:

$(G_{\mu\nu} = \eta_{\mu\nu} + s_{\mu\nu}, B_{\mu\nu} = b_{\mu\nu}, \Phi = \phi)$:

$$Q^\eta \Psi(s, b, \phi) = 0, \quad \Psi^s(s, b, \phi) \rightarrow \Psi(s, b, \phi) + Q^\eta \Lambda$$

in a semi-infinite complex associated to Virasoro module of Hilbert space of states for the "free" theory, associated to flat metric.

It was conjectured (A. Sen, B. Zwiebach,...) in the early 90s that Einstein equations with h -corrections are Generalized Maurer-Cartan (GMC) Equations:

$$Q^\eta \Psi + \frac{1}{2}[\Psi, \Psi]_h + \frac{1}{3!}[\Psi, \Psi, \Psi]_h + \dots = 0$$

$$\Psi \rightarrow \Psi + Q^\eta \Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

where $[\cdot, \cdot, \dots, \cdot]_h$ operations, together with differential Q^η satisfy certain bilinear relations and generate L_∞ -algebra (L stands for Lie).

In the early days of string theory:

Linearized Einstein Equations and their symmetries:

$(G_{\mu\nu} = \eta_{\mu\nu} + s_{\mu\nu}, B_{\mu\nu} = b_{\mu\nu}, \Phi = \phi)$:

$$Q^\eta \Psi(s, b, \phi) = 0, \quad \Psi^s(s, b, \phi) \rightarrow \Psi(s, b, \phi) + Q^\eta \Lambda$$

in a semi-infinite complex associated to Virasoro module of Hilbert space of states for the "free" theory, associated to flat metric.

It was conjectured (A. Sen, B. Zwiebach,...) in the early 90s that Einstein equations with \hbar -corrections are Generalized Maurer-Cartan (GMC) Equations:

$$Q^\eta \Psi + \frac{1}{2}[\Psi, \Psi]_h + \frac{1}{3!}[\Psi, \Psi, \Psi]_h + \dots = 0$$

$$\Psi \rightarrow \Psi + Q^\eta \Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

where $[\cdot, \cdot, \dots, \cdot]_h$ operations, together with differential Q^η satisfy certain bilinear relations and generate L_∞ -algebra (L stands for Lie).

In this talk:

i) Introducing complex structure:

Proper chiral "free action" \rightarrow sheaves of vertex algebras/vertex algebroids.

Metric, B -field \rightarrow Beltrami-Courant differential.

ii) Vertex algebroids $\rightarrow G_\infty$ -algebras (G stands for Gerstenhaber).

Quasiclassical limit:

vertex algebroid \rightarrow Courant algebroid, G_∞ algebra is truncated.

iii) Einstein equations and their \hbar -corrections via Generalized Maurer-Cartan equation for L_∞ -subalgebra of $G_\infty \otimes \bar{G}_\infty$.

Outline

**Sigma-models and
conformal invariance
conditions**

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

In this talk:

i) Introducing complex structure:

Proper chiral "free action" \rightarrow sheaves of vertex algebras/vertex algebroids.

Metric, B -field \rightarrow Beltrami-Courant differential.

ii) Vertex algebroids $\rightarrow G_\infty$ -algebras (G stands for Gerstenhaber).

Quasiclassical limit:

vertex algebroid \rightarrow Courant algebroid, G_∞ algebra is truncated.

iii) Einstein equations and their \hbar -corrections via Generalized Maurer-Cartan equation for L_∞ -subalgebra of $G_\infty \otimes \bar{G}_\infty$.

In this talk:

i) Introducing complex structure:

Proper chiral "free action" \rightarrow sheaves of vertex algebras/vertex algebroids.

Metric, B -field \rightarrow Beltrami-Courant differential.

ii) Vertex algebroids $\rightarrow G_\infty$ -algebras (G stands for Gerstenhaber).

Quasiclassical limit:

vertex algebroid \rightarrow Courant algebroid, G_∞ algebra is truncated.

iii) Einstein equations and their \hbar -corrections via Generalized Maurer-Cartan equation for L_∞ -subalgebra of $G_\infty \otimes \bar{G}_\infty$.

First order version of sigma-model action

We start from the action functional:

$$S_0 = \frac{1}{2\pi i h} \int_{\Sigma} \mathcal{L}_0, \quad \mathcal{L}_0 = \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle,$$

where p, \bar{p} are sections of $X^*(\Omega^{(1,0)}(M)) \otimes \Omega^{(1,0)}(\Sigma)$,
 $X^*(\Omega^{(0,1)}(M)) \otimes \Omega^{(0,1)}(\Sigma)$ correspondingly.

Infinitesimal local symmetries:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_0 + d\xi$$

For holomorphic transformations we have:

$$X^i \rightarrow X^i - v^i(X), \quad X^{\bar{i}} \rightarrow X^{\bar{i}} - \bar{v}^{\bar{i}}(\bar{X}),$$

$$p_i \rightarrow p_i + \partial_i v^k p_k, \quad p_{\bar{i}} \rightarrow p_{\bar{i}} + \partial_{\bar{i}} \bar{v}^{\bar{k}} p_{\bar{k}}$$

$$p_i \rightarrow p_i - \partial X^k (\partial_k \omega_i - \partial_i \omega_k), \quad p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}} (\partial_{\bar{k}} \omega_{\bar{i}} - \partial_{\bar{i}} \omega_{\bar{k}}).$$

Not invariant under general diffeomorphisms, i.e.

$$\delta \mathcal{L}_0 = -\langle \bar{\partial} v, p \wedge \bar{\partial} X \rangle + \langle \partial \bar{v}, \bar{p} \wedge \partial X \rangle.$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

First order version of sigma-model action

We start from the action functional:

$$S_0 = \frac{1}{2\pi i h} \int_{\Sigma} \mathcal{L}_0, \quad \mathcal{L}_0 = \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle,$$

where p, \bar{p} are sections of $X^*(\Omega^{(1,0)}(M)) \otimes \Omega^{(1,0)}(\Sigma)$,
 $X^*(\Omega^{(0,1)}(M)) \otimes \Omega^{(0,1)}(\Sigma)$ correspondingly.

Infinitesimal local symmetries:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_0 + d\xi$$

For holomorphic transformations we have:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X), \quad X^{\bar{i}} \rightarrow X^{\bar{i}} - \bar{v}^{\bar{i}}(\bar{X}), \\ p_i &\rightarrow p_i + \partial_i v^k p_k, \quad p_{\bar{i}} \rightarrow p_{\bar{i}} + \partial_{\bar{i}} \bar{v}^{\bar{k}} p_{\bar{k}} \\ p_i &\rightarrow p_i - \partial X^k (\partial_k \omega_i - \partial_i \omega_k), \quad p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}} (\partial_{\bar{k}} \omega_{\bar{i}} - \partial_{\bar{i}} \omega_{\bar{k}}). \end{aligned}$$

Not invariant under general diffeomorphisms, i.e.

$$\delta \mathcal{L}_0 = -\langle \bar{\partial} v, p \wedge \bar{\partial} X \rangle + \langle \partial \bar{v}, \bar{p} \wedge \partial X \rangle.$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

First order version of sigma-model action

We start from the action functional:

$$S_0 = \frac{1}{2\pi i h} \int_{\Sigma} \mathcal{L}_0, \quad \mathcal{L}_0 = \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle,$$

where p, \bar{p} are sections of $X^*(\Omega^{(1,0)}(M)) \otimes \Omega^{(1,0)}(\Sigma)$,
 $X^*(\Omega^{(0,1)}(M)) \otimes \Omega^{(0,1)}(\Sigma)$ correspondingly.

Infinitesimal local symmetries:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_0 + d\xi$$

For holomorphic transformations we have:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X), \quad X^{\bar{i}} \rightarrow X^{\bar{i}} - \bar{v}^{\bar{i}}(\bar{X}), \\ p_i &\rightarrow p_i + \partial_i v^k p_k, \quad p_{\bar{i}} \rightarrow p_{\bar{i}} + \partial_{\bar{i}} \bar{v}^{\bar{k}} p_{\bar{k}} \\ p_i &\rightarrow p_i - \partial X^k (\partial_k \omega_i - \partial_i \omega_k), \quad p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}} (\partial_{\bar{k}} \omega_{\bar{i}} - \partial_{\bar{i}} \omega_{\bar{k}}). \end{aligned}$$

Not invariant under general diffeomorphisms, i.e.

$$\delta \mathcal{L}_0 = -\langle \bar{\partial} v, p \wedge \bar{\partial} X \rangle + \langle \partial \bar{v}, \bar{p} \wedge \partial X \rangle.$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

First order version of sigma-model action

We start from the action functional:

$$S_0 = \frac{1}{2\pi i h} \int_{\Sigma} \mathcal{L}_0, \quad \mathcal{L}_0 = \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle,$$

where p, \bar{p} are sections of $X^*(\Omega^{(1,0)}(M)) \otimes \Omega^{(1,0)}(\Sigma)$,
 $X^*(\Omega^{(0,1)}(M)) \otimes \Omega^{(0,1)}(\Sigma)$ correspondingly.

Infinitesimal local symmetries:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_0 + d\xi$$

For holomorphic transformations we have:

$$X^i \rightarrow X^i - v^i(X), \quad X^{\bar{i}} \rightarrow X^{\bar{i}} - \bar{v}^{\bar{i}}(\bar{X}),$$

$$p_i \rightarrow p_i + \partial_i v^k p_k, \quad p_{\bar{i}} \rightarrow p_{\bar{i}} + \partial_{\bar{i}} \bar{v}^{\bar{k}} p_{\bar{k}}$$

$$p_i \rightarrow p_i - \partial X^k (\partial_k \omega_i - \partial_i \omega_k), \quad p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}} (\partial_{\bar{k}} \omega_{\bar{i}} - \partial_{\bar{i}} \omega_{\bar{k}}).$$

Not invariant under general diffeomorphisms, i.e.

$$\delta \mathcal{L}_0 = -\langle \bar{\partial} v, p \wedge \bar{\partial} X \rangle + \langle \partial \bar{v}, \bar{p} \wedge \partial X \rangle.$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

It is necessary to add extra terms:

$$\delta \mathcal{L}_\mu = -\langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle,$$

where $\mu \in \Gamma(T^{(1,0)}M \otimes T^{*(0,1)}(M))$, $\bar{\mu} \in \Gamma(T^{(0,1)}M \otimes T^{*(1,0)}(M))$, so that: $\mu \rightarrow \mu - \bar{\partial} v + \dots$, $\bar{\mu} \rightarrow \bar{\mu} - \partial \bar{v} + \dots$.

Continuing the procedure:

$$\begin{aligned} \tilde{\mathcal{L}} = & \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \\ & \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle, \end{aligned}$$

where

$$\begin{aligned} \mu_j^i &\rightarrow \\ \mu_j^i - \partial_{\bar{j}} v^i + v^k \partial_k \mu_j^i + v^{\bar{k}} \partial_{\bar{k}} \mu_j^i + \mu_{\bar{k}}^i \partial_{\bar{j}} v^{\bar{k}} - \mu_j^k \partial_k v^i + \mu_{\bar{l}}^i \mu_j^k \partial_k v^{\bar{l}}, \\ b_{i\bar{j}} &\rightarrow \\ b_{i\bar{j}} + v^k \partial_k b_{i\bar{j}} + v^{\bar{k}} \partial_{\bar{k}} b_{i\bar{j}} + b_{i\bar{k}} \partial_{\bar{j}} v^{\bar{k}} + b_{i\bar{j}} \partial_i v^{\bar{l}} + b_{i\bar{k}} \mu_j^k \partial_k v^{\bar{l}} + b_{i\bar{l}} \bar{\mu}_i^{\bar{k}} \partial_{\bar{k}} v^{\bar{l}}, \end{aligned}$$

so that the transformations of X - and p - fields are:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X, \bar{X}), & p_i &\rightarrow p_i + p_k \partial_i v^k - p_k \mu_{\bar{l}}^k \partial_i v^{\bar{l}} - b_{j\bar{k}} \partial_i v^{\bar{k}} \partial X^j, \\ X^{\bar{i}} &\rightarrow X^{\bar{i}} - v^{\bar{i}}(X, \bar{X}), & \bar{p}_{\bar{i}} &\rightarrow \bar{p}_{\bar{i}} + \bar{p}_{\bar{k}} \partial_{\bar{i}} v^{\bar{k}} - \bar{p}_{\bar{k}} \bar{\mu}_{\bar{l}}^{\bar{k}} \partial_{\bar{i}} v^{\bar{l}} - b_{\bar{j}k} \partial_{\bar{i}} v^k \partial \bar{X}^{\bar{j}}. \end{aligned}$$

It is necessary to add extra terms:

$$\delta \mathcal{L}_\mu = -\langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle,$$

where $\mu \in \Gamma(T^{(1,0)}M \otimes T^{*(0,1)}(M))$, $\bar{\mu} \in \Gamma(T^{(0,1)}M \otimes T^{*(1,0)}(M))$, so that: $\mu \rightarrow \mu - \bar{\partial} v + \dots$, $\bar{\mu} \rightarrow \bar{\mu} - \partial \bar{v} + \dots$.

Continuing the procedure:

$$\begin{aligned} \tilde{\mathcal{L}} = & \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \\ & \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle, \end{aligned}$$

where

$$\begin{aligned} \mu_j^i &\rightarrow \\ \mu_j^i - \partial_{\bar{j}} v^i + v^k \partial_k \mu_j^i + v^{\bar{k}} \partial_{\bar{k}} \mu_j^i + \mu_{\bar{k}}^i \partial_{\bar{j}} v^{\bar{k}} - \mu_j^k \partial_k v^i + \mu_l^i \mu_j^k \partial_k v^{\bar{l}}, \\ b_{i\bar{j}} &\rightarrow \\ b_{i\bar{j}} + v^k \partial_k b_{i\bar{j}} + v^{\bar{k}} \partial_{\bar{k}} b_{i\bar{j}} + b_{i\bar{k}} \partial_{\bar{j}} v^{\bar{k}} + b_{j\bar{l}} \partial_i v^l + b_{i\bar{k}} \mu_j^k \partial_k v^{\bar{k}} + b_{j\bar{l}} \bar{\mu}_i^{\bar{k}} \partial_{\bar{k}} v^l, \end{aligned}$$

so that the transformations of X - and p - fields are:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X, \bar{X}), & p_i &\rightarrow p_i + p_k \partial_i v^k - p_k \mu_l^k \partial_i v^{\bar{l}} - b_{j\bar{k}} \partial_i v^{\bar{k}} \partial X^j, \\ X^{\bar{i}} &\rightarrow X^{\bar{i}} - v^{\bar{i}}(X, \bar{X}), & \bar{p}_{\bar{i}} &\rightarrow \bar{p}_{\bar{i}} + \bar{p}_{\bar{k}} \partial_{\bar{i}} v^{\bar{k}} - \bar{p}_{\bar{k}} \bar{\mu}_l^{\bar{k}} \partial_{\bar{i}} v^l - b_{j\bar{k}} \partial_{\bar{i}} v^k \partial \bar{X}^{\bar{j}}. \end{aligned}$$

It is necessary to add extra terms:

$$\delta \mathcal{L}_\mu = -\langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle,$$

where $\mu \in \Gamma(T^{(1,0)}M \otimes T^{*(0,1)}(M))$, $\bar{\mu} \in \Gamma(T^{(0,1)}M \otimes T^{*(1,0)}(M))$, so that: $\mu \rightarrow \mu - \bar{\partial} v + \dots$, $\bar{\mu} \rightarrow \bar{\mu} - \partial \bar{v} + \dots$.

Continuing the procedure:

$$\begin{aligned} \tilde{\mathcal{L}} = & \langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \\ & \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle, \end{aligned}$$

where

$$\begin{aligned} \mu_j^i &\rightarrow \\ \mu_j^i - \partial_{\bar{j}} v^i + v^k \partial_k \mu_j^i + v^{\bar{k}} \partial_{\bar{k}} \mu_j^i + \mu_{\bar{k}}^i \partial_{\bar{j}} v^{\bar{k}} - \mu_j^k \partial_k v^i + \mu_{\bar{l}}^i \mu_j^k \partial_k v^{\bar{l}}, \\ b_{i\bar{j}} &\rightarrow \\ b_{i\bar{j}} + v^k \partial_k b_{i\bar{j}} + v^{\bar{k}} \partial_{\bar{k}} b_{i\bar{j}} + b_{i\bar{k}} \partial_{\bar{j}} v^{\bar{k}} + b_{i\bar{j}} \partial_i v^{\bar{l}} + b_{i\bar{k}} \mu_j^k \partial_k v^{\bar{l}} + b_{i\bar{j}} \bar{\mu}_i^{\bar{k}} \partial_{\bar{k}} v^{\bar{l}}, \end{aligned}$$

so that the transformations of X - and p - fields are:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X, \bar{X}), & p_i &\rightarrow p_i + p_k \partial_i v^k - p_k \mu_{\bar{l}}^k \partial_i v^{\bar{l}} - b_{j\bar{k}} \partial_i v^{\bar{k}} \partial X^j, \\ X^{\bar{i}} &\rightarrow X^{\bar{i}} - v^{\bar{i}}(X, \bar{X}), & \bar{p}_{\bar{i}} &\rightarrow \bar{p}_{\bar{i}} + \bar{p}_{\bar{k}} \partial_{\bar{i}} v^{\bar{k}} - \bar{p}_{\bar{k}} \bar{\mu}_{\bar{l}}^{\bar{k}} \partial_{\bar{i}} v^{\bar{l}} - b_{j\bar{k}} \partial_{\bar{i}} v^k \partial X^{\bar{j}}. \end{aligned}$$

Similarly, for the 1-form transformation we obtain:

$$b_{i\bar{j}} \rightarrow b_{i\bar{j}} + \partial_{\bar{j}}\omega_i - \partial_i\omega_{\bar{j}} + \mu_{\bar{j}}^i(\partial_i\omega_k - \partial_k\omega_i) + \\ \bar{\mu}_i^{\bar{s}}(\partial_{\bar{j}}\omega_{\bar{s}} - \partial_{\bar{s}}\omega_{\bar{j}}) + \bar{\mu}_{\bar{j}}^{\bar{i}}\mu_{\bar{k}}^s(\partial_s\omega_{\bar{i}} - \partial_{\bar{i}}\omega_s)$$

and

$$p_i \rightarrow p_i - \partial X^k(\partial_k\omega_i - \partial_i\omega_k) - \partial_{\bar{r}}\omega_i\partial X^{\bar{r}} - \bar{\mu}_k^{\bar{s}}\partial_i\omega_{\bar{s}}\partial X^k, \\ p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}}(\partial_{\bar{k}}\omega_{\bar{i}} - \partial_{\bar{i}}\omega_{\bar{k}}) - \partial_r\omega_{\bar{i}}\bar{\partial} X^r - \mu_k^s\partial_i\omega_s\bar{\partial} X^{\bar{k}}.$$

For simplicity:

$$E = TM \oplus T^*M, \quad E = \mathcal{E} \oplus \bar{\mathcal{E}}, \\ \mathcal{E} = T^{(1,0)}M \oplus T^{*(1,0)}M, \quad \bar{\mathcal{E}} = T^{(0,1)}M \oplus T^{*(0,1)}M.$$

Similarly, for the 1-form transformation we obtain:

$$b_{i\bar{j}} \rightarrow b_{i\bar{j}} + \partial_{\bar{j}}\omega_i - \partial_i\omega_{\bar{j}} + \mu_{\bar{j}}^i(\partial_i\omega_k - \partial_k\omega_i) + \\ \bar{\mu}_i^{\bar{s}}(\partial_{\bar{j}}\omega_{\bar{s}} - \partial_{\bar{s}}\omega_{\bar{j}}) + \bar{\mu}_{\bar{j}}^{\bar{i}}\mu_{\bar{k}}^s(\partial_s\omega_{\bar{i}} - \partial_{\bar{i}}\omega_s)$$

and

$$p_i \rightarrow p_i - \partial X^k(\partial_k\omega_i - \partial_i\omega_k) - \partial_{\bar{r}}\omega_i\partial X^{\bar{r}} - \bar{\mu}_k^{\bar{s}}\partial_i\omega_{\bar{s}}\partial X^k, \\ p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}}(\partial_{\bar{k}}\omega_{\bar{i}} - \partial_{\bar{i}}\omega_{\bar{k}}) - \partial_r\omega_{\bar{i}}\bar{\partial} X^r - \mu_k^s\partial_i\omega_s\bar{\partial} X^{\bar{k}}.$$

For simplicity:

$$E = TM \oplus T^*M, \quad E = \mathcal{E} \oplus \bar{\mathcal{E}}, \\ \mathcal{E} = T^{(1,0)}M \oplus T^{*(1,0)}M, \quad \bar{\mathcal{E}} = T^{(0,1)}M \oplus T^{*(0,1)}M.$$

Let $\tilde{M} \in \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$\tilde{M} = \begin{pmatrix} 0 & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

Introduce $\alpha \in \Gamma(E)$, i.e. $\alpha = (v, \bar{v}, \omega, \bar{\omega})$. Let $D : \Gamma(E) \rightarrow \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$D\alpha = \begin{pmatrix} 0 & \bar{\partial}v \\ \partial\bar{v} & \partial\bar{\omega} - \bar{\partial}\omega \end{pmatrix}.$$

Then the transformation of \tilde{M} is:

$$\tilde{M} \rightarrow \tilde{M} - D\alpha + \phi_1(\alpha, \tilde{M}) + \phi_2(\alpha, \tilde{M}, \tilde{M}).$$

Let us describe ϕ_1, ϕ_2 algebraically. In order to do that we need to pass to jet bundles, i.e.

$$\alpha \in J^\infty(\mathcal{O}_M) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}})) \oplus J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}_M),$$

$$\tilde{M} \in J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$$

Let $\tilde{M} \in \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$\tilde{M} = \begin{pmatrix} 0 & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

Introduce $\alpha \in \Gamma(E)$, i.e. $\alpha = (v, \bar{v}, \omega, \bar{\omega})$. Let $D : \Gamma(E) \rightarrow \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$D\alpha = \begin{pmatrix} 0 & \bar{\partial}v \\ \partial\bar{v} & \partial\bar{\omega} - \bar{\partial}\omega \end{pmatrix}.$$

Then the transformation of \tilde{M} is:

$$\tilde{M} \rightarrow \tilde{M} - D\alpha + \phi_1(\alpha, \tilde{M}) + \phi_2(\alpha, \tilde{M}, \tilde{M}).$$

Let us describe ϕ_1, ϕ_2 algebraically. In order to do that we need to pass to jet bundles, i.e.

$$\alpha \in J^\infty(\mathcal{O}_M) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}})) \oplus J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}_M),$$

$$\tilde{M} \in J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$$

Let $\tilde{M} \in \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$\tilde{M} = \begin{pmatrix} 0 & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

Introduce $\alpha \in \Gamma(E)$, i.e. $\alpha = (v, \bar{v}, \omega, \bar{\omega})$. Let $D : \Gamma(E) \rightarrow \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$D\alpha = \begin{pmatrix} 0 & \bar{\partial}v \\ \partial\bar{v} & \partial\bar{\omega} - \bar{\partial}\omega \end{pmatrix}.$$

Then the transformation of \tilde{M} is:

$$\tilde{M} \rightarrow \tilde{M} - D\alpha + \phi_1(\alpha, \tilde{M}) + \phi_2(\alpha, \tilde{M}, \tilde{M}).$$

Let us describe ϕ_1, ϕ_2 algebraically. In order to do that we need to pass to jet bundles, i.e.

$$\alpha \in J^\infty(\mathcal{O}_M) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}})) \oplus J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}_M),$$

$$\tilde{M} \in J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$$

Let $\tilde{\mathbb{M}} \in \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$\tilde{\mathbb{M}} = \begin{pmatrix} 0 & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

Introduce $\alpha \in \Gamma(E)$, i.e. $\alpha = (v, \bar{v}, \omega, \bar{\omega})$. Let $D : \Gamma(E) \rightarrow \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$D\alpha = \begin{pmatrix} 0 & \bar{\partial}v \\ \partial\bar{v} & \partial\bar{\omega} - \bar{\partial}\omega \end{pmatrix}.$$

Then the transformation of $\tilde{\mathbb{M}}$ is:

$$\tilde{\mathbb{M}} \rightarrow \tilde{\mathbb{M}} - D\alpha + \phi_1(\alpha, \tilde{\mathbb{M}}) + \phi_2(\alpha, \tilde{\mathbb{M}}, \tilde{\mathbb{M}}).$$

Let us describe ϕ_1, ϕ_2 algebraically. In order to do that we need to pass to jet bundles, i.e.

$$\alpha \in J^\infty(\mathcal{O}_M) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}})) \oplus J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}_M),$$

$$\tilde{\mathbb{M}} \in J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$$

One can write formally:

$$\alpha = \sum_J f^J \otimes \bar{b}^J + \sum_K b^K \otimes \bar{f}^K,$$

$$\tilde{\mathbb{M}} = \sum_I a^I \otimes \bar{a}^I,$$

where $a^I, b^J \in J^\infty(\mathcal{O}(\mathcal{E}))$, $f^I \in J^\infty(\mathcal{O}_M)$ and $\bar{a}^I, \bar{b}^J \in J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$, $\bar{f}^I \in J^\infty(\bar{\mathcal{O}}_M)$. Then

$$\phi_1(\alpha, \tilde{\mathbb{M}}) = \sum_{I,J} [b^J, a^I]_D \otimes \bar{f}^J \bar{a}^I + \sum_{I,K} f^K a^I \otimes [\bar{b}^K, \bar{a}^I]_D,$$

where $[\cdot, \cdot]_D$ is a *Dorfman bracket*:

$$\begin{aligned} [v_1, v_2]_D &= [v_1, v_2]^{Lie}, \quad [v, \omega]_D = L_v \omega, \\ [\omega, v]_D &= -i_v d\omega, \quad [\omega_1, \omega_2]_D = 0. \end{aligned}$$

Courant bracket is the antisymmetrized version of $[\cdot, \cdot]_D$.

Similarly:

$$\begin{aligned} \phi_2(\alpha, \tilde{\mathbb{M}}, \tilde{\mathbb{M}}) &= \tilde{\mathbb{M}} \cdot D\alpha \cdot \tilde{\mathbb{M}} \\ &= \frac{1}{2} \sum_{I,J,K} \langle b^I, a^K \rangle a^J \otimes \bar{a}^J (\bar{f}^I) \bar{a}^K + \frac{1}{2} \sum_{I,J,K} a^J (f^I) a^K \otimes \langle \bar{b}^I, \bar{a}^K \rangle \bar{a}^J. \end{aligned}$$

One can write formally:

$$\alpha = \sum_J f^J \otimes \bar{b}^J + \sum_K b^K \otimes \bar{f}^K,$$

$$\tilde{\mathbb{M}} = \sum_I a^I \otimes \bar{a}^I,$$

where $a^I, b^J \in J^\infty(\mathcal{O}(\mathcal{E}))$, $f^I \in J^\infty(\mathcal{O}_M)$ and $\bar{a}^I, \bar{b}^J \in J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$, $\bar{f}^I \in J^\infty(\bar{\mathcal{O}}_M)$. Then

$$\phi_1(\alpha, \tilde{\mathbb{M}}) = \sum_{I,J} [b^J, a^I]_D \otimes \bar{f}^J \bar{a}^I + \sum_{I,K} f^K a^I \otimes [\bar{b}^K, \bar{a}^I]_D,$$

where $[\cdot, \cdot]_D$ is a *Dorfman bracket*:

$$[v_1, v_2]_D = [v_1, v_2]^{Lie}, \quad [v, \omega]_D = L_v \omega,$$

$$[\omega, v]_D = -i_v d\omega, \quad [\omega_1, \omega_2]_D = 0.$$

Courant bracket is the antisymmetrized version of $[\cdot, \cdot]_D$.

Similarly:

$$\phi_2(\alpha, \tilde{\mathbb{M}}, \tilde{\mathbb{M}}) = \tilde{\mathbb{M}} \cdot D\alpha \cdot \tilde{\mathbb{M}}$$

$$\frac{1}{2} \sum_{I,J,K} \langle b^I, a^K \rangle a^J \otimes \bar{a}^J (\bar{f}^I) \bar{a}^K + \frac{1}{2} \sum_{I,J,K} a^J (f^I) a^K \otimes \langle \bar{b}^I, \bar{a}^K \rangle \bar{a}^J.$$

One can write formally:

$$\alpha = \sum_J f^J \otimes \bar{b}^J + \sum_K b^K \otimes \bar{f}^K,$$

$$\tilde{\mathbb{M}} = \sum_I a^I \otimes \bar{a}^I,$$

where $a^I, b^J \in J^\infty(\mathcal{O}(\mathcal{E}))$, $f^I \in J^\infty(\mathcal{O}_M)$ and $\bar{a}^I, \bar{b}^J \in J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$, $\bar{f}^I \in J^\infty(\bar{\mathcal{O}}_M)$. Then

$$\phi_1(\alpha, \tilde{\mathbb{M}}) = \sum_{I,J} [b^J, a^I]_D \otimes \bar{f}^J \bar{a}^I + \sum_{I,K} f^K a^I \otimes [\bar{b}^K, \bar{a}^I]_D,$$

where $[\cdot, \cdot]_D$ is a *Dorfman bracket*:

$$[v_1, v_2]_D = [v_1, v_2]^{Lie}, \quad [v, \omega]_D = L_v \omega,$$

$$[\omega, v]_D = -i_v d\omega, \quad [\omega_1, \omega_2]_D = 0.$$

Courant bracket is the antisymmetrized version of $[\cdot, \cdot]_D$.

Similarly:

$$\phi_2(\alpha, \tilde{\mathbb{M}}, \tilde{\mathbb{M}}) = \tilde{\mathbb{M}} \cdot D\alpha \cdot \tilde{\mathbb{M}}$$

$$\frac{1}{2} \sum_{I,J,K} \langle b^I, a^K \rangle a^J \otimes \bar{a}^J (\bar{f}^I) \bar{a}^K + \frac{1}{2} \sum_{I,J,K} a^J (f^I) a^K \otimes \langle \bar{b}^I, \bar{a}^K \rangle \bar{a}^J.$$

Hidden Homotopy Symmetries of Einstein Field Equations

Outline

Sigma-models and conformal invariance conditions

Beltrami-Courant differential

Vertex/Courant algebroids, G_∞ -algebras and quasiclassical limit

Einstein Equations

$$\mathbb{M} = \begin{pmatrix} g & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

$$G \rightarrow G - L_v G, \quad B \rightarrow B - L_v B$$

$$B \rightarrow B - 2d\omega$$

$$\alpha = (\mathbf{v}, \omega), \quad \mathbf{v} \in \Gamma(TM), \omega \in \Omega^1(M)$$

Hidden Homotopy Symmetries of Einstein Field Equations

$$S_{f_0} =$$

Anton Zeitlin

Relation to standard second order sigma-model: Let us fill in 0 in $\tilde{\mathbb{M}}$:

$$\mathbb{M} = \begin{pmatrix} g & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

$$S_{f_0} = \frac{1}{2\pi i h} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \langle g, p \wedge \bar{p} \rangle - \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \bar{p} \wedge \partial X \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle).$$

V.N. Popov, M.G. Zeitlin, Phys.Lett. B 163 (1985) 185, A. Losev, A. Marshakov, A.Z., Phys. Lett. B 633 (2006) 375

Same formulas express symmetries. If $\{g^{ij}\}$ is nondegenerate, then :

$$S_{so} = \frac{1}{4\pi h} \int_{\Sigma} (G_{\mu\nu}(X) dX^{\mu} \wedge *dX^{\nu} + X^* B),$$

$$G_{s\bar{k}} = g_{ij} \bar{\mu}_s^i \mu_{\bar{k}}^j + g_{s\bar{k}} - b_{s\bar{k}}, \quad B_{s\bar{k}} = g_{ij} \bar{\mu}_s^i \mu_{\bar{k}}^j - g_{s\bar{k}} - b_{s\bar{k}}$$

$$G_{si} = -g_{ij}\bar{\mu}_s^j - g_{sj}\bar{\mu}_i^j, \quad G_{\bar{s}\bar{i}} = -g_{\bar{s}j}\mu_{\bar{i}}^j - g_{\bar{i}j}\mu_{\bar{s}}^j$$

$$B_{si} = g_{s\bar{j}}\bar{\mu}_i^{\bar{j}} - g_{ij}\bar{\mu}_s^{\bar{j}}, \quad B_{\bar{s}\bar{i}} = g_{\bar{i}j}\mu_{\bar{s}}^j - g_{\bar{s}j}\mu_{\bar{i}}^j.$$

$$G \rightarrow G - L_v G, \quad B \rightarrow B - L_v B$$

$$B \rightarrow B - 2d\omega$$

$$\alpha = (\mathbf{v}, \omega), \quad \mathbf{v} \in \Gamma(TM), \omega \in \Omega^1(M)$$

Hidden Homotopy Symmetries of Einstein Field Equations

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Relation to standard second order sigma-model: Let us fill in 0 in \tilde{M} :

Sigma-models and conformal invariance conditions

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The quantum theory, corresponding to the chiral part of the free first order Lagrangian \mathcal{L}_0 is described (under certain constraints on M) via sheaves of VOA on M (V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob).

On the open set U of M we have VOA:

$$V = \sum_{n=0}^{\infty} V_n, \quad Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

generated by:

$$\begin{aligned} [X^i(z), p_j(w)] &= h\delta_j^i \delta(z-w), \quad i, j = 1, 2, \dots, D/2 \\ X^i(z) &= \sum_{r \in \mathbb{Z}} X_r^i z^{-r}, \quad p_j(z) = \sum_{s \in \mathbb{Z}} p_{j,s} z^{-s-1} \in \text{End}(V)[[z, z^{-1}]], \end{aligned}$$

so that

$$V = \text{Span}\{p_{j_1, -s_1}, \dots, p_{j_k, -s_k} X_{-r_1}^{i_1} \dots X_{-r_l}^{i_l}\} \otimes F(U) \otimes \mathbb{C}[h, h^{-1}],$$
$$r_m, s_n > 0,$$

$F(U)$ generated by X_0^i -modes.

The quantum theory, corresponding to the chiral part of the free first order Lagrangian \mathcal{L}_0 is described (under certain constraints on M) via sheaves of VOA on M (V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob).

On the open set U of M we have VOA:

$$V = \sum_{n=0}^{\infty} V_n, \quad Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

generated by:

$$[X^i(z), p_j(w)] = h \delta_j^i \delta(z-w), \quad i, j = 1, 2, \dots, D/2$$
$$X^i(z) = \sum_{r \in \mathbb{Z}} X_r^i z^{-r}, \quad p_j(z) = \sum_{s \in \mathbb{Z}} p_{j,s} z^{-s-1} \in \text{End}(V)[[z, z^{-1}]],$$

so that

$$V = \text{Span}\{p_{j_1, -s_1}, \dots, p_{j_k, -s_k} X_{-r_1}^{i_1} \dots X_{-r_l}^{i_l}\} \otimes F(U) \otimes \mathbb{C}[h, h^{-1}],$$
$$r_m, s_n > 0,$$

$F(U)$ generated by X_0^i -modes.

The quantum theory, corresponding to the chiral part of the free first order Lagrangian \mathcal{L}_0 is described (under certain constraints on M) via sheaves of VOA on M (V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob).

On the open set U of M we have VOA:

$$V = \sum_{n=0}^{\infty} V_n, \quad Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

generated by:

$$[X^i(z), p_j(w)] = h \delta_j^i \delta(z-w), \quad i, j = 1, 2, \dots, D/2$$

$$X^i(z) = \sum_{r \in \mathbb{Z}} X_r^i z^{-r}, \quad p_j(z) = \sum_{s \in \mathbb{Z}} p_{j,s} z^{-s-1} \in \text{End}(V)[[z, z^{-1}]],$$

so that

$$V = \text{Span}\{p_{j_1, -s_1}, \dots, p_{j_k, -s_k} X_{-r_1}^{i_1} \dots X_{-r_l}^{i_l}\} \otimes F(U) \otimes \mathbb{C}[h, h^{-1}],$$

$$r_m, s_n > 0,$$

$F(U)$ generated by X_0^i -modes.

The quantum theory, corresponding to the chiral part of the free first order Lagrangian \mathcal{L}_0 is described (under certain constraints on M) via sheaves of VOA on M (V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob).

On the open set U of M we have VOA:

$$V = \sum_{n=0}^{\infty} V_n, \quad Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

generated by:

$$[X^i(z), p_j(w)] = h\delta_j^i \delta(z-w), \quad i, j = 1, 2, \dots, D/2$$

$$X^i(z) = \sum_{r \in \mathbb{Z}} X_r^i z^{-r}, \quad p_j(z) = \sum_{s \in \mathbb{Z}} p_{j,s} z^{-s-1} \in \text{End}(V)[[z, z^{-1}]],$$

so that

$$V = \text{Span}\{p_{j_1, -s_1}, \dots, p_{j_k, -s_k} X_{-r_1}^{i_1} \dots X_{-r_l}^{i_l}\} \otimes F(U) \otimes \mathbb{C}[h, h^{-1}],$$

$$r_m, s_n > 0,$$

$F(U)$ generated by X_0^i -modes.

The quantum theory, corresponding to the chiral part of the free first order Lagrangian \mathcal{L}_0 is described (under certain constraints on M) via sheaves of VOA on M (V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob).

On the open set U of M we have VOA:

$$V = \sum_{n=0}^{\infty} V_n, \quad Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

generated by:

$$[X^i(z), p_j(w)] = h\delta_j^i \delta(z-w), \quad i, j = 1, 2, \dots, D/2$$
$$X^i(z) = \sum_{r \in \mathbb{Z}} X_r^i z^{-r}, \quad p_j(z) = \sum_{s \in \mathbb{Z}} p_{j,s} z^{-s-1} \in \text{End}(V)[[z, z^{-1}]],$$

so that

$$V = \text{Span}\{p_{j_1, -s_1}, \dots, p_{j_k, -s_k} X_{-r_1}^{i_1} \dots X_{-r_l}^{i_l}\} \otimes F(U) \otimes \mathbb{C}[h, h^{-1}],$$
$$r_m, s_n > 0,$$

$F(U)$ generated by X_0^i -modes.

The Virasoro element is:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{D}{12}(n^3 - n)\delta_{n,-m}$$

corresponding to correction:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_{\phi'} = \langle p \wedge \bar{\partial} X \rangle - 2\pi i h R^{(2)}(\gamma) \phi'(X)$$

where $\phi' = \log \Omega$, where $\Omega(X) dX^1 \wedge \cdots \wedge dX^n$ is a holomorphic volume form, i.e. for globally defined $T(z)$, M has to be Calabi-Yau.

The space V is a lowest weight module for the above Virasoro algebra.

V can be reproduced from V_0 and V_1 as a *vertex envelope*. The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebroid*.

In our case: $V_0 \rightarrow \mathcal{O}_U^h = \mathcal{O}_U \otimes \mathbb{C}[h, h^{-1}]$,

$V_1 \rightarrow \mathcal{V}^h = \mathcal{V} \otimes \mathbb{C}[h, h^{-1}]$,

$\mathcal{V} = \mathcal{O}(\mathcal{E}_U)$, generated by $: v_i(X) p_i : , \omega_i(X) \partial X^i$

The Virasoro element is:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{D}{12}(n^3 - n)\delta_{n,-m}$$

corresponding to correction:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_{\phi'} = \langle p \wedge \bar{\partial} X \rangle - 2\pi i h R^{(2)}(\gamma) \phi'(X)$$

where $\phi' = \log \Omega$, where $\Omega(X) dX^1 \wedge \cdots \wedge dX^n$ is a holomorphic volume form, i.e. for globally defined $T(z)$, M has to be Calabi-Yau.

The space V is a lowest weight module for the above Virasoro algebra.

V can be reproduced from V_0 and V_1 as a *vertex envelope*. The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebroid*.

In our case: $V_0 \rightarrow \mathcal{O}_U^h = \mathcal{O}_U \otimes \mathbb{C}[h, h^{-1}]$,

$V_1 \rightarrow \mathcal{V}^h = \mathcal{V} \otimes \mathbb{C}[h, h^{-1}]$,

$\mathcal{V} = \mathcal{O}(\mathcal{E}_U)$, generated by $: v_i(X) p_i : , \omega_i(X) \partial X^i$

The Virasoro element is:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{D}{12}(n^3 - n)\delta_{n,-m}$$

corresponding to correction:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_{\phi'} = \langle p \wedge \bar{\partial} X \rangle - 2\pi i h R^{(2)}(\gamma) \phi'(X)$$

where $\phi' = \log \Omega$, where $\Omega(X) dX^1 \wedge \cdots \wedge dX^n$ is a holomorphic volume form, i.e. for globally defined $T(z)$, M has to be Calabi-Yau.

The space V is a lowest weight module for the above Virasoro algebra.

V can be reproduced from V_0 and V_1 as a *vertex envelope*. The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebroid*.

In our case: $V_0 \rightarrow \mathcal{O}_U^h = \mathcal{O}_U \otimes \mathbb{C}[h, h^{-1}]$,

$V_1 \rightarrow \mathcal{V}^h = \mathcal{V} \otimes \mathbb{C}[h, h^{-1}]$,

$\mathcal{V} = \mathcal{O}(\mathcal{E}_U)$, generated by $: v_i(X) p_i : , \omega_i(X) \partial X^i$

The Virasoro element is:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{D}{12}(n^3 - n)\delta_{n,-m}$$

corresponding to correction:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_{\phi'} = \langle p \wedge \bar{\partial} X \rangle - 2\pi i h R^{(2)}(\gamma) \phi'(X)$$

where $\phi' = \log \Omega$, where $\Omega(X) dX^1 \wedge \cdots \wedge dX^n$ is a holomorphic volume form, i.e. for globally defined $T(z)$, M has to be Calabi-Yau.

The space V is a lowest weight module for the above Virasoro algebra.

V can be reproduced from V_0 and V_1 as a *vertex envelope*. The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebroid*.

In our case: $V_0 \rightarrow \mathcal{O}_U^h = \mathcal{O}_U \otimes \mathbb{C}[h, h^{-1}]$,

$V_1 \rightarrow \mathcal{V}^h = \mathcal{V} \otimes \mathbb{C}[h, h^{-1}]$,

$\mathcal{V} = \mathcal{O}(\mathcal{E}_U)$, generated by $: v_i(X) p_i : , \omega_i(X) \partial X^i$

The Virasoro element is:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{D}{12}(n^3 - n)\delta_{n,-m}$$

corresponding to correction:

$$\mathcal{L}_0 \rightarrow \mathcal{L}_{\phi'} = \langle p \wedge \bar{\partial} X \rangle - 2\pi i h R^{(2)}(\gamma) \phi'(X)$$

where $\phi' = \log \Omega$, where $\Omega(X) dX^1 \wedge \cdots \wedge dX^n$ is a holomorphic volume form, i.e. for globally defined $T(z)$, M has to be Calabi-Yau.

The space V is a lowest weight module for the above Virasoro algebra.

V can be reproduced from V_0 and V_1 as a *vertex envelope*. The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebroid*.

In our case: $V_0 \rightarrow \mathcal{O}_U^h = \mathcal{O}_U \otimes \mathbb{C}[h, h^{-1}]$,

$V_1 \rightarrow \mathcal{V}^h = \mathcal{V} \otimes \mathbb{C}[h, h^{-1}]$,

$\mathcal{V} = \mathcal{O}(\mathcal{E}_U)$, generated by $: v_i(X) p_i : , \omega_i(X) \partial X^i$

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$\begin{aligned}
 f * (g * v) - (fg) * v &= \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f), \\
 [v_1, f * v_2] &= \pi(v_1)(f) * v_2 + f * [v_1, v_2], \\
 [v_1, v_2] + [v_2, v_1] &= \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v), \\
 \langle f * v_1, v_2 \rangle &= f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)), \\
 \pi(v)(\langle v_1, v_2 \rangle) &= \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle, \\
 \partial(fg) &= f * \partial(g) + g * \partial(f), \\
 [v, \partial(f)] &= \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),
 \end{aligned}$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[\cdot, \cdot] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[\cdot, \cdot] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[\cdot, \cdot] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.

ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[\cdot, \cdot] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,

iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor

iv) a symmetric \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,

v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$,

naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$\begin{aligned}
 f * (g * v) - (fg) * v &= \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f), \\
 [v_1, f * v_2] &= \pi(v_1)(f) * v_2 + f * [v_1, v_2], \\
 [v_1, v_2] + [v_2, v_1] &= \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v), \\
 \langle f * v_1, v_2 \rangle &= f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)), \\
 \pi(v)(\langle v_1, v_2 \rangle) &= \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle, \\
 \partial(fg) &= f * \partial(g) + g * \partial(f), \\
 [v, \partial(f)] &= \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),
 \end{aligned}$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with

- i) \mathbb{C} -linear pairing $\mathcal{O}_M \otimes \mathcal{V} \rightarrow \mathcal{V}[h]$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$.
- ii) \mathbb{C} -linear bracket, satisfying Leibniz algebra $[,] : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{V}[h]$,
- iii) \mathbb{C} -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow h\Gamma(TM)[h]$ usually referred to as an anchor
- iv) a symmetric \mathbb{C} -bilinear pairing $\langle , \rangle : \mathcal{V} \otimes \mathcal{V} \rightarrow h\mathcal{O}_M[h]$,
- v) a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, naturally extending to \mathcal{O}_M^h and \mathcal{V}^h , and satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial \langle v_1, v_2 \rangle, \quad \pi(f * v) = f \pi(v),$$

$$\langle f * v_1, v_2 \rangle = f \langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}^h$, $f, g \in \mathcal{O}_M^h$.

For our considerations $\mathcal{V} = \mathcal{O}(\mathcal{E})$:

$$\begin{aligned}\partial f &= df, \quad \pi(v)f = -h\nu(f), \quad \pi(\omega) = 0, \\ f * v &= fv + h dX^i \partial_i \partial_j f v^j, \quad f * \omega = f\omega, \\ [v_1, v_2] &= -h[v_1, v_2]_D - h^2 dX^i \partial_i \partial_k v_1^s \partial_s v_2^k, \\ [v, \omega] &= -h[v, \omega]_D, \quad [\omega, v] = -h[\omega, v]_D, \quad [\omega_1, \omega_2] = 0, \\ \langle v, \omega \rangle &= -h\langle v, \omega \rangle^s, \quad \langle v_1, v_2 \rangle = -h^2 \partial_i v_1^j \partial_j v_2^i, \quad \langle \omega_1, \omega_2 \rangle = 0,\end{aligned}$$

where v and ω are vector fields and 1-forms correspondingly.

Together with $\text{div}_{\phi'}$ -the divergence operator with respect to ϕ' these operations generate vertex algebroid with Calabi-Yau structure.

For our considerations $\mathcal{V} = \mathcal{O}(\mathcal{E})$:

$$\begin{aligned}\partial f &= df, \quad \pi(v)f = -h\nu(f), \quad \pi(\omega) = 0, \\ f * v &= fv + h dX^i \partial_i \partial_j f v^j, \quad f * \omega = f\omega, \\ [v_1, v_2] &= -h[v_1, v_2]_D - h^2 dX^i \partial_i \partial_k v_1^s \partial_s v_2^k, \\ [v, \omega] &= -h[v, \omega]_D, \quad [\omega, v] = -h[\omega, v]_D, \quad [\omega_1, \omega_2] = 0, \\ \langle v, \omega \rangle &= -h\langle v, \omega \rangle^s, \quad \langle v_1, v_2 \rangle = -h^2 \partial_i v_1^j \partial_j v_2^i, \quad \langle \omega_1, \omega_2 \rangle = 0,\end{aligned}$$

where v and ω are vector fields and 1-forms correspondingly.

Together with $\text{div}_{\phi'}$ -the divergence operator with respect to ϕ' these operations generate vertex algebroid with Calabi-Yau structure.

Vertex algebra V is a Virasoro module. The corresponding semi-infinite complex V^{semi} (the analogue of Chevalley complex for Virasoro algebra) is a vertex algebra too:

$$V^{semi} = V \otimes \Lambda,$$

$$\Lambda \text{ generated by } [b(z), c(w)]_+ = \delta(z - w).$$

The corresponding differential

$$Q = j_0, \quad j(z) = \sum_{n \in \mathbb{Z}} j_n z^{-n-1} = c(z)T(z) + :c(z)\partial c(z)b(z):$$

is nilpotent when $D = 26$ (famous dimension 26!). However, we will consider subcomplex of light modes (i.e. $L_0 = 0$) denoted in the following as (\mathcal{F}_h, Q) , where we can drop this condition:

$$\begin{array}{ccccc}
 & \mathcal{V}^h & & \mathcal{V}^h & \\
 \nearrow \partial & & \searrow \partial & \nearrow \partial & \searrow \frac{1}{2}h\text{div} \\
 \mathcal{O}_M^h & \oplus & \mathcal{O}_M^h & \oplus & \mathcal{O}_M^h \\
 & \nwarrow \frac{1}{2}h\text{div} & \nearrow \text{id} & & \\
 & \mathcal{O}_M^h & & &
 \end{array}$$

Vertex algebra V is a Virasoro module. The corresponding semi-infinite complex V^{semi} (the analogue of Chevalley complex for Virasoro algebra) is a vertex algebra too:

$$V^{semi} = V \otimes \Lambda,$$

$$\Lambda \text{ generated by } [b(z), c(w)]_+ = \delta(z - w).$$

The corresponding differential

$$Q = j_0, \quad j(z) = \sum_{n \in \mathbb{Z}} j_n z^{-n-1} = c(z)T(z) + :c(z)\partial c(z)b(z):$$

is nilpotent when $D = 26$ (famous dimension 26!). However, we will consider subcomplex of light modes (i.e. $L_0 = 0$) denoted in the following as (\mathcal{F}_h, Q) , where we can drop this condition:

$$\begin{array}{ccccc}
 & \mathcal{V}^h & & \mathcal{V}^h & \\
 & \nearrow \partial & & \nearrow \partial & \\
 \mathcal{O}_M^h & & \oplus & & \mathcal{O}_M^h \\
 & \searrow \partial & & \searrow \partial & \\
 & \mathcal{O}_M^h & \xrightarrow{id} & \mathcal{O}_M^h & \\
 & & & & \mathcal{O}_M^h
 \end{array}$$

$\frac{1}{2} h \operatorname{div}$ (on the right arrow from \mathcal{V}^h to \mathcal{O}_M^h)
 $-\frac{1}{2} h \operatorname{div}$ (on the left arrow from \mathcal{V}^h to \mathcal{O}_M^h)

Vertex algebra V is a Virasoro module. The corresponding semi-infinite complex V^{semi} (the analogue of Chevalley complex for Virasoro algebra) is a vertex algebra too:

$$V^{semi} = V \otimes \Lambda,$$

$$\Lambda \text{ generated by } [b(z), c(w)]_+ = \delta(z - w).$$

The corresponding differential

$$Q = j_0, \quad j(z) = \sum_{n \in \mathbb{Z}} j_n z^{-n-1} = c(z)T(z) + :c(z)\partial c(z)b(z):$$

is nilpotent when $D = 26$ (famous dimension 26!). However, we will consider subcomplex of light modes (i.e. $L_0 = 0$) denoted in the following as (\mathcal{F}_h, Q) , where we can drop this condition:

$$\begin{array}{ccccc}
 & \mathcal{V}^h & & \mathcal{V}^h & \\
 \nearrow \partial & & \oplus & & \searrow \partial \\
 \mathcal{O}_M^h & & & & \mathcal{O}_M^h \\
 & \mathcal{O}_M^h & \xrightarrow{id} & \mathcal{O}_M^h & \\
 & \nwarrow -\frac{1}{2}hdiv & & \nearrow \frac{1}{2}hdiv &
 \end{array}$$

Vertex algebra V is a Virasoro module. The corresponding semi-infinite complex V^{semi} (the analogue of Chevalley complex for Virasoro algebra) is a vertex algebra too:

$$V^{semi} = V \otimes \Lambda,$$

$$\Lambda \text{ generated by } [b(z), c(w)]_+ = \delta(z - w).$$

The corresponding differential

$$Q = j_0, \quad j(z) = \sum_{n \in \mathbb{Z}} j_n z^{-n-1} = c(z)T(z) + :c(z)\partial c(z)b(z):$$

is nilpotent when $D = 26$ (famous dimension 26!). However, we will consider subcomplex of light modes (i.e. $L_0 = 0$) denoted in the following as (\mathcal{F}_h, Q) , where we can drop this condition:

The diagram illustrates the relationship between vertex algebras and Courant algebroids. It features three vertices:

- Top-left: \mathcal{V}^h
- Top-right: \mathcal{V}^h
- Bottom: \mathcal{O}_M^h

Arrows and labels:

- From \mathcal{O}_M^h to the top-left \mathcal{V}^h : arrow labeled ∂ .
- From the top-left \mathcal{V}^h to the bottom \mathcal{O}_M^h : arrow labeled \oplus .
- From the bottom \mathcal{O}_M^h to the top-right \mathcal{V}^h : arrow labeled ∂ .
- From the top-right \mathcal{V}^h to the bottom \mathcal{O}_M^h : arrow labeled \oplus .
- From the bottom \mathcal{O}_M^h to the top-right \mathcal{V}^h : arrow labeled $\frac{1}{2} h \text{div}$.
- From the top-right \mathcal{V}^h to the bottom \mathcal{O}_M^h : arrow labeled $\frac{1}{2} h \text{div}$.
- From the bottom \mathcal{O}_M^h to the bottom \mathcal{O}_M^h : arrow labeled id .

Vertex algebra V is a Virasoro module. The corresponding semi-infinite complex V^{semi} (the analogue of Chevalley complex for Virasoro algebra) is a vertex algebra too:

$$V^{semi} = V \otimes \Lambda,$$

$$\Lambda \text{ generated by } [b(z), c(w)]_+ = \delta(z - w).$$

The corresponding differential

$$Q = j_0, \quad j(z) = \sum_{n \in \mathbb{Z}} j_n z^{-n-1} = c(z)T(z) + :c(z)\partial c(z)b(z):$$

is nilpotent when $D = 26$ (famous dimension 26!). However, we will consider subcomplex of light modes (i.e. $L_0 = 0$) denoted in the following as (\mathcal{F}_h, Q) , where we can drop this condition:

$$\begin{array}{ccccc}
 & \mathcal{V}^h & & \mathcal{V}^h & \\
 & \nearrow \partial & & \nearrow \partial & \\
 \mathcal{O}_M^h & & \oplus & & \mathcal{O}_M^h \\
 & \searrow \partial & & \searrow \partial & \\
 & & \oplus & & \\
 & & & & \mathcal{O}_M^h
 \end{array}$$

$\mathcal{O}_M^h \xrightarrow{id} \mathcal{O}_M^h$

$\mathcal{O}_M^h \xrightarrow{-\frac{1}{2}h\text{div}} \mathcal{O}_M^h$

$\mathcal{O}_M^h \xrightarrow{\frac{1}{2}h\text{div}} \mathcal{O}_M^h$

Hidden Homotopy Symmetries of Einstein Field Equations

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

Outline

Sigma-models and conformal invariance conditions

Beltrami-Courant differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Anton Zeitlin

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

$$(A, B)_h = \text{Res}_z \frac{A(z)B}{z}$$

Outline

Sigma-models and conformal invariance conditions

Beltrami-Courant differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Anton Zeitlin

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

$$(A, B)_h = \text{Res}_z \frac{A(z)B}{z}$$

$$Q(a_1, a_2)_h = (Qa_1, a_2)_h + (-1)^{|a_1|}(a_1, Qa_2)_h,$$

$$(a_1, a_2)_h - (-1)^{|a_1||a_2|}(a_2, a_1)_h =$$

$$Qm(a_1, a_2) + m(Qa_1, a_2) + (-1)^{|a_1|} m(a_1, Qa_2),$$

$$Q(a_1, a_2, a_3)_h + (Qa_1, a_2, a_3)_h + (-1)^{|a_1|}(a_1, Qa_2, a_3)_h +$$

$$(-1)^{|a_1|+|a_2|}(a_1, a_2, Qa_3)_h = ((a_1, a_2)_h, a_3)_h - (a_1, (a_2, a_3)_h)_h$$

The homotopy Gerstenhaber algebra of Lian and Zuckerman

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

$$(A, B)_h = \text{Res}_z \frac{A(z)B}{z}$$

$$Q(a_1, a_2)_h = (Qa_1, a_2)_h + (-1)^{|a_1|}(a_1, Qa_2)_h,$$

$$(a_1, a_2)_h - (-1)^{|a_1||a_2|}(a_2, a_1)_h =$$

$$Qm(a_1, a_2) + m(Qa_1, a_2) + (-1)^{|a_1|}m(a_1, Qa_2),$$

$$Q(a_1, a_2, a_3)_h + (Qa_1, a_2, a_3)_h + (-1)^{|a_1|}(a_1, Qa_2, a_3)_h +$$

$$(-1)^{|a_1|+|a_2|}(a_1, a_2, Qa_3)_h = ((a_1, a_2)_h, a_3)_h - (a_1, (a_2, a_3)_h)_h$$

Operator **b** of degree -1 (0-mode of $b(z)$) on (\mathcal{F}_h, Q) which anticommutes with Q :

$$\begin{array}{ccc} \mathcal{V}^h & \xleftarrow{-id} & \mathcal{V}^h \\ \oplus & & \oplus \\ \mathcal{O}_M^h & \xleftarrow{id} & \mathcal{O}_M^h \end{array} \quad \begin{array}{ccc} \mathcal{V}^h & \xleftarrow{-id} & \mathcal{V}^h \\ \oplus & & \oplus \\ \mathcal{O}_M^h & \xleftarrow{id} & \mathcal{O}_M^h \end{array}$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The homotopy Gerstenhaber algebra of Lian and Zuckerman

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

$$(A, B)_h = \text{Res}_z \frac{A(z)B}{z}$$

$$Q(a_1, a_2)_h = (Qa_1, a_2)_h + (-1)^{|a_1|}(a_1, Qa_2)_h,$$

$$(a_1, a_2)_h - (-1)^{|a_1||a_2|}(a_2, a_1)_h =$$

$$Qm(a_1, a_2) + m(Qa_1, a_2) + (-1)^{|a_1|}m(a_1, Qa_2),$$

$$Q(a_1, a_2, a_3)_h + (Qa_1, a_2, a_3)_h + (-1)^{|a_1|}(a_1, Qa_2, a_3)_h +$$

$$(-1)^{|a_1|+|a_2|}(a_1, a_2, Qa_3)_h = ((a_1, a_2)_h, a_3)_h - (a_1, (a_2, a_3)_h)_h$$

Operator **b** of degree -1 (0-mode of $b(z)$) on (\mathcal{F}_h, Q) which anticommutes with Q :

$$\begin{array}{ccc} \mathcal{V}^h & \xleftarrow{-id} & \mathcal{V}^h \\ \oplus & & \oplus \\ \mathcal{O}_M^h & \xleftarrow{id} & \mathcal{O}_M^h \end{array} \quad \begin{array}{ccc} \mathcal{V}^h & \xleftarrow{-id} & \mathcal{V}^h \\ \oplus & & \oplus \\ \mathcal{O}_M^h & \xleftarrow{id} & \mathcal{O}_M^h \end{array}$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

One can define a bracket:

$$(-1)^{|a_1|} \{a_1, a_2\}_h = \mathbf{b}(a_1, a_2)_h - (\mathbf{b}a_1, a_2)_h - (-1)^{|a_1|} (a_1 \mathbf{b}a_2)_h,$$

so that together with Q , $(\cdot, \cdot)_h$ it satisfies the relations of homotopy Gerstenhaber algebra:

$$\begin{aligned} & \{a_1, a_2\}_h + (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, a_1\}_h = \\ & (-1)^{|a_1|-1} (Qm'_h(a_1, a_2) - m'_h(Qa_1, a_2) - (-1)^{|a_2|} m'_h(a_1, Qa_2)), \\ & \{a_1, (a_2, a_3)_h\}_h = (\{a_1, a_2\}_h, a_3)_h + (-1)^{(|a_1|-1)|a_2|} (a_2, \{a_1, a_3\}_h)_h, \\ & \{(a_1, a_2)_h, a_3\}_h - (a_1, \{a_2, a_3\}_h)_h - (-1)^{(|a_3|-1)|a_2|} (\{a_1, a_3\}_h, a_2)_h = \\ & (-1)^{|a_1|+|a_2|-1} (Qn'_h(a_1, a_2, a_3) - n'_h(Qa_1, a_2, a_3) - \\ & (-1)^{|a_1|} n'_h(a_1, Qa_2, a_3) - (-1)^{|a_1|+|a_2|} n'_h(a_1, a_2, Qa_3), \\ & \{\{a_1, a_2\}_h, a_3\}_h - \{a_1, \{a_2, a_3\}_h\}_h + \\ & (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, \{a_1, a_3\}_h\}_h = 0. \end{aligned}$$

The conjecture of Lian and Zuckerman, which was later proven by series of papers (Kimura, Zuckerman, Voronov; Huang, Zhao; Voronov) says that the symmetrized product and bracket of homotopy Gerstenhaber algebra constructed above can be lifted to G_∞ -algebra.

One can define a bracket:

$$(-1)^{|a_1|} \{a_1, a_2\}_h = \mathbf{b}(a_1, a_2)_h - (\mathbf{b}a_1, a_2)_h - (-1)^{|a_1|} (a_1 \mathbf{b}a_2)_h,$$

so that together with Q , $(\cdot, \cdot)_h$ it satisfies the relations of homotopy Gerstenhaber algebra:

$$\begin{aligned} & \{a_1, a_2\}_h + (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, a_1\}_h = \\ & (-1)^{|a_1|-1} (Qm'_h(a_1, a_2) - m'_h(Qa_1, a_2) - (-1)^{|a_2|} m'_h(a_1, Qa_2)), \\ & \{a_1, (a_2, a_3)_h\}_h = (\{a_1, a_2\}_h, a_3)_h + (-1)^{(|a_1|-1)|a_2|} (a_2, \{a_1, a_3\}_h)_h, \\ & \{(a_1, a_2)_h, a_3\}_h - (a_1, \{a_2, a_3\}_h)_h - (-1)^{(|a_3|-1)|a_2|} (\{a_1, a_3\}_h, a_2)_h = \\ & (-1)^{|a_1|+|a_2|-1} (Qn'_h(a_1, a_2, a_3) - n'_h(Qa_1, a_2, a_3) - \\ & (-1)^{|a_1|} n'_h(a_1, Qa_2, a_3) - (-1)^{|a_1|+|a_2|} n'_h(a_1, a_2, Qa_3), \\ & \{\{a_1, a_2\}_h, a_3\}_h - \{a_1, \{a_2, a_3\}_h\}_h + \\ & (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, \{a_1, a_3\}_h\}_h = 0. \end{aligned}$$

The conjecture of Lian and Zuckerman, which was later proven by series of papers (Kimura, Zuckerman, Voronov; Huang, Zhao; Voronov) says that the symmetrized product and bracket of homotopy Gerstenhaber algebra constructed above can be lifted to G_∞ -algebra.

One can define a bracket:

$$(-1)^{|a_1|} \{a_1, a_2\}_h = \mathbf{b}(a_1, a_2)_h - (\mathbf{b}a_1, a_2)_h - (-1)^{|a_1|} (a_1 \mathbf{b}a_2)_h,$$

so that together with Q , $(\cdot, \cdot)_h$ it satisfies the relations of homotopy Gerstenhaber algebra:

$$\begin{aligned} & \{a_1, a_2\}_h + (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, a_1\}_h = \\ & (-1)^{|a_1|-1} (Qm'_h(a_1, a_2) - m'_h(Qa_1, a_2) - (-1)^{|a_2|} m'_h(a_1, Qa_2)), \\ & \{a_1, (a_2, a_3)_h\}_h = (\{a_1, a_2\}_h, a_3)_h + (-1)^{(|a_1|-1)|a_2|} (a_2, \{a_1, a_3\}_h)_h, \\ & \{ (a_1, a_2)_h, a_3 \}_h - (a_1, \{a_2, a_3\}_h)_h - (-1)^{(|a_3|-1)|a_2|} (\{a_1, a_3\}_h, a_2)_h = \\ & (-1)^{|a_1|+|a_2|-1} (Qn'_h(a_1, a_2, a_3) - n'_h(Qa_1, a_2, a_3) - \\ & (-1)^{|a_1|} n'_h(a_1, Qa_2, a_3) - (-1)^{|a_1|+|a_2|} n'_h(a_1, a_2, Qa_3), \\ & \{ \{a_1, a_2\}_h, a_3 \}_h - \{a_1, \{a_2, a_3\}_h\}_h + \\ & (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, \{a_1, a_3\}_h\}_h = 0. \end{aligned}$$

The conjecture of Lian and Zuckerman, which was later proven by series of papers (Kimura, Zuckerman, Voronov; Huang, Zhao; Voronov) says that the symmetrized product and bracket of homotopy Gerstenhaber algebra constructed above can be lifted to G_∞ -algebra.

Homotopy algebras: G_∞ , L_∞ , C_∞

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Homotopy algebras: G_∞ , L_∞ , C_∞

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Homotopy algebras: G_∞ , L_∞ , C_∞

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Homotopy algebras: G_∞ , L_∞ , C_∞

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Let A be a graded vector space, consider free graded Lie algebra $Lie(A)$.

$$Lie^{k+1}(A) = [A, Lie^k A], \quad Lie^1(A) = A.$$

Consider free graded commutative algebra GA on the suspension $(Lie(A))[-1]$, i.e.

$$GA = \bigoplus_n \bigwedge^n Lie(A)[-n]$$

There are natural $[\cdot, \cdot]$, \wedge operations on GA of degree -1, 0 correspondingly, generating a Gerstenhaber algebra.

A G_∞ -algebra (Tamarkin, Tsygan, 2000) is a graded space V with a differential ∂ of degree 1 of $G(V[1]^*)$, such that ∂ is a derivation w.r.t bracket and the product.

Multiplicative Ideals, preserved by ∂ : I_1 -generated by the commutant of $Lie(V[1]^*)$, $I_2 = \bigwedge_{n \geq 2} (Lie(V[1]^*)[-n])$. That induces differentials on corresponding factors: $\bigwedge_{n \geq 1} (V[1]^*)[-n]$ and $Lie(V[1]^*)[-1]$. The resulting structures on V are called L_∞ -algebra and C_∞ -algebra correspondingly.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow \text{Lie}^{k_1}(V[1]^*) \wedge \cdots \wedge \text{Lie}^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product, $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow \text{Lie}^{k_1}(V[1]^*) \wedge \cdots \wedge \text{Lie}^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product, $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow Lie^{k_1}(V[1]^*) \wedge \cdots \wedge Lie^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product, $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow Lie^{k_1}(V[1]^*) \wedge \cdots \wedge Lie^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product, $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow Lie^{k_1}(V[1]^*) \wedge \cdots \wedge Lie^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product, $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow Lie^{k_1}(V[1]^*) \wedge \cdots \wedge Lie^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product,
 $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated
 by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is
 of degree 2) :

$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Restriction of ∂ on $V[1]^*$:

$$V[1]^* \rightarrow Lie^{k_1}(V[1]^*) \wedge \cdots \wedge Lie^{k_n}(V[1]^*).$$

Conjugated map:

$$m_{k_1, k_2, \dots, k_n} : V^{\otimes k_1} \otimes \cdots \otimes V^{\otimes k_n} \rightarrow V.$$

of degree $3 - n - k_1 - \dots - k_n$, satisfying bilinear relations.

In our previous notation $m_1 = Q$, m_2 -symmetrized LZ product,
 $m_{1,1}$ -antisymmetrized LZ bracket.

L_∞ is generated by $m_1 \equiv Q$, $m_{1,1,\dots,1} \equiv [\cdot, \dots, \cdot]$ and C_∞ is generated
 by $m_1 \equiv Q$, $m_k \equiv (\cdot, \dots, \cdot)$.

An important feature of L_∞ algebra is a Maurer-Cartan equation (Φ is
 of degree 2) :


$$Q\Phi + \sum_{n \geq 2} \frac{1}{n!} \underbrace{[\Phi, \dots, \Phi]}_n + \cdots = 0,$$

which has infinitesimal symmetries:

$$\Phi \rightarrow \Phi + Q\Lambda + \sum_{n \geq 1} \frac{1}{n!} \underbrace{[\Phi \dots \Phi, \Lambda]}_n$$

Quasiclassical limit of LZ G_∞ algebra

The following complex (\mathcal{F}, Q) :



is a subcomplex of (\mathcal{F}_h, Q) . Then

$$(\cdot, \cdot)_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow \mathcal{F}^{i+j}[h], \quad \{\cdot, \cdot\}_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow h\mathcal{F}_{i+j-1}[h],$$

$$\mathbf{b} : \mathcal{F}^i \rightarrow h\mathcal{F}^{i-1}[h],$$

so that

$$(\cdot, \cdot)_0 = \lim_{h \rightarrow 0} (\cdot, \cdot)_h, \quad \{\cdot, \cdot\}_0 = \lim_{h \rightarrow 0} h^{-1} \{\cdot, \cdot\}_h, \quad \mathbf{b}_0 = \lim_{h \rightarrow 0} h^{-1} \mathbf{b}$$

are well defined.

Outline

Sigma-models and
conformal invariance
conditions


Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Quasiclassical limit of LZ G_∞ algebra

The following complex (\mathcal{F}, Q) :



is a subcomplex of (\mathcal{F}_h, Q) . Then

$$(\cdot, \cdot)_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow \mathcal{F}^{i+j}[h], \quad \{\cdot, \cdot\}_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow h\mathcal{F}_{i+j-1}[h],$$

$$\mathbf{b} : \mathcal{F}^i \rightarrow h\mathcal{F}^{i-1}[h],$$

so that

$$(\cdot, \cdot)_0 = \lim_{h \rightarrow 0} (\cdot, \cdot)_h, \quad \{\cdot, \cdot\}_0 = \lim_{h \rightarrow 0} h^{-1} \{\cdot, \cdot\}_h, \quad \mathbf{b}_0 = \lim_{h \rightarrow 0} h^{-1} \mathbf{b}$$

are well defined.

Outline

Sigma-models and
conformal invariance
conditions


Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Quasiclassical limit of LZ G_∞ algebra

The following complex (\mathcal{F}, Q) :



is a subcomplex of (\mathcal{F}_h, Q) . Then

$$(\cdot, \cdot)_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow \mathcal{F}^{i+j}[h], \quad \{\cdot, \cdot\}_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow h\mathcal{F}_{i+j-1}[h],$$

$$\mathbf{b} : \mathcal{F}^i \rightarrow h\mathcal{F}^{i-1}[h],$$

so that

$$(\cdot, \cdot)_0 = \lim_{h \rightarrow 0} (\cdot, \cdot)_h, \quad \{\cdot, \cdot\}_0 = \lim_{h \rightarrow 0} h^{-1} \{\cdot, \cdot\}_h, \quad \mathbf{b}_0 = \lim_{h \rightarrow 0} h^{-1} \mathbf{b}$$

are well defined.

Outline

Sigma-models and
conformal invariance
conditions


Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Quasiclassical limit of LZ G_∞ algebra

The following complex (\mathcal{F}, Q) :



is a subcomplex of (\mathcal{F}_h, Q) . Then

$$(\cdot, \cdot)_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow \mathcal{F}^{i+j}[h], \quad \{\cdot, \cdot\}_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow h\mathcal{F}_{i+j-1}[h],$$

$$\mathbf{b} : \mathcal{F}^i \rightarrow h\mathcal{F}^{i-1}[h],$$

so that

$$(\cdot, \cdot)_0 = \lim_{h \rightarrow 0} (\cdot, \cdot)_h, \quad \{\cdot, \cdot\}_0 = \lim_{h \rightarrow 0} h^{-1} \{\cdot, \cdot\}_h, \quad \mathbf{b}_0 = \lim_{h \rightarrow 0} h^{-1} \mathbf{b}$$

are well defined.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The symmetrized operations $(\cdot, \cdot)_0, \{\cdot, \cdot\}_0, \dots$ satisfy the relations of the homotopy Gerstenhaber algebra, so that all non-covariant higher-order terms disappear from the multilinear operations.

The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Conjecture: This G_∞ -algebra is the G_3 -algebra (no homotopies beyond trilinear operations).

Classical limit procedure for vertex algebroid (due to P. Bressler):

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \lim_{h \rightarrow 0} \frac{1}{h} \langle \cdot, \cdot \rangle.$$

The resulting operations form a Courant algebroid (Z.-J. Liu, A. Weinstein, P. Xu, 1997)

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

**Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit**

Einstein Equations

The symmetrized operations $(\cdot, \cdot)_0$, $\{\cdot, \cdot\}_0, \dots$ satisfy the relations of the homotopy Gerstenhaber algebra, so that all non-covariant higher-order terms disappear from the multilinear operations.

The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Conjecture: This G_∞ -algebra is the G_3 -algebra (no homotopies beyond trilinear operations).

Classical limit procedure for vertex algebroid (due to P. Bressler):

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \lim_{h \rightarrow 0} \frac{1}{h} \langle \cdot, \cdot \rangle.$$

The resulting operations form a Courant algebroid (Z.-J. Liu, A. Weinstein, P. Xu, 1997)

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The symmetrized operations $(\cdot, \cdot)_0, \{\cdot, \cdot\}_0, \dots$ satisfy the relations of the homotopy Gerstenhaber algebra, so that all non-covariant higher-order terms disappear from the multilinear operations.

The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Conjecture: This G_∞ -algebra is the G_3 -algebra (no homotopies beyond trilinear operations).

Classical limit procedure for vertex algebroid (due to P. Bressler):

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \lim_{h \rightarrow 0} \frac{1}{h} \langle \cdot, \cdot \rangle.$$

The resulting operations form a Courant algebroid (Z.-J. Liu, A. Weinstein, P. Xu, 1997)

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The symmetrized operations $(\cdot, \cdot)_0, \{\cdot, \cdot\}_0, \dots$ satisfy the relations of the homotopy Gerstenhaber algebra, so that all non-covariant higher-order terms disappear from the multilinear operations.

The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Conjecture: This G_∞ -algebra is the G_3 -algebra (no homotopies beyond trilinear operations).

Classical limit procedure for vertex algebroid (due to P. Bressler):

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \lim_{h \rightarrow 0} \frac{1}{h} \langle \cdot, \cdot \rangle.$$

The resulting operations form a Courant algebroid (Z.-J. Liu, A. Weinstein, P. Xu, 1997)

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

The symmetrized operations $(\cdot, \cdot)_0, \{\cdot, \cdot\}_0, \dots$ satisfy the relations of the homotopy Gerstenhaber algebra, so that all non-covariant higher-order terms disappear from the multilinear operations.

The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Conjecture: This G_∞ -algebra is the G_3 -algebra (no homotopies beyond trilinear operations).

Classical limit procedure for vertex algebroid (due to P. Bressler):

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \lim_{h \rightarrow 0} \frac{1}{h} \langle \cdot, \cdot \rangle.$$

The resulting operations form a Courant algebroid (Z.-J. Liu, A. Weinstein, P. Xu, 1997)

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

A Courant \mathcal{O}_M -algebroid is an \mathcal{O}_M -module \mathcal{Q} equipped with a structure of a Leibniz \mathbb{C} -algebra $[\cdot, \cdot]_0 : \mathcal{Q} \otimes_{\mathbb{C}} \mathcal{Q} \rightarrow \mathcal{Q}$, an \mathcal{O}_M -linear map of Leibniz algebras (the anchor map) $\pi_0 : \mathcal{Q} \rightarrow \Gamma(TM)$, a symmetric \mathcal{O}_M -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{Q} \otimes_{\mathcal{O}_M} \mathcal{Q} \rightarrow \mathcal{O}_M$, a derivation $\partial : \mathcal{O}_M \rightarrow \mathcal{Q}$ which satisfy

$$\begin{aligned}\pi_0 \circ \partial &= 0, & [q_1, fq_2]_0 &= f[q_1, q_2]_0 + \pi_0(q_1)(f)q_2 \\ \langle [q, q_1], q_2 \rangle + \langle q_1, [q, q_2] \rangle &= \pi_0(q)(\langle q_1, q_2 \rangle_0), \\ [q, \partial(f)]_0 &= \partial(\pi_0(q)(f)) \\ \langle q, \partial(f) \rangle &= \pi_0(q)(f) & [q_1, q_2]_0 + [q_2, q_1]_0 &= \partial \langle q_1, q_2 \rangle_0\end{aligned}$$

for $f \in \mathcal{O}_M$ and $q, q_1, q_2 \in \mathcal{Q}$.

First it was obtained as an analogue of Manin's double for Lie bialgebroid by Z-J. Liu, A. Weinstein, P. Xu.

In our case $\mathcal{Q} \cong \mathcal{O}(\mathcal{E})$, π_0 is just a projection on $\mathcal{O}(TM)$

$$[q_1, q_2]_0 = -[q_1, q_2]_D, \quad \langle q_1, q_2 \rangle_0 = -\langle q_1, q_2 \rangle^s, \quad \partial = d.$$

Outline

Sigma-models and
conformal invariance
conditionsBeltrami-Courant
differentialVertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

A Courant \mathcal{O}_M -algebroid is an \mathcal{O}_M -module \mathcal{Q} equipped with a structure of a Leibniz \mathbb{C} -algebra $[\cdot, \cdot]_0 : \mathcal{Q} \otimes_{\mathbb{C}} \mathcal{Q} \rightarrow \mathcal{Q}$, an \mathcal{O}_M -linear map of Leibniz algebras (the anchor map) $\pi_0 : \mathcal{Q} \rightarrow \Gamma(TM)$, a symmetric \mathcal{O}_M -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{Q} \otimes_{\mathcal{O}_M} \mathcal{Q} \rightarrow \mathcal{O}_M$, a derivation $\partial : \mathcal{O}_M \rightarrow \mathcal{Q}$ which satisfy

$$\pi \circ \partial = 0, \quad [q_1, fq_2]_0 = f[q_1, q_2]_0 + \pi_0(q_1)(f)q_2$$

$$\langle [q, q_1], q_2 \rangle + \langle q_1, [q, q_2] \rangle = \pi_0(q)(\langle q_1, q_2 \rangle_0),$$

$$[q, \partial(f)]_0 = \partial(\pi_0(q)(f))$$

$$\langle q, \partial(f) \rangle = \pi_0(q)(f) \quad [q_1, q_2]_0 + [q_2, q_1]_0 = \partial \langle q_1, q_2 \rangle_0$$

for $f \in \mathcal{O}_M$ and $q, q_1, q_2 \in \mathcal{Q}$.

First it was obtained as an analogue of Manin's double for Lie bialgebroid by Z-J. Liu, A. Weinstein, P. Xu.

In our case $\mathcal{Q} \cong \mathcal{O}(\mathcal{E})$, π_0 is just a projection on $\mathcal{O}(TM)$

$$[q_1, q_2]_0 = -[q_1, q_2]_D, \quad \langle q_1, q_2 \rangle_0 = -\langle q_1, q_2 \rangle^s, \quad \partial = d.$$

A Courant \mathcal{O}_M -algebroid is an \mathcal{O}_M -module \mathcal{Q} equipped with a structure of a Leibniz \mathbb{C} -algebra $[\cdot, \cdot]_0 : \mathcal{Q} \otimes_{\mathbb{C}} \mathcal{Q} \rightarrow \mathcal{Q}$, an \mathcal{O}_M -linear map of Leibniz algebras (the anchor map) $\pi_0 : \mathcal{Q} \rightarrow \Gamma(TM)$, a symmetric \mathcal{O}_M -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{Q} \otimes_{\mathcal{O}_M} \mathcal{Q} \rightarrow \mathcal{O}_M$, a derivation $\partial : \mathcal{O}_M \rightarrow \mathcal{Q}$ which satisfy

$$\begin{aligned}\pi_0 \circ \partial &= 0, & [q_1, fq_2]_0 &= f[q_1, q_2]_0 + \pi_0(q_1)(f)q_2 \\ \langle [q, q_1], q_2 \rangle + \langle q_1, [q, q_2] \rangle &= \pi_0(q)(\langle q_1, q_2 \rangle_0), \\ [q, \partial(f)]_0 &= \partial(\pi_0(q)(f)) \\ \langle q, \partial(f) \rangle &= \pi_0(q)(f) & [q_1, q_2]_0 + [q_2, q_1]_0 &= \partial \langle q_1, q_2 \rangle_0\end{aligned}$$

for $f \in \mathcal{O}_M$ and $q, q_1, q_2 \in \mathcal{Q}$.

First it was obtained as an analogue of Manin's double for Lie bialgebroid by Z-J. Liu, A. Weinstein, P. Xu.

In our case $\mathcal{Q} \cong \mathcal{O}(\mathcal{E})$, π_0 is just a projection on $\mathcal{O}(TM)$

$$[q_1, q_2]_0 = -[q_1, q_2]_D, \quad \langle q_1, q_2 \rangle_0 = -\langle q_1, q_2 \rangle^s, \quad \partial = d.$$

The corresponding L_3 -algebra on the half-complex for Courant algebroid was constructed by D. Roytenberg and A. Weinstein (1998).

We show that it is a part of a more general structure, homotopy Gerstenhaber algebra.

Question: Is there a direct path (avoiding vertex algebra) from Courant algebroid to G_3 -algebra? Odd analogue of Manin double?

Remark. C_3 -algebra is related to gauge theory. The appropriate "metric" deformation gives a Yang-Mills C_3 -algebra on a flat space.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

The corresponding L_3 -algebra on the half-complex for Courant algebroid was constructed by D. Roytenberg and A. Weinstein (1998).

We show that it is a part of a more general structure, homotopy Gerstenhaber algebra.

Question: Is there a direct path (avoiding vertex algebra) from Courant algebroid to G_3 -algebra? Odd analogue of Manin double?

Remark. C_3 -algebra is related to gauge theory. The appropriate "metric" deformation gives a Yang-Mills C_3 -algebra on a flat space.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

The corresponding L_3 -algebra on the half-complex for Courant algebroid was constructed by D. Roytenberg and A. Weinstein (1998).

We show that it is a part of a more general structure, homotopy Gerstenhaber algebra.


Question: Is there a direct path (avoiding vertex algebra) from Courant algebroid to G_3 -algebra? Odd analogue of Manin double?

Remark. C_3 -algebra is related to gauge theory. The appropriate "metric" deformation gives a Yang-Mills C_3 -algebra on a flat space.

A.Z., Comm. Math. Phys. 303 (2011) 331-359.

Simplest version: $G_\infty \rightarrow$ Gerstenhaber algebra

Subcomplex (\mathcal{F}_{sm}, Q) :



The G_∞ algebra degenerates to G-algebra. Moreover, due to \mathbf{b}_0 it is a BV-algebra. Combine chiral and antichiral part:

$$\mathbf{F}_{sm} = \mathcal{F}_{sm} \otimes \bar{\mathcal{F}}_{sm}$$

$$(-1)^{|a_1|} \{a_1, a_2\} = \mathbf{b}^-(a_1, a_2) - (\mathbf{b}^- a_1, a_2) - (-1)^{|a_1|} (a_1 \mathbf{b}^- a_2),$$

where $\mathbf{b}^- = \mathbf{b} - \bar{\mathbf{b}}$.

Outline

Sigma-models and
conformal invariance
conditions


Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Simplest version: $G_\infty \rightarrow$ Gerstenhaber algebra

Subcomplex (\mathcal{F}_{sm}, Q) :



The G_∞ algebra degenerates to G-algebra. Moreover, due to \mathbf{b}_0 it is a BV-algebra. Combine chiral and antichiral part:

$$\mathbf{F}_{sm} = \mathcal{F}_{sm} \otimes \bar{\mathcal{F}}_{sm}$$

$$(-1)^{|a_1|} \{a_1, a_2\} = \mathbf{b}^-(a_1, a_2) - (\mathbf{b}^- a_1, a_2) - (-1)^{|a_1|} (a_1 \mathbf{b}^- a_2),$$

where $\mathbf{b}^- = \mathbf{b} - \bar{\mathbf{b}}$.

Outline

Sigma-models and
conformal invariance
conditions


Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Simplest version: $G_\infty \rightarrow$ Gerstenhaber algebra

Subcomplex (\mathcal{F}_{sm}, Q) :



The G_∞ algebra degenerates to G-algebra. Moreover, due to \mathbf{b}_0 it is a BV-algebra. Combine chiral and antichiral part:

$$\mathbf{F}_{sm} = \mathcal{F}_{sm} \otimes \bar{\mathcal{F}}_{sm}$$

$$(-1)^{|a_1|} \{a_1, a_2\} = \mathbf{b}^-(a_1, a_2) - (\mathbf{b}^- a_1, a_2) - (-1)^{|a_1|} (a_1 \mathbf{b}^- a_2),$$

where $\mathbf{b}^- = \mathbf{b} - \bar{\mathbf{b}}$.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M) \oplus \mathcal{O}(T^{(1,0)}(M) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

A.Z., Nucl. Phys. B 794 (2008) 370-398; A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

- 1). Vector field $\text{div}_\Omega g$, where $\log \Omega = -2\Phi_0 = -2(\phi' + \bar{\phi}' + \phi + \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.
- 2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_\Omega(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_\Omega(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

- 3). $\text{div}_\Omega \text{div}_\Omega(g) = 0$.

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(1,0)}(M)) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

A.Z., Nucl. Phys. B 794 (2008) 370-398; A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

- 1). Vector field $\text{div}_{\Omega} g$, where $\log \Omega = -2\Phi_0 = -2(\phi' + \bar{\phi}' + \phi + \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.
- 2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_{\Omega}(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_{\Omega}(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

- 3). $\text{div}_{\Omega} \text{div}_{\Omega}(g) = 0$.

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(1,0)}(M)) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

A.Z., Nucl. Phys. B 794 (2008) 370-398; A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

1). Vector field $\text{div}_\Omega g$, where $\log \Omega = -2\Phi_0 = -2(\phi' + \bar{\phi}' + \phi + \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.

2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_\Omega(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_\Omega(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

3). $\text{div}_\Omega \text{div}_\Omega(g) = 0$.

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(1,0)}(M)) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

A.Z., Nucl. Phys. B 794 (2008) 370-398; A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

- 1). Vector field $\text{div}_{\Omega} g$, where $\log \Omega = -2\Phi_0 = -2(\phi' + \bar{\phi}' + \phi + \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.
- 2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_{\Omega}(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_{\Omega}(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

- 3). $\text{div}_{\Omega} \text{div}_{\Omega}(g) = 0$.

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(1,0)}(M)) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

A.Z., Nucl. Phys. B 794 (2008) 370-398; A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

- 1). Vector field $\text{div}_{\Omega} g$, where $\log \Omega = -2\Phi_0 = -2(\phi' + \bar{\phi}' + \phi + \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.
- 2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_{\Omega}(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_{\Omega}(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

- 3). $\text{div}_{\Omega} \text{div}_{\Omega}(g) = 0$.

These are Einstein equations with the following constraints:

$$\begin{aligned} G_{i\bar{k}} &= g_{i\bar{k}}, & B_{i\bar{k}} &= -g_{i\bar{k}}, & \Phi &= \log \sqrt{g} + \Phi_0, \\ G_{ik} &= G_{\bar{i}\bar{k}} = G_{ik} = G_{\bar{i}\bar{k}} = 0, \end{aligned}$$

Physically:

$$\begin{aligned} &\int [dp][d\bar{p}][dX][d\bar{X}] e^{-\frac{1}{2\pi i\hbar} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \langle g, p \wedge \bar{p} \rangle) + \int_{\Sigma} R^{(2)}(\gamma) \Phi_0(X)} = \\ &\int [dX][d\bar{X}] e^{\frac{-1}{4\pi\hbar} \int d^2z (G_{\mu\nu} + B_{\mu\nu}) \partial X^\mu \bar{\partial} X^\nu + \int R^{(2)}(\gamma) (\Phi_0(X) + \log \sqrt{g})} \end{aligned}$$

based on computations of

A. Tseytlin and A. Schwarz, Nucl.Phys. B399 (1993) 691-708.

Outline

Sigma-models and
 conformal invariance
 conditions

Beltrami-Courant
 differential

Vertex/Courant
 algebroids,
 G_{∞} -algebras and
 quasiclassical limit

Einstein Equations

These are Einstein equations with the following constraints:

$$\begin{aligned} G_{i\bar{k}} &= g_{i\bar{k}}, & B_{i\bar{k}} &= -g_{i\bar{k}}, & \Phi &= \log \sqrt{g} + \Phi_0, \\ G_{ik} &= G_{\bar{i}\bar{k}} = G_{ik} = G_{\bar{i}\bar{k}} = 0, \end{aligned}$$

Physically:

$$\begin{aligned} \int [dp][d\bar{p}][dX][d\bar{X}] e^{-\frac{1}{2\pi i\hbar} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \langle g, p \wedge \bar{p} \rangle) + \int_{\Sigma} R^{(2)}(\gamma) \Phi_0(X)} = \\ \int [dX][d\bar{X}] e^{\frac{-1}{4\pi\hbar} \int d^2z (G_{\mu\nu} + B_{\mu\nu}) \partial X^{\mu} \bar{\partial} X^{\nu} + \int R^{(2)}(\gamma) (\Phi_0(X) + \log \sqrt{g})} \end{aligned}$$

based on computations of

A. Tseytlin and A. Schwarz, Nucl.Phys. B399 (1993) 691-708.

Consider

$$\mathbf{F}_{b-} = \mathcal{F} \otimes \bar{\mathcal{F}}|_{b-=0}$$

with the L_∞ -algebra structure given by Lian-Zuckerman construction.

One can explicitly check that GMC symmetry

($\Psi = \Psi(\mathbb{M}, \Phi, \text{auxiliary fields})$)

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

reproduces

$$\mathbb{M} \rightarrow \mathbb{M} - D\alpha + \phi_1(\alpha, \mathbb{M}) + \phi_2(\alpha, \mathbb{M}, \mathbb{M}).$$

A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

Conjecture: The corresponding Maurer-Cartan equation gives Einstein equations on G, B, Φ expressed in terms of Beltrami-Courant differential. The symmetries of the Maurer-Cartan equation reproduce mentioned above symmetries of Einstein equations.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Consider

$$\mathbf{F}_{b-} = \mathcal{F} \otimes \bar{\mathcal{F}}|_{b-=0}$$

with the L_∞ -algebra structure given by Lian-Zuckerman construction.
One can explicitly check that GMC symmetry
($\Psi = \Psi(\mathbb{M}, \Phi, \text{auxiliary fields})$)

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

reproduces

$$\mathbb{M} \rightarrow \mathbb{M} - D\alpha + \phi_1(\alpha, \mathbb{M}) + \phi_2(\alpha, \mathbb{M}, \mathbb{M}).$$

A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

Conjecture: The corresponding Maurer-Cartan equation gives Einstein equations on G, B, Φ expressed in terms of Beltrami-Courant differential. The symmetries of the Maurer-Cartan equation reproduce mentioned above symmetries of Einstein equations.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Consider

$$\mathbf{F}_{b-} = \mathcal{F} \otimes \bar{\mathcal{F}}|_{b-=0}$$

with the L_∞ -algebra structure given by Lian-Zuckerman construction.
One can explicitly check that GMC symmetry
($\Psi = \Psi(\mathbb{M}, \Phi, \text{auxiliary fields})$)

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

reproduces

$$\mathbb{M} \rightarrow \mathbb{M} - D\alpha + \phi_1(\alpha, \mathbb{M}) + \phi_2(\alpha, \mathbb{M}, \mathbb{M}).$$

A.Z., Adv. Theor. Math. Phys. 19 (2015) 1249-1275

Conjecture: The corresponding Maurer-Cartan equation gives Einstein equations on G, B, Φ expressed in terms of Beltrami-Courant differential. The symmetries of the Maurer-Cartan equation reproduce mentioned above symmetries of Einstein equations.

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_∞ -algebras and
quasiclassical limit

Einstein Equations

Outline

Sigma-models and
conformal invariance
conditions

Beltrami-Courant
differential

Vertex/Courant
algebroids,
 G_{∞} -algebras and
quasiclassical limit

Einstein Equations

Thank you!