

Beltrami-Courant Differentials and Homotopy Gerstenhaber algebras

Anton M. Zeitlin

Columbia University, Department of Mathematics
Institute des Hautes Études Scientifiques

30th International Colloquium on Group Theoretical Methods in Physics

Ghent, Belgium

July 13 - July 19, 2014

Outline

Sigma-models and
conformal invariance
conditions

Reformulation of
Sigma-model in
first-order form

Vertex algebroids,
 G_{∞} -algebra and
quasiclassical limit

Einstein Equations



Outline

Sigma-models and
conformal invariance
conditions

Reformulation of
Sigma-model in
first-order form

Vertex algebroids,
 G_∞ -algebra and
quasiclassical limit

Einstein Equations

Sigma-models and conformal invariance conditions

Reformulation of Sigma-model in first-order form

Vertex algebroids, G_∞ -algebra and quasiclassical limit

Einstein Equations from G_∞ -algebras via Beltrami-Courant differential

Sigma-model:

$$S_{so} = \frac{1}{4\pi h} \int d^2z (G_{\mu\nu} + B_{\mu\nu}) \partial X^\mu \bar{\partial} X^\nu + \int_\Sigma \sqrt{\gamma} R^{(2)}(\gamma) \Phi(X)$$

Symmetries: Diff symmetry and $B \rightarrow B + d\lambda$.

Conformal invariance conditions:

$$\begin{aligned} \mu \frac{d}{d\mu} G_{\mu\nu} &= \beta_{\mu\nu}^G(G, B, \Phi, h) = 0, & \mu \frac{d}{d\mu} B_{\mu\nu} &= \beta_{\mu\nu}^B(G, B, \Phi, h) = 0, \\ \mu \frac{d}{d\mu} \Phi &= \beta^\Phi(G, B, \Phi, h) = 0 \end{aligned}$$

at the level h^0 turn out to be Einstein Equations:

$$\begin{aligned} R^{\mu\nu} &= \frac{1}{4} H^{\mu\lambda\rho} H_{\lambda\rho}^\nu - 2\nabla^\mu \nabla^\nu \Phi, \\ \nabla_\mu H^{\mu\nu\rho} - 2(\nabla_\lambda \Phi) H^{\lambda\nu\rho} &= 0, \\ 4(\nabla_\mu \Phi)^2 - 4\nabla_\mu \nabla^\mu \Phi + R + \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} &= 0. \end{aligned}$$

Outline

Sigma-models and
conformal invariance
conditions

Reformulation of
Sigma-model in
first-order form

Vertex algebroids,
 G_∞ -algebra and
quasiclassical limit

Einstein Equations

Early days of string theory: linearized Einstein Equations and their symmetries: ($G_{\mu\nu} = \eta_{\mu\nu} + s_{\mu\nu}$):

$$Q^\eta \Psi^s = 0, \quad \Psi^s \rightarrow \Psi^s + Q\Lambda$$

in a BRST complex associated to certain Virasoro module.

It was conjectured (Sen, Zwiebach,...) that Einstein equations with h -corrections and their symmetries are Generalized Maurer-Cartan (GMC) Equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi]_h + \frac{1}{3!}[\Psi, \Psi, \Psi]_h + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

where $[\cdot, \cdot, \dots, \cdot]_h$ operations generate L_∞ -algebra.

We show that for a proper background, there is a richer structure, namely G_∞ -algebra, as well as a well-defined classical limit of the L_∞ -subalgebra, so that GMC equations are equivalent to Einstein Equations.

First order version of sigma-model action

We start from free action:

$$S_0 = \frac{1}{2\pi i\hbar} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle),$$

where $p \in X^*(\Omega^{(1,0)}(M)) \otimes \Omega^{(1,0)}(\Sigma)$, $\bar{p} \in X^*(\Omega^{(0,1)}(M)) \otimes \Omega^{(0,1)}(\Sigma)$.

Symmetries: $X^i \rightarrow X^i - v^i(X)$, $X^{\bar{i}} \rightarrow X^{\bar{i}} - v^{\bar{i}}(\bar{X})$,

$$p_i \rightarrow p_i + \partial_i v^k p_k, \quad p_{\bar{i}} \rightarrow p_{\bar{i}} + \partial_{\bar{i}} v^{\bar{k}} p_{\bar{k}}$$

$$p_i \rightarrow p_i - \partial X^k (\partial_k \omega_i - \partial_i \omega_k), \quad p_{\bar{i}} \rightarrow p_{\bar{i}} - \bar{\partial} X^{\bar{k}} (\partial_{\bar{k}} \omega_{\bar{i}} - \partial_{\bar{i}} \omega_{\bar{k}}).$$

Not invariant under general diffeomorphisms, i.e.

$$\delta S_0 = -\frac{1}{2\pi i\hbar} \int (\langle \bar{\partial} v, p \wedge \bar{\partial} X \rangle + \langle \partial \bar{v}, \bar{p} \wedge \partial X \rangle).$$

It is necessary to add extra terms:

$$\delta S_{\mu} = -\frac{1}{2\pi i\hbar} \int (\langle \mu, p \wedge \bar{\partial} X \rangle + \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle),$$

where $\mu \in \Gamma(T^{(1,0)}M \otimes T^{*(0,1)}(M))$, $\bar{\mu} \in \Gamma(T^{(0,1)}M \otimes T^{*(1,0)}(M))$, so

that: $\mu \rightarrow \mu - \bar{\partial} v + \dots$, $\bar{\mu} \rightarrow \bar{\mu} - \partial \bar{v} + \dots$

Continuing the procedure:

$$\tilde{S} = \frac{1}{2\pi i h} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \partial X \wedge \bar{p} \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle),$$

where

$$\begin{aligned} \mu_j^i &\rightarrow \mu_j^i - \partial_{\bar{j}} v^i + v^k \partial_k \mu_j^i + v^{\bar{k}} \partial_{\bar{k}} \mu_j^i + \mu_{\bar{k}}^i \partial_{\bar{j}} v^{\bar{k}} - \mu_j^k \partial_k v^i + \mu_{\bar{j}}^i \mu_j^k \partial_k v^{\bar{j}}, \\ b_{i\bar{j}} &\rightarrow b_{i\bar{j}} + v^k \partial_k b_{i\bar{j}} + v^{\bar{k}} \partial_{\bar{k}} b_{i\bar{j}} + b_{i\bar{k}} \partial_{\bar{j}} v^{\bar{k}} + b_{\bar{j}} \partial_i v^{\bar{j}} + b_{i\bar{k}} \mu_j^k \partial_k v^{\bar{k}} + b_{\bar{j}} \bar{\mu}_i^{\bar{k}} \partial_{\bar{k}} v^{\bar{j}}, \end{aligned}$$

so that:

$$\begin{aligned} X^i &\rightarrow X^i - v^i(X, \bar{X}), & p_i &\rightarrow p_i + p_k \partial_i v^k - p_k \mu_{\bar{j}}^k \partial_i v^{\bar{j}} - b_{j\bar{k}} \partial_i v^{\bar{k}} \partial X^j, \\ X^{\bar{i}} &\rightarrow X^{\bar{i}} - v^{\bar{i}}(X, \bar{X}), & \bar{p}_{\bar{i}} &\rightarrow \bar{p}_{\bar{i}} + \bar{p}_{\bar{k}} \partial_{\bar{i}} v^{\bar{k}} - \bar{p}_{\bar{k}} \bar{\mu}_i^{\bar{k}} \partial_i v^{\bar{j}} - b_{j\bar{k}} \partial_{\bar{i}} v^{\bar{k}} \bar{\partial} X^{\bar{j}}. \end{aligned}$$

For simplicity: $E = TM \oplus T^*M$ and $\mathcal{E} = T^{(1,0)}M \oplus T^{*(1,0)}M$,
 $\bar{\mathcal{E}} = T^{(0,1)}M \oplus T^{*(0,1)}M$, i.e. $E = \mathcal{E} \oplus \bar{\mathcal{E}}$.

Let $\tilde{M} \in \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$, such that

$$\tilde{M} = \begin{pmatrix} 0 & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

Introduce $\alpha \in \Gamma(E)$, i.e. $\alpha = (v, \bar{v}, \omega, \bar{\omega})$. Let $D : \Gamma(E) \rightarrow \Gamma(\mathcal{E} \otimes \bar{\mathcal{E}})$,
 such that

$$D\alpha = \begin{pmatrix} 0 & \bar{\partial}v \\ \partial\bar{v} & \partial\bar{\omega} - \bar{\partial}\omega \end{pmatrix}.$$

Then the transformation of \tilde{M} can be expressed:

$$\tilde{M} \rightarrow \tilde{M} - D\alpha + \phi_1(\alpha, \tilde{M}) + \phi_2(\alpha, \tilde{M}, \tilde{M}).$$

Let us describe ϕ_1, ϕ_2 algebraically. In order to do that we need to pass
 to jet bundles, i.e.

$$\alpha \in J^\infty(\mathcal{O}_M) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}})) \oplus J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}_M),$$

$$\tilde{M} \in J^\infty(\mathcal{O}(\mathcal{E})) \otimes J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$$

Then:

$$\alpha = \sum_J f^J \otimes \bar{b}^J + \sum_K b^K \otimes \bar{f}^K,$$

$$\tilde{\mathbb{M}} = \sum_I a^I \otimes \bar{a}^I,$$

where $a^I, b^J \in J^\infty(\mathcal{O}(\mathcal{E}))$, $f^I \in J^\infty(\mathcal{O}_M)$ and $\bar{a}^I, \bar{b}^J \in J^\infty(\bar{\mathcal{O}}(\bar{\mathcal{E}}))$, $\bar{f}^I \in J^\infty(\bar{\mathcal{O}}_M)$. Then

$$\phi_1(\alpha, \tilde{\mathbb{M}}) = \sum_{I,J} [b^J, a^I]_D \otimes \bar{f}^J \bar{a}^I + \sum_{I,K} f^K a^I \otimes [\bar{b}^K, \bar{a}^I]_D,$$

where $[\cdot, \cdot]_D$ is a Dorfman bracket:

$$[v_1, v_2]_D = [v_1, v_2]^{Lie}, \quad [v, \omega]_D = L_v \omega,$$

$$[\omega, v]_D = -i_v d\omega, \quad [\omega_1, \omega_2]_D = 0.$$

Similarly:

$$\phi_2(\alpha, \tilde{\mathbb{M}}, \tilde{\mathbb{M}}) = \tilde{\mathbb{M}} \cdot D\alpha \cdot \tilde{\mathbb{M}}$$

$$\frac{1}{2} \sum_{I,J,K} \langle b^I, a^K \rangle a^J \otimes \bar{a}^J (\bar{f}^I) \bar{a}^K + \frac{1}{2} \sum_{I,J,K} a^J (f^I) a^K \otimes \langle \bar{b}^I, \bar{a}^K \rangle \bar{a}^J.$$

Relation to standard second order sigma-model: Let us fill in 0 in $\tilde{\mathbb{M}}$:

$$\mathbb{M} = \begin{pmatrix} g & \mu \\ \bar{\mu} & b \end{pmatrix}.$$

$$S_{fo} = \frac{1}{2\pi i h} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle + \langle \bar{p} \wedge \partial X \rangle - \\ - \langle g, p \wedge \bar{p} \rangle - \langle \mu, p \wedge \bar{\partial} X \rangle - \langle \bar{\mu}, \bar{p} \wedge \partial X \rangle - \langle b, \partial X \wedge \bar{\partial} X \rangle).$$

Same formulas express symmetries. If $\{g^{ij}\}$ is nondegenerate, then :

$$S_{so} = \frac{1}{4\pi h} \int d^2 z (G_{\mu\nu} + B_{\mu\nu}) \partial X^\mu \bar{\partial} X^\nu,$$

$$G_{s\bar{k}} = g_{ij} \bar{\mu}_s^i \mu_{\bar{k}}^j + g_{s\bar{k}} - b_{s\bar{k}}, \quad B_{s\bar{k}} = g_{ij} \bar{\mu}_s^i \mu_{\bar{k}}^j - g_{s\bar{k}} - b_{s\bar{k}}$$

$$G_{si} = -g_{ij} \bar{\mu}_s^j - g_{s\bar{j}} \bar{\mu}_i^j, \quad G_{\bar{s}\bar{i}} = -g_{\bar{s}j} \mu_i^j - g_{ij} \mu_{\bar{s}}^j$$

$$B_{si} = g_{s\bar{j}} \bar{\mu}_i^j - g_{ij} \bar{\mu}_s^j, \quad B_{\bar{s}\bar{i}} = g_{ij} \mu_{\bar{s}}^j - g_{\bar{s}j} \mu_i^j,$$

Symmetries: infinitesimal diffeomorphism transformations and the 2-form B symmetry

$$G \rightarrow G - L_v G, \quad B \rightarrow B - L_v B$$

$$B \rightarrow B - 2d\omega$$

if $\alpha = (\mathbf{v}, \omega)$, so that $\mathbf{v} \in \Gamma(TM)$, $\omega \in \Omega^1(M)$.

The CFT corresponding to the chiral part of the free first order action

$$S_0 = \frac{1}{2\pi i h} \int_{\Sigma} \langle p \wedge \bar{\partial} X \rangle - \int R^{(2)}(\gamma) \phi'(X)$$

is locally described via VOA:

$$X^i(z) p_j(w) \sim \frac{h \delta_j^i}{z - w}$$

with Virasoro element

$$T(z) = \frac{1}{h} : \langle p(z) \partial X(z) \rangle : + \partial^2 \phi'(X(z)).$$

The corresponding space of states $V = \sum_{n=0}^{+\infty} V_n$,

$$V_0 \rightarrow \mathcal{O}_M \otimes \mathbb{C}[h] = \mathcal{O}_M^h, \quad V_1 \rightarrow \mathcal{V} = \mathcal{O}(\mathcal{E}) \otimes \mathbb{C}[h] \equiv \mathcal{O}(\mathcal{E})^h$$

The structure of vertex algebra imposes algebraic relations on $V_0 \oplus V_1$ giving it a structure of a *vertex algebra*.

[Outline](#)[Sigma-models and
conformal invariance
conditions](#)[Reformulation of
Sigma-model in
first-order form](#)[Vertex algebras,
 \$G_\infty\$ -algebra and
quasiclassical limit](#)[Einstein Equations](#)

A *vertex \mathcal{O}_M -algebroid* is a sheaf of \mathbb{C} -vector spaces \mathcal{V} with a pairing $\mathcal{O}_M \otimes_{\mathbb{C}[h]} \mathcal{V} \rightarrow \mathcal{V}$, i.e. $f \otimes v \mapsto f * v$ such that $1 * v = v$, equipped with a structure of a Leibniz $\mathbb{C}[h]$ -algebra $[\cdot, \cdot] : \mathcal{V} \otimes_{\mathbb{C}[h]} \mathcal{V} \rightarrow \mathcal{V}$, a $\mathbb{C}[h]$ -linear map of Leibniz algebras $\pi : \mathcal{V} \rightarrow \Gamma(TM)$ usually referred to as an anchor, a symmetric \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{V} \otimes_{\mathbb{C}[h]} \mathcal{V} \rightarrow \mathcal{O}_M^h$ a \mathbb{C} -linear map $\partial : \mathcal{O}_M \rightarrow \mathcal{V}$ such that $\pi \circ \partial = 0$, which satisfy the relations

$$f * (g * v) - (fg) * v = \pi(v)(f) * \partial(g) + \pi(v)(g) * \partial(f),$$

$$[v_1, f * v_2] = \pi(v_1)(f) * v_2 + f * [v_1, v_2],$$

$$[v_1, v_2] + [v_2, v_1] = \partial(\langle v_1, v_2 \rangle), \quad \pi(f * v) = f\pi(v),$$

$$\langle f * v_1, v_2 \rangle = f\langle v_1, v_2 \rangle - \pi(v_1)(\pi(v_2)(f)),$$

$$\pi(v)(\langle v_1, v_2 \rangle) = \langle [v, v_1], v_2 \rangle + \langle v_1, [v, v_2] \rangle,$$

$$\partial(fg) = f * \partial(g) + g * \partial(f),$$

$$[v, \partial(f)] = \partial(\pi(v)(f)), \quad \langle v, \partial(f) \rangle = \pi(v)(f),$$

where $v, v_1, v_2 \in \mathcal{V}$, $f, g \in \mathcal{O}_M^h$.

$$\partial f = df, \quad \pi(v)f = -hv(f), \quad \pi(\omega) = 0,$$

$$f * v = fv + h dX^i \partial_i \partial_j f v^j, \quad f * \omega = f\omega,$$

$$[v_1, v_2] = -h[v_1, v_2]_D - h^2 dX^i \partial_i \partial_k v_1^s \partial_s v_2^k,$$

$$[v, \omega] = -h[v, \omega]_D, \quad [\omega, v] = -h[\omega, v]_D, \quad [\omega_1, \omega_2] = 0,$$

$$\langle v, \omega \rangle = -h\langle v, \omega \rangle^s, \quad \langle v_1, v_2 \rangle = -h^2 \partial_i v_1^j \partial_j v_2^i, \quad \langle \omega_1, \omega_2 \rangle = 0,$$

Given a holomorphic volume form on open neighborhood U of M , one can associate a homotopy Gerstenhaber algebra to the vertex algebroid on U .

Consider the light modes of the corresponding BRST complex, (i.e. $L_0 = 0$). The resulting complex (\mathcal{F}_h, Q) is:

$$\begin{array}{ccccc}
 & \mathcal{V} & & \mathcal{V} & \\
 \nearrow \partial & \oplus & \nwarrow \partial & \nearrow \partial & \searrow \frac{1}{2} h \text{div} \\
 \mathcal{O}_M^h & & \mathcal{O}_M^h & \xrightarrow{id} & \mathcal{O}_M^h \\
 & & \nwarrow \frac{1}{2} h \text{div} & &
 \end{array}$$

div stands for divergence operator with respect to the nonvanishing volume form applied to sections of $\Gamma(U, T^{(1,0)}(M))$.

The homotopy Gerstenhaber algebra of Lian and Zuckerman

The homotopy associative and homotopy commutative product of Lian and Zuckerman:

$$(A, B) = \text{Res}_z \frac{A(z)B}{z}$$

$$\begin{aligned} Q(a_1, a_2)_h &= (Qa_1, a_2)_h + (-1)^{|a_1|}(a_1, Qa_2)_h, \\ (a_1, a_2)_h - (-1)^{|a_1||a_2|}(a_2, a_1)_h &= \\ Qm(a_1, a_2) + m(Qa_1, a_2) + (-1)^{|a_1|}m(a_1, Qa_2), \\ Q(a_1, a_2, a_3)_h + (Qa_1, a_2, a_3)_h + (-1)^{|a_1|}(a_1, Qa_2, a_3)_h + \\ (-1)^{|a_1|+|a_2|}(a_1, a_2, Qa_3)_h &= ((a_1, a_2)_h, a_3)_h - (a_1, (a_2, a_3)_h)_h \end{aligned}$$

Operator \mathbf{b} of degree -1 on (\mathcal{F}_h, Q) which anticommutes with Q :

$$\begin{array}{ccc} \mathcal{V} & \xleftarrow{-id} & \mathcal{V} \\ \oplus & & \oplus \\ \mathcal{O}_M^h & \xleftarrow{id} & \mathcal{O}_M^h \end{array} \quad \mathcal{O}_M^h \xleftarrow{-id} \mathcal{O}_M^h$$

One can define a bracket:

$$(-1)^{|a_1|} \{a_1, a_2\}_h = \mathbf{b}(a_1, a_2)_h - (\mathbf{b}a_1, a_2)_h - (-1)^{|a_1|} (a_1 \mathbf{b}a_2)_h,$$


$$\begin{aligned} & \{a_1, a_2\} + (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, a_1\} = \\ & (-1)^{|a_1|-1} (Qm'_h(a_1, a_2) - m'_h(Qa_1, a_2) - (-1)^{|a_2|} m'_h(a_1, Qa_2)), \\ & \{a_1, (a_2, a_3)_h\}_h = (\{a_1, a_2\}_h, a_3)_h + (-1)^{(|a_1|-1)|a_2|} (a_2, \{a_1, a_3\}_h)_h, \\ & \{(a_1, a_2)_h, a_3\}_h - (a_1, \{a_2, a_3\}_h)_h - (-1)^{(|a_3|-1)|a_2|} (\{a_1, a_3\}_h, a_2)_h = \\ & (-1)^{|a_1|+|a_2|-1} (Qn'_h(a_1, a_2, a_3) - n'_h(Qa_1, a_2, a_3) - \\ & (-1)^{|a_1|} n'_h(a_1, Qa_2, a_3) - (-1)^{|a_1|+|a_2|} n'_h(a_1, a_2, Qa_3), \\ & \{\{a_1, a_2\}_h, a_3\}_h - \{a_1, \{a_2, a_3\}_h\}_h + \\ & (-1)^{(|a_1|-1)(|a_2|-1)} \{a_2, \{a_1, a_3\}_h\}_h = 0. \end{aligned}$$

The conjecture of Lian and Zuckerman, which was later proven by series of papers (Kimura, Zuckerman, Voronov; Huang, Zhao; Voronov) says that the symmetrized product and bracket of homotopy Gerstenhaber algebra constructed above can be lifted to G_∞ -algebra.

Quasiclassical limit of LZ G_∞ algebra

Let $\mathcal{V}|_{h=0} = \mathcal{V}^0$.

One can see that $(\mathcal{F}, Q) \cong (\mathcal{F}_1, Q)$ is a subcomplex of (\mathcal{F}_h, Q) , which is:



Then

$$(\cdot, \cdot)_h : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow \mathcal{F}^{i+j}[h], \quad \{\cdot, \cdot\} : \mathcal{F}^i \otimes \mathcal{F}^j \rightarrow h\mathcal{F}_{i+j-1}[h],$$

$$\mathbf{b} : \mathcal{F}^i \rightarrow h\mathcal{F}^{i-1}[h],$$

so that

$$(\cdot, \cdot)_0 = \lim_{h \rightarrow 0} (\cdot, \cdot)_h, \quad \{\cdot, \cdot\}_0 = \lim_{h \rightarrow 0} h^{-1} \{\cdot, \cdot\}_h, \quad \mathbf{b}_0 = \lim_{h \rightarrow 0} h^{-1} \mathbf{b}$$

are well defined.

The symmetrized operations $(\cdot, \cdot)_0$, $\{\cdot, \cdot\}_0$ satisfy the relations of homotopy Gerstenhaber algebra. The resulting C_∞ and L_∞ algebras are reduced to C_3 and L_3 algebras.

Conjecture: This G_∞ -algebra is G_3 -algebra (no higher homotopies).

This is very close to the classical limit procedure for vertex algebroid:

$$[\cdot, \cdot]_0 = \lim_{h \rightarrow 0} \frac{1}{h} [\cdot, \cdot], \quad \pi_0 = \lim_{h \rightarrow 0} \frac{1}{h} \pi, \quad \langle \cdot, \cdot \rangle_0 = \frac{1}{h} \langle \cdot, \cdot \rangle.$$

As a result we get a Courant algebroid:

A Courant \mathcal{O}_M -algebroid is an \mathcal{O}_M -module \mathcal{Q} equipped with a structure of a Leibniz \mathbb{C} -algebra $[\cdot, \cdot]_0 : \mathcal{Q} \otimes_{\mathbb{C}} \mathcal{Q} \rightarrow \mathcal{Q}$, an \mathcal{O}_M -linear map of Leibniz algebras (the anchor map) $\pi_0 : \mathcal{Q} \rightarrow \Gamma(TM)$, a symmetric \mathcal{O}_M -bilinear pairing $\langle \cdot, \cdot \rangle : \mathcal{Q} \otimes_{\mathcal{O}_M} \mathcal{Q} \rightarrow \mathcal{O}_M$, a derivation $\partial : \mathcal{O}_M \rightarrow \mathcal{Q}$ which satisfy


$$\begin{aligned} \pi \circ \partial &= 0, \quad [q_1, f q_2]_0 = f [q_1, q_2]_0 + \pi_0(q_1)(f) q_2 \\ \langle [q, q_1], q_2 \rangle + \langle q_1, [q, q_2] \rangle &= \pi_0(q)(\langle q_1, q_2 \rangle_0), \\ [q, \partial(f)]_0 &= \partial(\pi_0(q)(f)) \\ \langle q, \partial(f) \rangle &= \pi_0(q)(f) \quad [q_1, q_2]_0 + [q_2, q_1]_0 = \partial(\langle q_1, q_2 \rangle_0) \end{aligned}$$

for $f \in \mathcal{O}_M$ and $q, q_1, q_2 \in \mathcal{Q}$. In our case $\mathcal{Q} \cong \mathcal{O}(\mathcal{E})$, π_0 is just a projection on $\mathcal{O}(TM)$

$$[q_1, q_2]_0 = -[q_1, q_2]_D, \quad \langle q_1, q_2 \rangle_0 = -\langle q_1, q_2 \rangle^s, \quad \partial = d.$$

Question: Is there a direct path (avoiding vertex algebra) from Courant algebroid to G_3 -algebra?

Subcomplex (\mathcal{F}_{sm}, Q) :



The G_∞ algebra degenerates to G -algebra. Moreover, due to \mathbf{b}_0 it is a BV-algebra. Combine chiral and antichiral part:

$$\mathbf{F}_{sm} = \mathcal{F}_{sm} \otimes \bar{\mathcal{F}}_{sm}$$

$$(-1)^{|a_1|} \{a_1, a_2\} = \mathbf{b}^-(a_1, a_2) - (\mathbf{b}^- a_1, a_2) - (-1)^{|a_1|} (a_1 \mathbf{b}^- a_2),$$

where $\mathbf{b}^- = \mathbf{b} - \bar{\mathbf{b}}$.

Maurer-Cartan elements, closed under \mathbf{b}^- :

$$\Gamma(T^{(1,0)}(M) \otimes T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(0,1)}(M)) \oplus \mathcal{O}(T^{(1,0)}(M)) \oplus \mathcal{O}_M \oplus \bar{\mathcal{O}}_M$$

Components: $(g, \bar{v}, v, \phi, \bar{\phi})$.

The Maurer-Cartan equation is equivalent to:

- 1). Vector field $\text{div}_\Omega g$, where $\Omega = \Omega' e^{-2\phi+2\bar{\phi}}$ is determined by $f \equiv -2\Phi_0 = -2(\Phi'_0 + \phi - \bar{\phi})$ and $\partial_i \partial_{\bar{j}} \Phi_0 = 0$, is such that its $\Gamma(T^{(1,0)}M)$, $\Gamma(T^{(0,1)}M)$ components are correspondingly holomorphic and antiholomorphic.
- 2). Bivector field $g \in \Gamma(T^{(1,0)}M \otimes T^{(0,1)}M)$ obeys the following equation:

$$[[g, g]] + \mathcal{L}_{\text{div}_\Omega(g)} g = 0,$$

where $\mathcal{L}_{\text{div}_\Omega(g)}$ is a Lie derivative with respect to the corresponding vector fields and

$$[[g, h]]^{k\bar{l}} \equiv (g^{i\bar{j}} \partial_i \partial_{\bar{j}} h^{k\bar{l}} + h^{i\bar{j}} \partial_i \partial_{\bar{j}} g^{k\bar{l}} - \partial_i g^{k\bar{j}} \partial_{\bar{j}} h^{i\bar{l}} - \partial_i h^{k\bar{j}} \partial_{\bar{j}} g^{i\bar{l}})$$

- 3). $\text{div}_\Omega \text{div}_\Omega(g) = 0$.

The infinitesimal symmetries of the Maurer-Cartan equation coincide with the holomorphic coordinate transformations of the volume form and tensor $\{g^{i\bar{j}}\}$.

These are Einstein equations with the following constraints:

$$\begin{aligned} G_{i\bar{k}} &= g_{i\bar{k}}, & B_{i\bar{k}} &= -g_{i\bar{k}}, & \Phi &= \log \sqrt{g} + \Phi_0, \\ G_{ik} &= G_{i\bar{k}} = G_{ik} = G_{i\bar{k}} = 0, \end{aligned}$$

Physically:

$$\begin{aligned} &\int [dp][d\bar{p}][dX][d\bar{X}] e^{-\frac{1}{2\pi i\hbar} \int_{\Sigma} (\langle p \wedge \bar{\partial} X \rangle - \langle \bar{p} \wedge \partial X \rangle - \langle g, p \wedge \bar{p} \rangle) + \int_{\Sigma} R^{(2)}(\gamma) \Phi_0(X)} = \\ &\int [dX][d\bar{X}] e^{\frac{-1}{4\pi\hbar} \int d^2z (G_{\mu\nu} + B_{\mu\nu}) \partial X^{\mu} \bar{\partial} X^{\nu} + \int R^{(2)}(\gamma) (\Phi_0(X) + \sqrt{g})} \end{aligned}$$

Consider

$$\mathbf{F}_{b-} = \mathcal{F} \otimes \bar{\mathcal{F}}|_{b-=0}$$

with the L_∞ -algebra structure given by Lian-Zuckerman construction.
One can explicitly check that GMC symmetry

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda]_h + \frac{1}{2}[\Psi, \Psi, \Lambda]_h + \dots,$$

reproduces

$$\mathbb{M} \rightarrow \mathbb{M} - D\alpha + \phi_1(\alpha, \mathbb{M}) + \phi_2(\alpha, \mathbb{M}, \mathbb{M}).$$

up to the second order in \mathbb{M} .

Conjecture: The corresponding Maurer-Cartan equation gives Einstein equations on G, B, Φ expressed in terms of Beltrami-Courant differential. The symmetries of the Maurer-Cartan equation reproduce mentioned above symmetries of Einstein equations.

Outline

Sigma-models and
conformal invariance
conditions

Reformulation of
Sigma-model in
first-order form

Vertex algebroids,
 G_∞ -algebra and
quasiclassical limit

Einstein Equations

Thank you!