

Algebraic and geometric structures of mathematical physics

Anton M. Zeitlin

Columbia University, Department of Mathematics

University of Nottingham

Nottingham

May 27, 2016



Outline

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

I work in Representation Theory with applications in Algebra, Geometry, Topology and Mathematical Physics.

The "gameplay" of modern mathematics is the interplay between algebra and geometry. In particular, this applies to mathematics emerging from mathematical physics problems.

The famous examples involve mirror symmetry, Khovanov homology, topological quantum field theories, geometric Langlands correspondence, umbral moonshine, which target a lot of areas of mathematics, from algebraic topology to number theory.

Outline

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

I work in Representation Theory with applications in Algebra, Geometry, Topology and Mathematical Physics.

The "gameplay" of modern mathematics is the interplay between algebra and geometry. In particular, this applies to mathematics emerging from mathematical physics problems.

The famous examples involve mirror symmetry, Khovanov homology, topological quantum field theories, geometric Langlands correspondence, umbral moonshine, which target a lot of areas of mathematics, from algebraic topology to number theory.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

I work in Representation Theory with applications in Algebra, Geometry, Topology and Mathematical Physics.

The "gameplay" of modern mathematics is the interplay between algebra and geometry. In particular, this applies to mathematics emerging from mathematical physics problems.

The famous examples involve mirror symmetry, Khovanov homology, topological quantum field theories, geometric Langlands correspondence, umbral moonshine, which target a lot of areas of mathematics, from algebraic topology to number theory.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

In this presentation:

Coordinates on super-Teichmüller spaces

Homotopy algebras, Conformal Field Theory, and Einstein equations

Towards the continuous Kazhdan-Lusztig correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶ $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶ $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline

Super-Teichmüller spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶ $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶ $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline

Super-Teichmüller spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶ $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶ $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶ $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶ $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ {complex structures on F }/isotopy
- ▶ {conformal structures on F }/isotopy
- ▶ {hyperbolic structures on F }/isotopy

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Coordinates on Super-Teichmüller spaces

Let $F_s^g \equiv F$ be the Riemann surface of genus g and s punctures.

Teichmüller space $T(F)$ has many incarnations:

- ▶ $\{\text{complex structures on } F\}/\text{isotopy}$
- ▶ $\{\text{conformal structures on } F\}/\text{isotopy}$
- ▶ $\{\text{hyperbolic structures on } F\}/\text{isotopy}$

Representation-theoretic definition:

$$T(F) = \text{Hom}'(\pi_1(F), PSL(2, \mathbb{R}))/PSL(2, \mathbb{R}).$$

The mapping class group $MC(F)$: group of homotopy classes of orientation preserving homeomorphisms. It acts on $T(F)$ by outer automorphisms of $\pi_1(F)$.

The primary object of interest is the *moduli space*:

$$M(F) = T(F)/MC(F).$$

The goal is to find system of coordinates on $T(F)$, so that $MC(F)$ is realized in the simplest possible way.

Outline


Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence


Extra topics

Penner's work in the 1980s: a construction of coordinates associated to the ideal triangulation of F (assume $s > 0$ and $2 - 2g - s < 0$):




so that one assigns one positive number for every edge.

The action of $MC(F)$ can be described combinatorially using elementary transformations called flips:




Ptolemy relation: $ef = ac + bd$.

Penner's work in the 1980s: a construction of coordinates associated to the ideal triangulation of F (assume $s > 0$ and $2 - 2g - s < 0$):




so that one assigns one positive number for every edge.

The action of $MC(F)$ can be described combinatorially using elementary transformations called flips:




Ptolemy relation: $ef = ac + bd$.

Penner's work in the 1980s: a construction of coordinates associated to the ideal triangulation of F (assume $s > 0$ and $2 - 2g - s < 0$):



so that one assigns one positive number for every edge.

The action of $MC(F)$ can be described combinatorially using elementary transformations called flips:



Ptolemy relation: $ef = ac + bd$.

Change of coordinates via the change of triangulation is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by S. Fomin and A. Zelevinsky in the early 2000s.

Penner's coordinates can be used for the quantization of $T(F)$ (V. Fock, R. Kashaev, late 90s, early 2000s).

Higher (super)Teichmüller spaces: $PSL(2, \mathbb{R})$ is replaced by some reductive (super)group G . In the case of reductive groups G the construction of coordinates was given by V.Fock and A. Goncharov (2003).

Change of coordinates via the change of triangulation is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by S. Fomin and A. Zelevinsky in the early 2000s.

Penner's coordinates can be used for the quantization of $T(F)$ (V. Fock, R. Kashaev, late 90s, early 2000s).

Higher (super)Teichmüller spaces: $PSL(2, \mathbb{R})$ is replaced by some reductive (super)group G . In the case of reductive groups G the construction of coordinates was given by V.Fock and A. Goncharov (2003).

Change of coordinates via the change of triangulation is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by S. Fomin and A. Zelevinsky in the early 2000s.

Penner's coordinates can be used for the quantization of $T(F)$ (V. Fock, R. Kashaev, late 90s, early 2000s).

Higher (super)Teichmüller spaces: $PSL(2, \mathbb{R})$ is replaced by some reductive (super)group G . In the case of reductive groups G the construction of coordinates was given by V.Fock and A. Goncharov (2003).

Change of coordinates via the change of triangulation is therefore governed by Ptolemy relations. This leads to the prominent geometric example of *cluster algebra*, introduced by S. Fomin and A. Zelevinsky in the early 2000s.

Penner's coordinates can be used for the quantization of $T(F)$ (V. Fock, R. Kashaev, late 90s, early 2000s).

Higher (super)Teichmüller spaces: $PSL(2, \mathbb{R})$ is replaced by some reductive (super)group G . In the case of reductive groups G the construction of coordinates was given by V.Fock and A. Goncharov (2003).

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Basic objects in superstring theory are:

$N = 1$ and $N = 2$ Super-Teichmüller spaces $ST(F)$, related to supergroups $OSP(1|2)$, $SL(1|2)$ correspondingly. In the late 80s the problem of construction of Penner's coordinates on $ST(F)$ was introduced on Yu.I. Manin's Moscow seminar.

The $N=1$ case was solved nearly 30 years later in:

R. Penner, A. Zeitlin, [arXiv:1509.06302](https://arxiv.org/abs/1509.06302).

The $N=2$ case is solved in collaboration with

I. Ip and R. Penner (in [arXiv](https://arxiv.org/abs/1509.06302) in a few weeks).

Further directions of study:

- ▶ Cluster algebras with anticommuting variables
- ▶ Quantization of super-Teichmüller spaces (first attempt by J. Teschner et al. [arXiv:1512.02617](https://arxiv.org/abs/1512.02617))
- ▶ Application to supermoduli theory and calculation of superstring amplitudes, which are highly nontrivial due to recent results of R. Donagi and E. Witten
- ▶ Higher super-Teichmüller theory for supergroups of higher rank

Homotopy algebras, Conformal Field Theory and Einstein equations

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

**Homotopy algebras,
CFT, and Einstein
equations**

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Early days of String Theory (the 1980s):

Einstein equations with matter fields emerged as consistency conditions for two-dimensional models.

At the same time it was discovered that linearized Einstein equations and their symmetries can be represented in terms of formalism of homological algebra:

$$Q\Psi = 0, \quad \Psi \rightarrow \Psi + Q\Lambda,$$

where Q ($Q^2 = 0$) is the differential in certain complex, known to physicists as BRST complex.

Homotopy algebras, Conformal Field Theory and Einstein equations

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

**Homotopy algebras,
CFT, and Einstein
equations**

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Early days of String Theory (the 1980s):

Einstein equations with matter fields emerged as consistency conditions for two-dimensional models.

At the same time it was discovered that linearized Einstein equations and their symmetries can be represented in terms of formalism of homological algebra:

$$Q\Psi = 0, \quad \Psi \rightarrow \Psi + Q\Lambda,$$

where Q ($Q^2 = 0$) is the differential in certain complex, known to physicists as BRST complex.

Homotopy algebras, Conformal Field Theory and Einstein equations

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

**Homotopy algebras,
CFT, and Einstein
equations**

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Early days of String Theory (the 1980s):

Einstein equations with matter fields emerged as consistency conditions for two-dimensional models.

At the same time it was discovered that linearized Einstein equations and their symmetries can be represented in terms of formalism of homological algebra:

$$Q\Psi = 0, \quad \Psi \rightarrow \Psi + Q\Lambda,$$

where Q ($Q^2 = 0$) is the differential in certain complex, known to physicists as BRST complex.

Homotopy algebras, Conformal Field Theory and Einstein equations

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

**Homotopy algebras,
CFT, and Einstein
equations**

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Early days of String Theory (the 1980s):

Einstein equations with matter fields emerged as consistency conditions for two-dimensional models.

At the same time it was discovered that linearized Einstein equations and their symmetries can be represented in terms of formalism of homological algebra:

$$Q\Psi = 0, \quad \Psi \rightarrow \Psi + Q\Lambda,$$

where Q ($Q^2 = 0$) is the differential in certain complex, known to physicists as BRST complex.

Homotopy algebras, Conformal Field Theory and Einstein equations

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

**Homotopy algebras,
CFT, and Einstein
equations**

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Early days of String Theory (the 1980s):

Einstein equations with matter fields emerged as consistency conditions for two-dimensional models.

At the same time it was discovered that linearized Einstein equations and their symmetries can be represented in terms of formalism of homological algebra:

$$Q\Psi = 0, \quad \Psi \rightarrow \Psi + Q\Lambda,$$

where Q ($Q^2 = 0$) is the differential in certain complex, known to physicists as BRST complex.

It was conjectured in the 90s (A. Sen, B. Zwiebach,...) that Einstein equations and their symmetries are Generalized Maurer-Cartan equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi] + \frac{1}{3!}[\Psi, \Psi, \Psi] + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda] + \frac{1}{2}[\Psi, \Psi, \Lambda] + \dots,$$

where polylinear operations satisfy quadratic relations and generate L_∞ -algebra.

The above L_∞ algebra structure involves perturbative expansion around flat metric (local, destroys geometry).

Claim:

There is a much deeper structure, called G_∞ -algebra (defined by D. Tamarkin and B. Tsygan in the early 2000s), governing Einstein equations.

The corresponding L_∞ algebra structure does not involve any perturbative expansion of metric.

It was conjectured in the 90s (A. Sen, B. Zwiebach,...) that Einstein equations and their symmetries are Generalized Maurer-Cartan equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi] + \frac{1}{3!}[\Psi, \Psi, \Psi] + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda] + \frac{1}{2}[\Psi, \Psi, \Lambda] + \dots,$$

where polylinear operations satisfy quadratic relations and generate L_∞ -algebra.

The above L_∞ algebra structure involves perturbative expansion around flat metric (local, destroys geometry).

Claim:

There is a much deeper structure, called G_∞ -algebra (defined by D. Tamarkin and B. Tsygan in the early 2000s), governing Einstein equations.

The corresponding L_∞ algebra structure does not involve any perturbative expansion of metric.

It was conjectured in the 90s (A. Sen, B. Zwiebach,...) that Einstein equations and their symmetries are Generalized Maurer-Cartan equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi] + \frac{1}{3!}[\Psi, \Psi, \Psi] + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda] + \frac{1}{2}[\Psi, \Psi, \Lambda] + \dots,$$

where polylinear operations satisfy quadratic relations and generate L_∞ -algebra.

The above L_∞ algebra structure involves perturbative expansion around flat metric (local, destroys geometry).

Claim:

There is a much deeper structure, called G_∞ -algebra (defined by D. Tamarkin and B. Tsygan in the early 2000s), governing Einstein equations.

The corresponding L_∞ algebra structure does not involve any perturbative expansion of metric.

It was conjectured in the 90s (A. Sen, B. Zwiebach,...) that Einstein equations and their symmetries are Generalized Maurer-Cartan equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi] + \frac{1}{3!}[\Psi, \Psi, \Psi] + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda] + \frac{1}{2}[\Psi, \Psi, \Lambda] + \dots,$$

where polylinear operations satisfy quadratic relations and generate L_∞ -algebra.

The above L_∞ algebra structure involves perturbative expansion around flat metric (local, destroys geometry).

Claim:

There is a much deeper structure, called G_∞ -algebra (defined by D. Tamarkin and B. Tsygan in the early 2000s), governing Einstein equations.

The corresponding L_∞ algebra structure does not involve any perturbative expansion of metric.

It was conjectured in the 90s (A. Sen, B. Zwiebach,...) that Einstein equations and their symmetries are Generalized Maurer-Cartan equations and their symmetries:

$$Q\Psi + \frac{1}{2}[\Psi, \Psi] + \frac{1}{3!}[\Psi, \Psi, \Psi] + \dots = 0$$

$$\Psi \rightarrow \Psi + Q\Lambda + [\Psi, \Lambda] + \frac{1}{2}[\Psi, \Psi, \Lambda] + \dots,$$

where polylinear operations satisfy quadratic relations and generate L_∞ -algebra.

The above L_∞ algebra structure involves perturbative expansion around flat metric (local, destroys geometry).

Claim:

There is a much deeper structure, called G_∞ -algebra (defined by D. Tamarkin and B. Tsygan in the early 2000s), governing Einstein equations.

The corresponding L_∞ algebra structure does not involve any perturbative expansion of metric.

The construction involves two important mathematical objects:

- ▶ Sheaves of vertex operator algebras
- ▶ Courant algebroids

Sheaves of vertex operator algebras (VOA) is a mathematical tool to describe certain two-dimensional physical models (2d conformal field theories) on a given manifold M (introduced by V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob in the early 2000s).

Courant algebroids are now primarily related to the subject called "Generalized Geometry", introduced by N. Hitchin in the late 90s.

Important object is the Courant bracket:

in the simplest case it is the generalization of standard Lie bracket on the sections of direct sum of tangent and cotangent bundles of M .

Unlike the Lie bracket, Courant bracket satisfies Jacobi identity only up to homotopy.

A. Losev, A. Marshakov, A. Zeitlin, Phys. Lett. B, 633, 375-381, 2006;
A. Zeitlin, Nucl. Phys. B, 794 381-401, 2008;
A. Zeitlin, J. Phys. A, 42, 355401-355412, 2009;
A. Zeitlin, Comm. Math. Phys. 303, 331-359, 2011;
A. Zeitlin, Contemp. Math., 623, 267-280, 2014;
A. Zeitlin, Adv.Theor.Math.Phys., 19, 1249-1275, 2015

The construction involves two important mathematical objects:

- ▶ Sheaves of vertex operator algebras
- ▶ Courant algebroids

Sheaves of vertex operator algebras (VOA) is a mathematical tool to describe certain two-dimensional physical models (2d conformal field theories) on a given manifold M (introduced by V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob in the early 2000s).

Courant algebroids are now primarily related to the subject called "Generalized Geometry", introduced by N. Hitchin in the late 90s.

Important object is the Courant bracket:

in the simplest case it is the generalization of standard Lie bracket on the sections of direct sum of tangent and cotangent bundles of M .

Unlike the Lie bracket, Courant bracket satisfies Jacobi identity only up to homotopy.

A. Losev, A. Marshakov, A. Zeitlin, Phys. Lett. B, 633, 375-381, 2006;

A. Zeitlin, Nucl. Phys. B, 794 381-401, 2008;

A. Zeitlin, J. Phys. A, 42, 355401-355412, 2009;

A. Zeitlin, Comm. Math. Phys. 303, 331-359, 2011;

A. Zeitlin, Contemp. Math., 623, 267-280, 2014;

A. Zeitlin, Adv.Theor.Math.Phys., 19, 1249-1275, 2015

The construction involves two important mathematical objects:

- ▶ Sheaves of vertex operator algebras
- ▶ Courant algebroids

Sheaves of vertex operator algebras (VOA) is a mathematical tool to describe certain two-dimensional physical models (2d conformal field theories) on a given manifold M (introduced by V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob in the early 2000s).

Courant algebroids are now primarily related to the subject called "Generalized Geometry", introduced by N. Hitchin in the late 90s.

Important object is the Courant bracket:

in the simplest case it is the generalization of standard Lie bracket on the sections of direct sum of tangent and cotangent bundles of M .

Unlike the Lie bracket, Courant bracket satisfies Jacobi identity only up to homotopy.

A. Losev, A. Marshakov, A. Zeitlin, Phys. Lett. B, 633, 375-381, 2006;

A. Zeitlin, Nucl. Phys. B, 794 381-401, 2008;

A. Zeitlin, J. Phys. A, 42, 355401-355412, 2009;

A. Zeitlin, Comm. Math. Phys. 303, 331-359, 2011;

A. Zeitlin, Contemp. Math., 623, 267-280, 2014;

A. Zeitlin, Adv.Theor.Math.Phys., 19, 1249-1275, 2015

The construction involves two important mathematical objects:

- ▶ Sheaves of vertex operator algebras
- ▶ Courant algebroids

Sheaves of vertex operator algebras (VOA) is a mathematical tool to describe certain two-dimensional physical models (2d conformal field theories) on a given manifold M (introduced by V. Gorbounov, F. Malikov, V. Schechtman, A. Vaintrob in the early 2000s).

Courant algebroids are now primarily related to the subject called "Generalized Geometry", introduced by N. Hitchin in the late 90s.

Important object is the Courant bracket:

in the simplest case it is the generalization of standard Lie bracket on the sections of direct sum of tangent and cotangent bundles of M .

Unlike the Lie bracket, Courant bracket satisfies Jacobi identity only up to homotopy.

A. Losev, A. Marshakov, A. Zeitlin, Phys. Lett. B, 633, 375-381, 2006;

A. Zeitlin, Nucl. Phys. B, 794 381-401, 2008;

A. Zeitlin, J. Phys. A, 42, 355401-355412, 2009;

A. Zeitlin, Comm. Math. Phys. 303, 331-359, 2011;

A. Zeitlin, Contemp. Math., 623, 267-280, 2014;

A. Zeitlin, Adv.Theor.Math.Phys., 19, 1249-1275, 2015

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

**Continuous
Kazhdan-Lusztig
correspondence**

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

**Continuous
Kazhdan-Lusztig
correspondence**

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

Towards the continuous Kazhdan-Lusztig correspondence

Algebraic and
geometric structures
of mathematical
physics

Anton Zeitlin

Outline

Super-Teichmüller
spaces

Homotopy algebras,
CFT, and Einstein
equations

Continuous
Kazhdan-Lusztig
correspondence

Extra topics

Kazhdan-Lusztig correspondence (the early 90s).

Equivalence between

- ▶ Braided tensor category of finite-dimensional representations of quantum group $U_q(\mathfrak{g})$ (algebraic),
- ▶ Braided tensor category of standard highest weight modules of affine Lie algebra \mathfrak{g} (geometric).

Physical origins of this correspondence come from 3d Chern-Simons theory for Lie group G_c , corresponding to the compact form \mathfrak{g}_c of \mathfrak{g} .

Correlation functions give knot polynomials (Jones polynomial for $\mathfrak{g}_c = \mathfrak{su}(2)$), which can be described via representation theory $U_q(\mathfrak{g}_c)$.

On the other hand, the boundary theory is Wess-Zumino-Witten (WZW) model, described by the representation theory of $\hat{\mathfrak{g}}_c$.

This juxtaposition of theories found applications in many exciting topics in mathematics and physics like

- ▶ Topological Field Theories
- ▶ Mirror Symmetry
- ▶ Khovanov Homology
- ▶ Geometry of Hilbert schemes

Currently physicists are interested in the understanding of Chern-Simons theory for noncompact real reductive Lie groups $G_{\mathbb{R}}$, e.g. for $SL(2, \mathbb{R})$.

In particular, $SL(2, \mathbb{R})$ Chern-Simons theory is related to 3d gravity, many aspects of which were studied in the research groups of J.W. Barrett and K. Krasnov.

This juxtaposition of theories found applications in many exciting topics in mathematics and physics like

- ▶ Topological Field Theories
- ▶ Mirror Symmetry
- ▶ Khovanov Homology
- ▶ Geometry of Hilbert schemes

Currently physicists are interested in the understanding of Chern-Simons theory for noncompact real reductive Lie groups $G_{\mathbb{R}}$, e.g. for $SL(2, \mathbb{R})$.

In particular, $SL(2, \mathbb{R})$ Chern-Simons theory is related to 3d gravity, many aspects of which were studied in the research groups of J.W. Barrett and K. Krasnov.

This juxtaposition of theories found applications in many exciting topics in mathematics and physics like

- ▶ Topological Field Theories
- ▶ Mirror Symmetry
- ▶ Khovanov Homology
- ▶ Geometry of Hilbert schemes

Currently physicists are interested in the understanding of Chern-Simons theory for noncompact real reductive Lie groups $G_{\mathbb{R}}$, e.g. for $SL(2, \mathbb{R})$.

In particular, $SL(2, \mathbb{R})$ Chern-Simons theory is related to 3d gravity, many aspects of which were studied in the research groups of J.W. Barrett and K. Krasnov.

This juxtaposition of theories found applications in many exciting topics in mathematics and physics like

- ▶ Topological Field Theories
- ▶ Mirror Symmetry
- ▶ Khovanov Homology
- ▶ Geometry of Hilbert schemes

Currently physicists are interested in the understanding of Chern-Simons theory for noncompact real reductive Lie groups $G_{\mathbb{R}}$, e.g. for $SL(2, \mathbb{R})$.

In particular, $SL(2, \mathbb{R})$ Chern-Simons theory is related to 3d gravity, many aspects of which were studied in the research groups of J.W. Barrett and K. Krasnov.

This juxtaposition of theories found applications in many exciting topics in mathematics and physics like

- ▶ Topological Field Theories
- ▶ Mirror Symmetry
- ▶ Khovanov Homology
- ▶ Geometry of Hilbert schemes

Currently physicists are interested in the understanding of Chern-Simons theory for noncompact real reductive Lie groups $G_{\mathbb{R}}$, e.g. for $SL(2, \mathbb{R})$.

In particular, $SL(2, \mathbb{R})$ Chern-Simons theory is related to 3d gravity, many aspects of which were studied in the research groups of J.W. Barrett and K. Krasnov.

One can use the following approach:

Construct WZW model on the boundary for $G_{\mathbb{R}}$. In other words, describe the braided tensor category of unitary modules for $\hat{\mathfrak{g}}_{\mathbb{R}}$.

i) Quantum group side of the story is known in this case.

Modular double representations of $U_q(\mathfrak{g}_{\mathbb{R}})$: L. Faddeev and J. Teschner (for $U_q(\mathfrak{sl}(2, \mathbb{R}))$) in the early 2000s; I.B. Frenkel and I. Ip recently generalized these formulas for higher rank. This is the proper quantum analogue of principal series representations.

ii) The corresponding modules for $\hat{\mathfrak{sl}}(2, \mathbb{R})$ were constructed in: I.B. Frenkel, A. Zeitlin, *Comm.Math.Phys.* 326, pp. 145-165, 2014 based on the earlier work A. Zeitlin, *J.Funct.Anal.* 263, pp. 529-548, 2012. The construction relies on the renormalization technique similar to the one used in field theory.

A recent review: A. Zeitlin, *Proc. Symp. Pure Math.*, 92, 2016

The description of tensor structure, braiding and the continuous version of KL correspondence for $\mathfrak{sl}(2, \mathbb{R})$ is currently in progress.

One can use the following approach:

Construct WZW model on the boundary for $G_{\mathbb{R}}$. In other words, describe the braided tensor category of unitary modules for $\hat{\mathfrak{g}}_{\mathbb{R}}$.

i) Quantum group side of the story is known in this case.

Modular double representations of $U_q(\mathfrak{g}_{\mathbb{R}})$: L. Faddeev and J. Teschner (for $U_q(\mathfrak{sl}(2, \mathbb{R}))$) in the early 2000s; I.B. Frenkel and I. Ip recently generalized these formulas for higher rank. This is the proper quantum analogue of principal series representations.

ii) The corresponding modules for $\widehat{\mathfrak{sl}}(2, \mathbb{R})$ were constructed in: I.B. Frenkel, A. Zeitlin, *Comm.Math.Phys.* 326, pp. 145-165, 2014 based on the earlier work A. Zeitlin, *J.Funct.Anal.* 263, pp. 529-548, 2012. The construction relies on the renormalization technique similar to the one used in field theory.

A recent review: A. Zeitlin, *Proc. Symp. Pure Math.*, 92, 2016

The description of tensor structure, braiding and the continuous version of KL correspondence for $\mathfrak{sl}(2, \mathbb{R})$ is currently in progress.

[Outline](#)[Super-Teichmüller spaces](#)[Homotopy algebras, CFT, and Einstein equations](#)[Continuous Kazhdan-Lusztig correspondence](#)[Extra topics](#)

One can use the following approach:

Construct WZW model on the boundary for $G_{\mathbb{R}}$. In other words, describe the braided tensor category of unitary modules for $\hat{\mathfrak{g}}_{\mathbb{R}}$.

i) Quantum group side of the story is known in this case.

Modular double representations of $U_q(\mathfrak{g}_{\mathbb{R}})$: L. Faddeev and J. Teschner (for $U_q(\mathfrak{sl}(2, \mathbb{R}))$) in the early 2000s; I.B. Frenkel and I. Ip recently generalized these formulas for higher rank. This is the proper quantum analogue of principal series representations.

ii) The corresponding modules for $\hat{\mathfrak{sl}}(2, \mathbb{R})$ were constructed in: I.B. Frenkel, A. Zeitlin, *Comm.Math.Phys.* 326, pp. 145-165, 2014 based on the earlier work A. Zeitlin, *J.Funct.Anal.* 263, pp. 529-548, 2012. The construction relies on the renormalization technique similar to the one used in field theory.

A recent review: A. Zeitlin, *Proc. Symp. Pure Math.*, 92, 2016

The description of tensor structure, braiding and the continuous version of KL correspondence for $\mathfrak{sl}(2, \mathbb{R})$ is currently in progress.

One can use the following approach:

Construct WZW model on the boundary for $G_{\mathbb{R}}$. In other words, describe the braided tensor category of unitary modules for $\hat{\mathfrak{g}}_{\mathbb{R}}$.

i) Quantum group side of the story is known in this case.

Modular double representations of $U_q(\mathfrak{g}_{\mathbb{R}})$: L. Faddeev and J. Teschner (for $U_q(\mathfrak{sl}(2, \mathbb{R}))$) in the early 2000s; I.B. Frenkel and I. Ip recently generalized these formulas for higher rank. This is the proper quantum analogue of principal series representations.

ii) The corresponding modules for $\widehat{\mathfrak{sl}}(2, \mathbb{R})$ were constructed in:

I.B. Frenkel, A. Zeitlin, *Comm.Math.Phys.* 326, pp. 145-165, 2014
based on the earlier work A. Zeitlin, *J.Funct.Anal.* 263, pp. 529-548,
2012. The construction relies on the renormalization technique similar
to the one used in field theory.

A recent review: A. Zeitlin, *Proc. Symp. Pure Math.*, 92, 2016

The description of tensor structure, braiding and the continuous version
of KL correspondence for $\mathfrak{sl}(2, \mathbb{R})$ is currently in progress.

- ▶ Quantum groups via semi-infinite cohomology,
I. Frenkel, A. Zeitlin,
J. Noncomm. Geom., 7, 1007-1026, 2013;
Comm.Math.Phys., 297, 687-732, 2010
- ▶ Geometry of compactified moduli spaces
and homotopy algebras of CFT,
A. Zeitlin,
Contemp. Math., 623, 267-280, 2014;
Int. J. Math., 23, 1250012, 2012
- ▶ Geometric Langlands correspondence for superalgebras.
Simplest example considered in:
A. Zeitlin, Lett. Math. Phys., 105, 149-167, 2015
- ▶ Integrable models of statistical physics via quantum K-theory,
P. Pushkar, A. Smirnov, A. Zeitlin, to appear in 2016