

Lab 4: Spectroscopy

Answer Key

point allocation is ultimately at instructor discretion - just use good judgement

BEFORE LAB: Instructors must set up 3 diffusion tubes and a light-bulb. Noor will help make sure these are easy for lab instructors to find. **TODO(Noor):** test out 3 good diffusion tubes and label what element they are

1 Introduction: The Rainbow Connection

Spectroscopy is the study of the light that objects emit or absorb. The Universe is filled with many "flavors" of light, spanning orders of magnitude in wavelength. Spectroscopy hones in on a specific object and studies the range of wavelengths emitted or absorbed by that object – the set of light emitted or absorbed by an object is called the object's *spectrum*. An object's spectrum can provide us a lot of useful information: the chemical makeup of the object, its motion and speed, temperature, and the distance (just to name a few). Thus, spectroscopy is an extremely popular technique for studying planets, stars, and galaxies. Much of what we know about these objects is *because* of spectroscopy.


2 Looking at Light

Let's look at some lamps (or, diffusion tubes)! If you do not already have a *diffraction grating* or *spectroscope*, come get one from me. **Keep track of your diffraction grating/spectroscope and ensure it is returned at the end of the lab.** A diffraction grating splits light into its component colors, much like a prism. There are several lamps set up in the room; each lamp is actually heated gas of a specific element. Go up to each lamp and look at it through your diffraction grating. **Please do not touch the lamps, they are hot!**

Refer to the "Spectra_Handout" PDF for help identifying the elements from their spectra.

1. [9 points] **Sketch each lamp's spectrum in your notebook.** Do all three lamps. For each one, write down the element and draw what you see.

Draw a horizontal strip for each lamp then mark roughly where the vertical lines fall. No need to color the lines, but try to describe the colors ("bright green," "two close reds").



Answer: Should look like a few separated lines, different for each lamp. Elements in the diffusion lamps still needs to be confirmed but, roughly: - Hydrogen: red 656, blue-green 486, violet 434 nm. - Helium: bright yellow 588, red 668, green 502 nm. - Neon: bunch of

red/orange lines. -Mercury: violet 436, green 546, yellow pair 577/579 nm.

Points: 3 points per lamp $\times$ 3 lamps — 1 pt for naming the element, 1 pt for a sketch showing distinct lines in roughly the right places, 1 pt for sketch annotation.

2. [2 points] Note the similarities and differences in these spectra.

Answer: Similar: all are narrow/thin bright lines. Different: how many, where they fall, their colors, their spacing.

Points: 1 pt for a similarity, 1 pt for a difference.

3. [1 point] Why do the lamps look different from each other through the grating? Choose an option from below.

- (a) Each lamp holds a different element, and each element emits its own set of wavelengths.
- (b) Each lamp is a different temperature.
- (c) Each lamp is a different brightness.

Answer: (a).

Points: 1 pt, it's either right or wrong.

4. [2 points] Now look at one of the light bulbs (a source of white light) in the room through your diffraction grating. Describe what you see. How is the light bulb's spectrum different from the lamps?

Answer: An complete/smooth/uninterrupted rainbow, every color is present with no gaps. The lamps gave only a few separate bright lines with nothing between them.

Points: 1 pt for saying a complete rainbow; 1 pt for the contrast with the lamps' emission/isolated lines.

3 Blackbody Radiation

The light bulb gave you an unbroken rainbow. Where does that rainbow come from, and what decides which color is brightest in it? To answer that, we'll consider the simplified, idealized model of a **blackbody**. A blackbody is an object that perfectly *absorbs* all wavelengths of light, with no reflection. In addition to absorbing all wavelengths of light, a perfect blackbody will also *emit* light at all wavelengths.

5. [3 points] Consider the processes of *emission*, *absorption*, and *reflection* (or “*scattering*”) of light. Define each one in your own words. A short phrase each is plenty — think about what happens to a beam of light when it meets an object.

Answer: Emission: the object gives off its own light. Absorption: light goes in and doesn't come back out. Reflection/scattering: light bounces off.

Points: 1 pt per correct definition (emission, absorption, reflection).

If one plots the intensity (or brightness) of light at each wavelength emitted by a blackbody, it yields the *blackbody spectrum*. Blackbody spectra are smooth - we will soon find out that stars and other astronomical objects are not, but we can usually approximate them as blackbodies!

For now, though, let's examine the blackbody spectrum hands-on. Navigate to https://phet.colorado.edu/sims/html/blackbody-spectrum/latest/blackbody-spectrum_en.html. The plot to the left of the screen shows a blackbody spectrum, with spectral power density (the amount of light emitted) on the vertical axis and wavelength on the horizontal axis; don't worry too much about these units. **Make sure to check the boxes for “Graph Values” and “Labels” in the box** to the upper right; after turning on these features, you should see a pair of yellow numbers labeling the *peak* of the blackbody curve.

To the right of the screen is a thermometer showing the temperature of the blackbody. You can click and drag the slider on the thermometer to change the temperature of the object. If you need to adjust the size of your graph to see the full curve, you can use the zoom buttons (the magnifying glasses) to the upper left and bottom right of the plot.

6. [3 points] **Make a table in your notebook** with four columns:

Temperature, T (K)	Peak wavelength, λ_{peak} (μm)	$1/\lambda_{\text{peak}}$ ($1/\mu\text{m}$)	$\lambda_{\text{peak}} \times T$ ($\mu\text{m K}$)
-------------------------	---	--	---

Test **at least 5** different temperatures spread across the full range of the thermometer, and fill in the first three columns. Three of your five rows should be for the ones marked on the thermometer itself: the **light bulb**, the **Sun**, and **Sirius A**. Use *two* significant figures. **Leave the last column empty for now.**

Answer: Final table:

T (K)	λ_{peak} (μm)	$1/\lambda_{\text{peak}}$	$\lambda_{\text{peak}} \times T$
300	9.7	0.10	2900
3000 (bulb)	0.97	1.0	2900
5800 (Sun)	0.50	2.0	2900
9950 (Sirius A)	0.29	3.4	2900
11000	0.26	3.8	2900

Can accept $\pm 5\%$ on λ_{peak} , and ± 50 K on temperature

Points: 1 pt for at least 5 temperatures (a.k.a 5 rows), 1 pt for including the bulb, the Sun and Sirius A; 1 pt for correct λ_{peak} values. If no 2 sig. fig, subtract 0.5 or 1 pt..

7. [2 points] Look at your first two columns. As temperature goes *up*, what happens to the peak wavelength? What is this type of relationship called?




Figure 1: Answer for question 9

Answer: Peak wavelength goes down as temperature goes up. Inverse relationship.

Points: 1 pt for (basically) saying peak wavelength decreases as temperature increases; 1 pt for naming it an inverse (or inversely proportional) relationship.

8. [3 points] **Plot your data:** $1/\lambda_{\text{peak}}$ on the vertical axis, temperature on the horizontal axis. **Label both axes with the appropriate units.** What shape do your points make?

Answer: A straight line.

Points: 1 pt for plotting the points correctly; 1 pt for both axes labelled with units; 1 pt for identifying the shape as a straight line.

9. [2 points] Now fill in the last column of your table ($\lambda_{\text{peak}} \times T$, two sig figs). What do you notice about those five values?.

Answer: All five products are about the same: $\approx 2900 \mu\text{m K}$. ± 100 should be fine

Points: 1 pt for computing the fourth column; 1 pt for noticing the values are all (roughly) the same constant.

Congratulations! You've just derived a key equation in astronomy, **Wien's law**. Because the product of the peak wavelength and the temperature is always the same number, we can determine the surface temperature of an object (like a star) by measuring the wavelength at which the object emits most of its light. In math, Wien's law reads

$$\lambda_{\text{peak}} \times T = (2.9 \times 10^3) \mu\text{m K}, \quad (1)$$

where λ_{peak} is the peak wavelength (in μm) and T is the temperature (in K). Let's use Wien's law to answer a few questions.

10. [1 point] Are redder objects hotter or cooler than bluer objects? (this applies to both the flames on our kitchen stoves and to the stars illuminating the night sky).

Answer: Cooler. Blue is hot, red is cool.

Points: 1 pt if correct

11. The surface temperature of the Sun is around 5780 K.

- (a) [2 points] At which wavelength does the Sun emit most of its light (give your answer in μm to two sig figs)? Show your work.
- (b) [1 point] What color of light does the Sun emit the most? (Note: you can use the tool from the previous section, too).

Answer: (a) $\lambda_{\text{peak}} = (2.9 \times 10^3)/5780 = 0.50 \mu\text{m} = 500 \text{ nm}$ (b) Green

Points: (a) 1 pt for showing the rearranged Wien's law / showing work, 1 pt for the correct value to two sig figs. (b) 1 pt for green.

4 Fingerprints of the Elements

The blackbody spectrum is a perfectly smooth spectrum — like the light bulb. In reality, however, objects (like stars and galaxies) possess many “features” — or *lines* — in their spectra, like the ones you sketched from the lamps, that can tell us much more than just the temperature of the object. Let's explore some real spectra now.

Figure 2 shows examples of two different real spectra. On the horizontal axis, we have the wavelength of light (as was also the case in the blackbody spectrum); in general, spectra can cover wavelengths of light spanning the entire electromagnetic spectrum. On the vertical axis, we have *flux*, another measure for how much light is being emitted and detected (in particular, flux measures the amount of light passing through a given area in a given amount of time).


12. [1 point] The color bar below Figure 2a (the stellar spectrum) is an alternative representation of the spectrum. How do the features in the Flux vs. Wavelength plot correspond to the features on the color bar?

Answer: Each dip/absorption line in the flux curve is a dark line at the same spot on the color bar.


Points: 1 pt for connecting the dips in flux to the dark lines in the color bar.

13. [3 points] What differences do you notice between the stellar spectrum and the galaxy spectrum? What are some similarities? Hypothesize as to *why* the spectrum of a star differs from that of a galaxy.

Answer: Both show a continuum with narrow features on it. The star's is smoother, with clean absorption dips; the galaxy's is noisier and has emission lines as well. Why: a star is one



(a) The spectrum of a star.



(b) The spectrum of a galaxy.

Figure 2

object at one temperature; a galaxy is MANY stars plus gas plus dust, all blended together (makes the spectra complicated).

Points: 1 pt for a difference, 1 pt for a similarity, 1 pt for a sensible hypothesis.

Now, let's look at the spectrum of our Sun in Figure 3.

14. [1 point] How is the Sun's spectrum different from the lamp spectra and the light bulb spectrum?

Answer: It's a full rainbow like the bulb, but with dark lines cut out of it.

Points: 1 pt for identifying the dark lines on an otherwise continuous rainbow.

Let's put everything we've seen in this lab together! Figure 4 shows three different types of spectra and how they're formed — you have now seen all three today. Blackbodies emit across all wavelengths and thus have a **continuous spectrum** (the light bulb). A sample of gas is usually a conglomerate of multiple different chemical elements, so when the atoms of a gas are heated, they emit light at certain wavelengths, thus forming an **emission spectrum** (the lamps). By contrast, if light from a blackbody or a continuous spectrum passes *through* a gas, the atoms of the gas




Figure 3: The spectrum of the Sun. Note that the actual spectrum should be one long, horizontal color gradient (like the ones seen through your diffraction gratings) – this spectrum has been broken up into rows to fit onto the page and to show more detail.

absorb some light, yielding an **absorption spectrum** (the Sun). Due to the quantum nature of atoms, each element and molecule absorbs or emits at a unique set of wavelengths; each element has its own spectral “fingerprint” that can be used to uniquely identify the chemical makeup of an object.

15. [3 points] Which type of spectrum matches the Sun’s spectrum? The light bulb spectrum? The lamp spectra?

Answer: Sun = absorption. Light bulb = continuous. Lamps = emission.

Points: 1 pt for each type of spectrum (total 3 points).

16. [2 points] Why does the Sun have that particular type, but not one of the other two?
Hint: we observe light coming from the Sun’s photosphere, but the Sun’s corona lies between the photosphere and us...

Answer: The photosphere is hot and dense, so it makes a blackbody (basically) continuum. That light then passes through gas in the corona, which absorbs certain wavelengths on the way out, so we get a continuum with lines missing. The lamps look different because they have nothing glowing behind or in front of them.

Points: 1 pt for the photosphere producing a continuum; 1 pt for that light being absorbed on its way out through the gas above it.




Figure 4: Types of spectra.

4.1 Decomposing the Stars

The spectral fingerprints we just explored allow astronomers to probe the chemical compositions of stars, galaxies, and planets. If we see an absorption line that we know is from carbon, then we know that there must be carbon in the star. The deeper the carbon absorption line in the spectrum, the more carbon there must be.

Check Courseworks for the "Stellar Spectra Activity" document in the Lab 4 folder. This should be a PDF document with a few pages. Within this document, there are two real spectra from stars, labelled in orange, and some example spectra of seven specific elements (like the ones you just looked at with the lamps) labelled in blue. The stellar spectra has already been (roughly) lined up with each element for you. Each page has different elements on it that are lined up with the stellar spectra.

17. [2 points] Go through the activity and list which elements you can see lines for in **Star 1**, then do the same for **Star 2**.

Answer: **Star 1:** hydrogen, lithium, helium. **Star 2:** hydrogen, helium.

For simplicity, students should be checking which elements have *all* their lines in the stellar spectra. (Of course, this isn't always the case in reality - but it goes to show that it takes careful work to figure out what elements a star is composed of)

Points: 1 pt for Star 1's list, 1 pt for Star 2's list.

18. [2 points] Which elements are found in *both* stars? Which elements are only found in one star?

Answer: Hydrogen and helium in both. Lithium is in Star 1 only.

Points: 1 pt for the elements in both, 1 pt for the element in only one. Grade consistently with their answer to Q17.

5 Spectra and Motion: Oneshift, Twoshift, Redshift, Blueshift

Spectra are not only useful for chemistry – we can also use spectroscopy to study the *motions* of astronomical bodies. Much like how the pitch of an ambulance changes as it moves towards or away from you, the wavelength of light changes depending on the motion of the light source – this is a manifestation of the *Doppler effect*. As a source of light moves towards you, the emitted light waves are *compressed*, yielding shorter, bluer wavelengths: this is called *blueshift*. Similarly, if a source of light is moving away from you, the light waves are *stretched*, yielding longer, redder wavelengths: this is called *redshift*. For fun, you can check out https://javablab.org/en/doppler_effect_and_redshift_en/ to see blueshift and redshift in action.




Figure 5: Redshift vs. Blueshift

This redshift or blueshift manifests itself in the spectra of moving objects, as shown in Figure 5. The middle row of Figure 5 shows the spectrum in the *rest-frame* of the object – that is, this is the spectrum we would observe if the object were perfectly stationary.

19. [3 points] Look at the **top row** of Figure 5, where the source is moving *away*. Compared to the stationary spectrum in the middle, which way did the lines move, and are those wavelengths longer or shorter? What is this called?

Answer: Moved right, toward red. Longer wavelengths. Redshift.

Points: 1 pt for the direction, 1 pt for longer wavelengths, 1 pt for naming redshift.

20. [3 points] Now look at the **bottom row**, where the source is moving *towards*. Which way did the lines move, and are those wavelengths longer or shorter? What is this called?

Answer: Moved left, toward blue. Shorter wavelengths. Blueshift.

Points: 1 pt for the direction, 1 pt for shorter wavelengths, 1 pt for naming blueshift.

21. [1 point] Because the Universe is expanding, most objects are moving *away* from us. So what should we expect to see in the spectrum of a typical galaxy, compared to its rest-frame spectrum?

Answer: Redshifted — the same pattern of lines, shifted as a group to longer wavelengths.

Points: 1 pt for redshifted.

6 Reflection

22. [1 point] Take a few minutes to reflect on your work today. What did you learn? Did anything click, or connect to something you already knew — from another class, or from your own field? Did today's lab raise any questions?

Answer: This is a free point (as long as they write *something* here).

Points: 1 pt for writing something

Total: 50 points.