

# BedSAT: Antarctica

Exploring what lies beneath using big data and modelling

Ana Fabela Hinojosa<sup>1</sup>  
March, 2025

Supervisors:  
Dr. Felicity McCormack  
Dr. Jason Roberts  
Dr. Richard Jones

Panel:  
Dr. Fabio Capitanio (Chair)  
Dr. Andrew Gunn  
Dr. Ariaan Purich



MONASH University



**SAEF**

Securing Antarctica's  
Environmental Future

Australian Research Council Special Research Initiative

---

<sup>1</sup>ana.fabelahinojosa1@monash.edu

## **Impact Statement**

Antarctica's bed topography data currently has local uncertainties of hundreds of meters in elevation due to sparse and unevenly distributed radar surveys, significantly limiting our ability to predict ice sheet behaviour and sea level rise contributions. Through the BedSAT project, I am developing a novel modelling approach that integrates remote sensing data and airborne-derived estimates, with mathematical and numerical ice flow models to substantially improve bed topography resolution and accuracy. I aim to derive a continent-wide bed topography dataset and conduct sensitivity analyses of dynamic ice loss to different realisations of topographic roughness through 2300CE. My work will quantify how bed topography uncertainties affect ice mass loss projections. This improved understanding can provide more reliable sea level rise predictions, and enable evidence-based policy decisions for climate adaptation strategies. This open-source approach and FAIR data principles will ensure these improvements benefit the broader scientific community and support more effective climate change mitigation planning.

# Contents

<b>1</b>	<b>Antarctica's Landscape</b>	<b>4</b>
1.1	Climate Impacts and Global Significance . . . . .	4
<b>2</b>	<b>Topography of Antarctica</b>	<b>5</b>
2.1	Approaches to Bed Topography Reconstruction . . . . .	6
2.2	Theoretical Frameworks . . . . .	8
2.3	Current Opportunities . . . . .	9
<b>3</b>	<b>Methods</b>	<b>10</b>
3.1	Aims . . . . .	10
3.2	Research plan methodology . . . . .	10
<b>4</b>	<b>Resources</b>	<b>13</b>
<b>5</b>	<b>Progress</b>	<b>15</b>

# Antarctica's Landscape

## 1.1 Climate Impacts and Global Significance


The polar regions are losing ice, and their oceans are changing rapidly [1]. The consequences of this extend to the whole planet and it is crucial for us to understand them to be able to evaluate the costs and benefits of potential mitigation.

Changes in different kinds of polar ice affect many connected systems. Of particular concern is the accelerating loss of continental ice sheets (glacial ice masses on land) in both Greenland and Antarctica, which has become a major contributor to global sea level rise [1]. Impacts extend beyond direct ice loss: as fresh water from melting ice sheets is added into the ocean, it increases ocean stratification disrupting global thermohaline circulation [2]. In addition, cold freshwater can dissolve larger amounts of  $\text{CO}_2$  than regular ocean water creating corrosive conditions for marine life [1].

While there is high confidence in current ice loss and retreat observations in many areas, there is more uncertainty about the mechanisms driving these changes and their future progression [3]. Uncertainty increases in regions with variable bed conditions, where characteristics like “slipperiness” and “roughness” are difficult to verify via direct observations. Other problematic areas involve the ice sheet’s grounding line (GL), the zone that delineates ice grounded on bedrock from ice shelves floating over the ocean. The retreat rate depends crucially on topographical features like pinning points [3], which lead to increased buttressing by the ice shelf on the upstream ice sheet. Although this mechanism is established, major knowledge gaps persist in mapping bed topography across Antarctic ice sheet margins - with over half of all margin areas having insufficient data within 5 km of the grounding zone [4]. Addressing this data gap through both systematic mapping and improved interpolation utilising auxiliary data streams with more complete coverage would significantly improve both our understanding of current ice dynamics and the accuracy of ice-sheet models projecting future changes.

# Topography of Antarctica

Bed topography is one of the most crucial boundary conditions that influences ice flow and loss from the Antarctic Ice Sheet (AIS) [5]. Bed topography datasets are typically generated from airborne radar surveys, which are sparse and unevenly distributed across the Antarctic continent (see figure 2.1). Interpolation schemes to “gap fill” these sparse datasets yield bed topography estimates that have high uncertainties (i.e. multiple hundreds of metres in elevation uncertainty; Morlighem et al., 2020) which propagate through simulations of AIS evolution under climate change [6]. Given the logistical challenges of accessing large parts of the Antarctic continent, there is a crucial need for alternative approaches that integrate diverse and possibly more spatially complete data streams – including satellite data.



**Figure 2.1:** Distribution of BedMAP{1,2,3} data tracks (Source: bedmap.scar.org).

## 2.1 Approaches to Bed Topography Reconstruction

A key objective of this study is to understand the bed topography itself and how it influences ice dynamics. There are two ways that we can infer information about this relationship: Through forward modelling, with assumptions of the bed conditions; and through inverse modelling that relies on surface observations.

- **Forward models**

The aim of forward models is to see how bed properties impact ice dynamics. A key example is using a large ensemble of bed topographies to investigate how bed uncertainties impact simulated ice mass loss. In this example geostatistical methods can be used to generate bed topographies that either preserve elevation or texture:

- **Geostatistics** Statistical methods specialized for analyzing spatially correlated data. In glaciology, this approach is used to interpolate between sparse measurements and characterise spatial patterns in bed properties, often employing techniques like kriging [7].

- **Inversion models**

The aim of these models is to understand bed properties through knowledge of surface or other variables. A key example is the retrieval of bed topography or basal slipperiness from surface elevation and velocities.

- **Control method inversion:** A variational approach that minimizes mismatches between observed and simulated fields through a cost function approach. Remote sensing data and theoretical ice flow models are used to obtain basal conditions [8]. Often needs regularization terms to prevent non-physical features or over-fitting [9].
- **4dvar:** Four-dimensional variational data assimilation - Similar to the control method inversion algorithm, but adds a time dimension. Used to optimize model parameters and initial conditions [9]. Can handle time-varying data and evolving glacier states, making it more suitable for dynamic systems unlike control methods, this makes them more computationally demanding [9].
- **Mass conservation:** Used to constrain inversion models and fill data gaps by employing physical conservation laws, particularly effective for reconstructing bed topography where direct measurements are sparse [5, 10]. Requires (contemporary) measurements of ice thickness at the inflow boundary to properly constrain the system [9].
- **Markov Chain Monte Carlo (MCMC):** A probabilistic method that generates sample distributions to quantify uncertainties in ice sheet parameters and models [9]. While powerful for uncertainty quantification, these methods remain computationally intensive for continental-scale ice sheet models [9].
- **EnKF** Ensemble Kalman Filter. A sequential data assimilation method that uses an ensemble of model states to estimate uncertainty and update model parameters based on observations [9].

This study aims to develop an integrated method combining forward and inverse modeling to improve bed topography estimates by leveraging high-resolution satellite surface data in

regions where radar data is sparse. Despite revolutionary advances in satellite technology that provide unprecedented surface detail, a key challenge in glaciology remains: how to fully utilize this wealth of information in regions where our understanding of subglacial conditions is limited. Our approach will integrate more comprehensive dynamical ice models with modern computational capabilities to develop better bed topography.

## 2.2 Theoretical Frameworks

Understanding how bed features manifest in surface observations requires a theoretical framework that connects these two domains. The modelling approach used in this project relies on two different theoretical frameworks that relate bed topography and surface features. Using observations and these modelling frameworks, my goal is understanding the limitations of each approach and how they can be improved.

The first framework was originally developed by Budd [11]. This model relates ice flow over bedrock perturbations to surface expressions using a two-dimensional biharmonic stress equation. The model’s foundation rests on two key simplifications:

- Most shear deformation occurs at the base of the ice sheet
- Explicit consideration of longitudinal stresses and strain-rates

The modelling carried out in [11] determined ice-sliding velocities for wide ranges of roughness, normal stress, and shear stress relevant to real glaciers [11]. Despite its robustness, Budd’s mathematical framework remains notably underutilized in modern ice sheet modeling.

The second framework in my plan analyses how shape and mechanical properties of the ice bed significantly influence how ice flows, with changes at the bed potentially leading to large differences in predicted ice loss rates [12]. Recent work by Ockenden et al. demonstrates both the capabilities and limitations of current inversion approaches in addressing this problem. The core principle of the method by Ockenden et al. (2022) relies on the fact that variations in basal topography, slipperiness, and roughness cause measurable disturbances to the surface flow of the ice. Through linear perturbation analysis, they establish a systematic relationship between surface observations and bed conditions. This relationship can be expressed as  $y = f(x)$ , where  $y$  represents surface measurements (velocity and topography),  $x$  represents bed properties (topography and slipperiness), and  $f$  is the forward model [13]. The inversion process,  $x = f^{-1}(y)$ , estimates bed conditions from surface observations.

The method works best when analyzing perturbations that are small relative to mean properties, under specific conditions including:

1. A linear viscous medium ( $n = 1$ )
2. Non-linear sliding law ( $m > 0$ )
3. Steady-state conditions
4. Spatially constant zero-order solutions

Using high-resolution datasets (REMA surface elevation at 8m resolution and NASA ITS\_LIVE velocity at 120m resolution), their approach performs well for:

- Areas with moderate topographic gradients
- Features not aligned with ice flow direction
- Medium-wavelength (5-50km) bedrock features



Nevertheless, significant limitations emerge when:

- Dealing with steep topography where the shallow-ice-stream approximation breaks down
- Handling variable slipperiness parameters
- Attempting to validate slipperiness predictions due to lack of ground-truth data

## 2.3 Current Opportunities

Current Antarctic bed topography reconstruction methods fail to utilize the wealth of presently available satellite-derived surface data. While mathematical models linking bed to surface through ice dynamics (such as those by Ockenden and Budd) provide a foundation for inferring bed topography from satellite data, they have significant limitations. My approach with BedSAT builds upon Budd’s theoretical foundations and recent inversion methods to better understand how bed conditions—including slipperiness, roughness, and pinning points—affect both grounding line retreat rates and their surface expressions. BedSAT will connect surface observations with bed topography using more realistic rheological and geometric assumptions through an iterative process: initially inverted bed topography will feed into ice dynamics models with these improved assumptions, allowing comparison between model predictions and established datasets like NASA’s ITS\_LIVE. I expect to utilise Machine learning methods to systematize this process, enhancing the analytical capabilities for the project’s final phase.

# Methods

## 3.1 Aims

I will address the following research questions:

1. How does the bed topography manifest on the ice surface?
2. To what extent do interpolation uncertainties in bed topography datasets affect the accuracy of Antarctic Ice Sheet evolution simulations under different climate change scenarios?
3. What is the impact of variable bed conditions and topography influence the rate of GL retreat in continental ice sheets?

Underpinning these research questions are the following objectives:

1. Develop an ice sheet modelling approach to assimilate satellite remote sensing datasets to improve knowledge of the bed (BedSAT) informed by mathematical models of ice flow over topography;
2. Derive a new bed topography for Antarctica using BedSAT;
3. Evaluate the impact of the improved bed topography on projections of ice mass loss from Antarctica under climate warming through sensitivity analyses.

## 3.2 Research plan methodology

Objective 1 is to derive the BedSAT method. I will use a regional catchment in Antarctica for which relatively more radar data are available and has an indicative range of topography features, e.g. the Aurora Subglacial Basin, East Antarctica, extensively surveyed by Collaborative Exploration of the Cryosphere through Airborne Profiling (ICECAP) [14]. In developing my inversion approach, I will consider important factors like sliding variability, to distinguish between topographical signatures and sliding-induced features in surface expressions. I will implement a realistic sliding-law in the Ice-sheet and Sea-level System Model (ISSM) that incorporates Budd's determinations of ice-sliding velocities across varying roughness, normal stress, and shear stress conditions. By calibrating these advanced models against known radar-surveyed regions with varying sliding conditions and validating with other observational data, I can better isolate true bed topographical features from friction-related artifacts and create a more robust inversion framework.

The second phase of the project (objective 2) will apply the methodology developed in objective 1 to the whole Antarctic continent, deriving a continent-wide bed topography dataset. Using covariance properties from existing radar surveys, I will generate a number of realisations of bed topography with unique high-resolution, and statistically-consistent topographic roughness. In the third phase of the project will use the new bed topography datasets to conduct a sensitivity analysis of ice sheet model projections to 2300 CE, investigating the impact of the new topography and different realisations of roughness on ice and subglacial hydrological flow and ice mass loss from Antarctica.

## Objective 1

1. Develop an ice sheet modelling approach to assimilate satellite remote sensing datasets to improve knowledge of the bed (BedSAT) informed by mathematical models of ice flow over topography;

## Key Investigation Areas for Objective 1

### 1. Model Development Strategy

- I will maintain invariant (but spatially variable) bed traction throughout the modeling timeframe to isolate topographical effects in the inversion approach.
- My model will incorporate available thermal distribution, velocity field, and ice thickness data, as these parameters be constrained using observations.
- Through spectral analysis of surface expressions, I will identify the topographical features that most strongly influence surface patterns.
- To account for variations in ice behavior with thickness, I will simulate scenarios ranging from thick ice with slippery base to thin ice with sticky base, using observed velocity patterns as constraints.

### 2. Transfer Functions

- I will develop efficient transfer function methods for rapid bed topography inversion, validating against known bed configurations from radar data.
- My transfer functions will be tested across various ice thickness and flow conditions.
- Cross-validation against radargrams will provide direct verification of my inversion results.
- Spatial covariance analysis of existing radar data will inform my statistical framework and error propagation through the inversion process.
- I will account for friction roughness and high-amplitude variations, using observed surface velocity patterns as constraints.

### 3. Model Validation

- I will apply quantitative error reduction metrics and compare systematically against existing bed topography products.
- Model limitations and breaking points will be identified through systematic testing across extreme scenarios, constrained by physical principles.

- Grid independence testing will ensure solution robustness across different spatial resolutions.
- Sensitivity analysis will examine the impact of my model assumptions.

*Note: Detailed methodological outlines for Objectives 2 and 3 will be developed following the completion and refinement of the BedSAT method in Objective 1.*

# Resources

I require high performance computing resources (including compute and storage) from the National Computing Infrastructure (NCI). These resources are already available via a Flagship between NCI and the Monash-led Australian Research Council project Securing Antarctica’s Environmental Future (SAEF).

## Currently available data and Framework

The project will make use of a number of new remote sensing datasets and software tools.

1. **Reference Elevation Model of Antarctica (REMA)**

Constructed from hundreds of thousands of Digital Elevation Models (DEMs) derived from high-resolution Maxar satellite imagery, REMA is calibrated with Cryosat-2 and ICESat altimetry [15].

2. **ITS\_LIVE Antarctic surface velocities and elevation**

The NASA-administered ITS\_LIVE website provides automated, high-resolution datasets of Antarctic surface velocities and ice surface elevation change, derived from satellite observations [16].

3. **BedMAP**

A suite of gridded products describing surface elevation, ice-thickness and the seafloor and subglacial bed elevation of Antarctica, based on a compilation of data collected by a large number of researchers using a variety of techniques, with the aim of representing a snap-shot of understanding of the Antarctic region [17]. BEDMAP lacks detailed information on bedrock type, sediment layers, or geothermal heat flux, all of which affect ice dynamics intruding model uncertainty.

4. **BedMachine Antarctica**

A high-resolution map of Antarctic subglacial bed topography that provides unprecedented detail of basal features. The dataset combines multiple ice thickness measurements with mass conservation principles, satellite-derived ice flow velocities, and surface mass balance from regional atmospheric models [5]. Similarly to BedMAP, BedMachine does not explicitly model basal properties.

5. **Collaborative Exploration of the Cryosphere through Airborne Profiling (ICECAP)**

Since 2012, the project has obtained extensive data on ice thickness mapping and surface elevation in regions of the East Antarctic grounding zone, also comprehensive gravity mapping in areas beneath the Totten Glacier cavity [18].

#### 6. Ice-sheet and Sea-level System Model (ISSM)

ISSM is a finite-element numerical ice sheet model. It has been used to simulate the Antarctic Ice Sheet's response to various climate scenarios and assess future mass loss contributions to sea level rise [5, 8, 19].

## Data management and archiving

Data will be published adhering to FAIR principles (Findable, Accessible, Interoperable, Reusable), ensuring transparency and accessibility. The final bed topography datasets will be published at the Australian Antarctic Data Centre (AADC) under an open source licence. All production model outputs will be published with unique DOIs at repositories aligned with the corresponding journal articles. Model outputs will be archived to tape at NCI using existing SAEF resources, as well as backed up to storage available through Monash MASSIVE M3 account aligned with project supervisor Dr McCormack. All journal articles published through this project will be open source, and tier 1 journals will be targeted.

## Risk

The project is highly feasible and low risk, given that it is a desk-based modelling and data assimilation project. All the data to be used in this project are freely available for download, and project supervisors are experts in ice sheet modelling using ISSM.

## Fieldwork


Fieldwork is not necessary to achieve the objectives of the project; however, there may be the opportunity to participate in fieldwork through the ICECAP airborne geophysics project (led CI of ICECAP is project supervisor Dr Jason Roberts, Australian Antarctic Division), which will be instrumental in training of geophysical instruments and in developing broader expertise in the field.

## Career Development


I have secured a position in the highly competitive Karthaus Glaciology Summer School taking place in northern Italy in May 2025. This prestigious program will provide comprehensive training across all aspects of glaciology directly relevant to my research, offering not only technical knowledge but also valuable networking opportunities with leading researchers in the field.

# Progress


Budd’s sliding theory examines how stress propagates through an ice mass as it flows over an undulating bed. When ice encounters a bedrock bump, it creates a complex stress field that propagates upward through the ice column at an angle rather than vertically, (see figures 5.2 for the pressure and 5.3 horizontal velocity fields). The steady-state surface shape consists of waves similar to the bedrock perturbations but out of phase by approximately  $\pi$ . As a direct mathematical consequence, “the steepest slope occurs over the highest bedrock” [11] not the surface undulation itself. To verify Budd’s sliding law, this simulation implements a flowband model across a 10 km domain with 100 m resolution. The simulation features an 800 m mean ice thickness over bedrock at 1 km elevation, with a -0.1 radians downward slope in the x-direction and imposed cosine undulations (2.64 km wavelength, 0.1 km amplitude). The main model, executes a 300-year transient Full-Stokes simulation using 1-year time steps. The current implementation demonstrates the stress propagation through ice flowing over spatially varying basal friction according to a simplified version of Budd’s sliding law (see figure 5.1) and validates the non-linear rheology ( $n = 4.0$ ) that I use.



**Figure 5.1:** Slope parallel visualisation of the computational mesh with basal nodes highlighted. The gray dotted lines represent the complete finite element mesh with multiple vertical layers conforming to the undulating geometry. Basal nodes (yellow to purple) follow the periodic undulated bed topography. The colour gradient along the basal nodes display the imposed variations in basal friction coefficient implemented through a simplified version of Budd’s sliding law, with lighter colours indicating regions of higher basal drag as per Budd’s sliding theory.




**Figure 5.2:** Pressure field distribution at  $t = 300.00$  years shown in original coordinates. The color scale (ranging from  $-2000$  to  $8000$  Pa) reveals the spatially distributed pressure variations throughout the ice body, with higher pressures (yellow) concentrated before the peaks of the bed undulations. The triangular mesh elements display the finite element discretisation used for solving the Stokes equations. The development of low-pressure zones near the surface aligning with the undulating basal topography suggests stress transfer from the imposed variation in basal friction. Note: The white gaps visible within the mesh represent masked areas excluded from visualization processing, not physical features or data gaps in the original simulation.




**Figure 5.3:** Horizontal velocity field ( $V_x$ ) at  $t = 300.00$  years displayed in original coordinates. The color scale indicates velocity magnitude ( $10^{-12}$  km/s, equivalent to  $\approx 31.5$  mm/year at the maximum) with flow direction from left to right. The velocity pattern shows clear acceleration as the ice flows downslope, with highest velocities (yellow) occurring near the terminus. The fixed upstream boundary condition constrains flow at the inlet, while the progressive acceleration downstream results from gravitational forcing along the sloped bed. Note: The white gaps visible within the mesh represent masked areas excluded from visualization processing, not physical features or data gaps in the original simulation.






**Figure 5.4:** Cross-correlation between the base and surface slope signals across spatial lags. The maximum correlation corresponds to the lag distance of 0.7 km, which translates to a phase shift of  $0.53\pi$  (95.5 degrees), only  $0.03\pi$  radians away from Budd's theoretical value.



**Figure 5.5:** Visual comparison of topographic profile of bedrock compared to the surface slope signal in slope-parallel coordinates. Here it is clearly visible that the surface slope (orange) is shifted to the right of the peak in bed elevation (blue), with approximately the predicted phase shift. This result is strong evidence that my simple model is correctly capturing the physics described in Budd's paper. The surface slope is responding to bed undulations with the expected phase relationship - maximum surface slopes occur slightly downstream of the highest bed elevations.

# Project timeline



# Bibliography

- [1] M. Meredith, M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A. Mackintosh, J. Melbourne-Thomas, M. M. C. Muelbert, G. Ottersen, H. Pritchard, and E. A. G. Schuur. *The Ocean and Cryosphere in a changing climate*, pages 203—320. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2019). DOI: [10.1017/9781009157964.005](https://doi.org/10.1017/9781009157964.005). 4
- [2] S. S. Jacobs. *Bottom water production and its links with the thermohaline circulation*. Antarctic Science **16**, 427–437 (2004). DOI: [10.1017/S095410200400224X](https://doi.org/10.1017/S095410200400224X). 4
- [3] B. Fox-Kemper, H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S. S. Drijfhout, T. L. Edwards, N. R. Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, J.-B. Sallée, A. B. A. Slangen, and Y. Yu. *Ocean, Cryosphere and Sea Level Change*. In V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (editors), *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, chapter 9, pages 1211–1362. Cambridge University Press, Cambridge, UK and New York, NY, USA (2021). DOI: [10.1017/9781009157896.011](https://doi.org/10.1017/9781009157896.011). 4
- [4] R. A. Group. *RINGS: Collaborative international effort to map all Antarctic ice-sheet margins*. Scientific Committee on Antarctic Research (2022). DOI: <https://doi.org/10.5281/zenodo.6638327>. 4
- [5] M. Morlighem, E. Rignot, T. Binder, D. Blankenship, R. Drews, G. Eagles, O. Eisen, F. Ferraccioli, R. Forsberg, P. Fretwell, V. Goel, J. S. Greenbaum, H. Gudmundsson, J. Guo, V. Helm, C. Hofstede, I. Howat, A. Humbert, W. Jokat, N. B. Karlsson, W. L. Lee, K. Matsuoka, R. Millan, J. Mouginot, J. Paden, F. Pattyn, J. Roberts, S. Rosier, A. Ruppel, H. Seroussi, E. C. Smith, D. Steinhage, B. Sun, M. R. van den Broeke, T. D. van Ommen, M. van Wessem, and D. A. Young. *Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet*. Nature Geoscience **13**, 132 (2020). DOI: [10.1038/s41561-019-0510-8](https://doi.org/10.1038/s41561-019-0510-8). 5, 6, 13, 14
- [6] B. A. Castleman, N. J. Schlegel, L. Caron, E. Larour, and A. Khazendar. *Derivation of bedrock topography measurement requirements for the reduction of uncertainty in ice-sheet model projections of Thwaites Glacier*. The Cryosphere **16**, 761 (2022). DOI: [10.5194/tc-16-761-2022](https://doi.org/10.5194/tc-16-761-2022). 5
- [7] E. J. MacKie, D. M. S. J. Caers, M. R. Siegfried, and C. Scheidt. *Antarctic Topographic Realizations and Geostatistical Modeling Used to Map Sub-*

- glacial Lakes*. Journal of Geophysical Research: Earth Surface **125** (2020). DOI: <https://doi.org/10.1029/2019JF005420>. 6
- [8] J. D. Rydt, G. H. Gudmundsson, H. F. J. Corr, and P. Christoffersen. *Surface undulations of Antarctic ice streams tightly controlled by bedrock topography*. The Cryosphere **7**, 407 (2013). DOI: [10.5194/tc-7-407-2013](https://doi.org/10.5194/tc-7-407-2013). 6, 14
- [9] M. Morlighem and D. Goldberg. *Data Assimilation in Glaciology*, pages 93–111. Cambridge University Press (2024). DOI: [10.1017/9781009180412.007](https://doi.org/10.1017/9781009180412.007). 6
- [10] M. Morlighem, C. N. Williams, E. Rignot, L. An, J. E. Arndt, J. L. Bamber, G. Catania, N. Chauché, J. A. Dowdeswell, B. Dorschel, I. Fenty, K. Hogan, I. Howat, A. Hubbard, M. Jakobsson, T. M. Jordan, K. K. Kjeldsen, R. Millan, L. Mayer, J. Mouginot, B. P. Y. Noël, C. O’Cofaigh, S. Palmer, S. Rysgaard, H. Seroussi, M. J. Siegert, P. Slabon, F. Straneo, M. R. van den Broeke, W. Weinrebe, M. Wood, and K. B. Zinglensen. *Bedmachine v3: Complete bed topography and ocean bathymetry mapping of greenland from multibeam echo sounding combined with mass conservation*. Geophysical Research Letters **44**, 11,051 (2017). ARXIV: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL074954>, DOI: <https://doi.org/10.1002/2017GL074954>. 6
- [11] W. F. Budd. *Ice Flow Over Bedrock Perturbations*. Journal of Glaciology **9**, 29 (1970). DOI: [10.3189/S0022143000026770](https://doi.org/10.3189/S0022143000026770). 8, 15
- [12] H. Ockenden, R. G. Bingham, A. Curtis, and D. Goldberg. *Inverting ice surface elevation and velocity for bed topography and slipperiness beneath Thwaites Glacier*. The Cryosphere **16**, 3867 (2022). DOI: [10.5194/tc-16-3867-2022](https://doi.org/10.5194/tc-16-3867-2022). 8
- [13] G. H. Gudmundsson and M. Raymond. *On the limit to resolution and information on basal properties obtainable from surface data on ice streams*. The Cryosphere **2**, 167 (2008). DOI: [10.5194/tc-2-167-2008](https://doi.org/10.5194/tc-2-167-2008). 8
- [14] D. A. Young, A. P. Wright, J. L. Roberts, R. C. Warner, N. W. Young, J. S. Greenbaum, D. M. Schroeder, J. W. Holt, D. E. Sugden, D. D. Blankenship, and T. D. van Ommen. *A dynamic early East Antarctic Ice Sheet suggested by ice-covered fjord landscapes*. Nature **474**, 72 (2011). DOI: <https://doi.org/10.1038/nature10114>. 10
- [15] I. M. Howat, C. Porter, B. E. Smith, M.-J. Noh, and P. Morin. *The Reference Elevation Model of Antarctica*, (2019). The Cryosphere. DOI: <https://doi.org/10.5194/tc-13-665-2019>. 13
- [16] A. S. Gardner, M. A. Fahnestock, and T. A. Scambos. *ITS\_LIVE Regional Glacier and Ice Sheet Surface Velocities: Version 1*, (2019). Data archived at National Snow and Ice Data Center. DOI: <https://doi.org/10.5067/6II6VW8LLWJ7>. 13
- [17] P. Fretwell, H. D. Pritchard, D. G. Vaughan, J. L. Bamber, N. E. Barrand, R. Bell, C. Bianchi, R. G. Bingham, D. D. Blankenship, G. Casassa, G. Catania, D. Callens, H. Conway, A. J. Cook, H. F. J. Corr, D. Damaske, V. Damm, F. Ferraccioli, R. Forsberg, S. Fujita, Y. Gim, P. Gogineni, J. A. Griggs, R. C. A. Hindmarsh, P. Holmlund, J. W. Holt, R. W. Jacobel, A. Jenkins, W. Jokat, T. Jordan, E. C. King, J. Kohler, W. Krabill, M. Riger-Kusk, K. A. Langley, G. Leitchenkov, C. Leuschen, B. P. Luyendyk, K. Matsuoka, J. Mouginot, F. O. Nitsche, Y. Nogi, O. A.

- Nost, S. V. Popov, E. Rignot, D. M. Rippin, A. Rivera, J. Roberts, N. Ross, M. J. Siegert, A. M. Smith, D. Steinhage, M. Studinger, B. Sun, B. K. Tinto, B. C. Welch, D. Wilson, D. A. Young, C. Xiangbin, and A. Zirizzotti. *Bedmap2: improved ice bed, surface and thickness datasets for Antarctica*. The Cryosphere **7**, 375 (2013). DOI: [10.5194/tc-7-375-2013](https://doi.org/10.5194/tc-7-375-2013). 13
- [18] Australian Antarctic Division. *Australian Antarctic Program: Project 4077 - Sea Ice Thickness Measurements*, (2013). 13
- [19] E. Larour, H. Seroussi, M. Morlighem, and E. Rignot. *Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model*, (2012). J. Geophys. Res., 117, F01022. DOI: [https://doi:10.1029/2011JF002140](https://doi.org/10.1029/2011JF002140). 14