

# $\mathcal{PT}$ -symmetric quantum mechanics

HONOURS LITERATURE REVIEW

Ana Fabela Hinojosa  
Student ID: 27876594

Supervisors:

Dr. Jesper Levinsen  
Prof. Meera Parish

SCHOOL OF PHYSICS & ASTRONOMY



MONASH University

11th November 2022

## Abstract

An underlying assumption of quantum theory is Hermiticity, which is a strict mathematical postulate that guarantees that predictions of observables satisfy specific physical criteria. All natural physical systems are in contact with an environment and in many cases this describes crucial features of said systems. Generally non-Hermitian quantum mechanics can describe systems which incorporate gain and loss from their environment and so physicists have developed an alternative postulate to Hermiticity known as  $\mathcal{PT}$ -symmetry. This postulate has shown to be consistent with all other aspects of the conventional theory. However a non-Hermitian  $\mathcal{PT}$ -symmetric quantum system has not yet been experimentally demonstrated.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b><math>\mathcal{PT}</math>-symmetry</b>	<b>4</b>
2.1	The $\mathcal{PT}$ operator . . . . .	4
2.2	The $\mathcal{CPT}$ inner product . . . . .	5
2.2.1	The $\mathcal{C}$ operator . . . . .	6
2.3	$2 \times 2$ $\mathcal{PT}$ -symmetric Hamiltonian . . . . .	7
<b>3</b>	<b>Broken <math>\mathcal{PT}</math>-symmetry</b>	<b>8</b>
<b>4</b>	<b>Time evolution</b>	<b>11</b>
4.1	Faster than Hermitian time evolution? . . . . .	11
<b>5</b>	<b>Experiments</b>	<b>13</b>
5.1	Observation of a Topological Transition in the Bulk of a Non-Hermitian System . . . . .	13
5.2	Light Stops at Exceptional Points . . . . .	13
5.3	Topological protection in non-Hermitian Haldane honeycomb lattices . . .	14
<b>6</b>	<b>Project statement</b>	<b>15</b>
6.1	Quantum simulation: $\mathcal{PT}$ -symmetric time evolution . . . . .	15
<b>7</b>	<b>Conclusion</b>	<b>16</b>
<b>8</b>	<b>Appendix</b>	<b>17</b>
8.1	Bi-orthogonality and completeness . . . . .	17
8.2	Equivalent Hamiltonians with distinct symmetries . . . . .	18
8.3	Equivalent $\mathcal{PT}$ -symmetric and Hermitian quantum properties . . . . .	19
8.4	Stokes wedges and boundary conditions . . . . .	19

# Introduction

In “The principles of quantum mechanics”, Paul Dirac advises us that “it is important to remember that science is concerned only with *observable* things and that we can observe an object only by letting it interact with some outside influence”[1]. Observable interactions are measurable phenomena and overtime physicists have attempted to explain a number of experimental results with a set of ad-hoc rules that came to be known as the fundamental postulates of quantum theory. Nearly all these postulates are based in physical properties. For example, one postulate establishes that time evolution of a quantum system must be probability conserving. Another postulate requires that the energy spectrum of the system is bounded below so a lowest energy state can be measured. In quantum mechanics, the Hamiltonian operator ( $\hat{H}$ ) encapsulates the total energy of a system. A system’s energy spectrum is required to be bounded and real in order to be measurable. According to the fundamental postulates of the standard theory, if an operator  $\hat{O}$  satisfies the mathematical property known as Hermiticity, then  $\hat{O}$  will satisfy the other postulates of quantum theory.

If  $\hat{O}$  is Hermitian then it has the property that its effect on the vectors of the Hilbert space in which  $\hat{O}$  is defined is independent of the order in which  $\hat{O}$  acts on said vectors[2]

$$\langle\phi|\hat{O}|\psi\rangle=\langle\phi|\hat{O}^\dagger|\psi\rangle. \tag{1.1}$$

It may appear to the reader that there is no compelling reason to change this convenient mathematical assumption, since Hermiticity results in a “correct” theory of quantum mechanics. But despite the correctness of Hermiticity, this postulate may have lead us to an overly restrictive quantum theory. The philosophically curious physicist will perhaps be persuaded to expand their mathematical toolbox by learning that not all non-Hermitian Hamiltonians describe physically invalid systems. Here it is relevant to quote Gell-Mann’s totalitarian principle: “Anything not forbidden is compulsory”. This means that if there exists an alternative non-Hermitian quantum theory then we must use it to study this class of Hamiltonians. It is the goal of this review to demonstrate that new and physically relevant theories have been developed by considering an alternative postulate to Hermiticity. This alternative postulate is referred to as space–time reflection symmetry ( $\mathcal{PT}$  symmetry)[3].

# $\mathcal{PT}$ -symmetry

In the late nineties, Bender and Boettcher presented a  $\mathcal{PT}$ -symmetric theory of quantum mechanics. Their aim was to explain a conjecture on the reality and positiveness of the spectrum of a non-Hermitian Hamiltonian proposed by Bessis in their private correspondence[4]. Bessis' conjecture was surprising since it was widely thought that non-Hermitian Hamiltonians would not have physically valid energy spectra.  $\mathcal{PT}$ -symmetric theory has since then become established and the scientists involved have produced a multitude of theoretical works and experimental proposals demonstrating that non-Hermitian  $\mathcal{PT}$ -symmetric models can be used as an extension to conventional quantum theory.

An important question that must be answered, is whether a non Hermitian  $\mathcal{PT}$ -symmetric Hamiltonian defines a valid physical theory of quantum mechanics. By a physical theory we mean that the following conditions must be satisfied; The energy spectrum of a system described by  $\hat{H}$  must be real and bounded below. Another condition is related to the probabilistic interpretation of the norm of a state, a norm must be always positive to be a valid probability. Finally, time evolution under the theory must be unitary. This means that as a state vector evolves in time the state's probability does not leak away[3, 5]. To answer this question, in this chapter I will present the theoretical background behind  $\mathcal{PT}$ -symmetric quantum theory.

## 2.1 The $\mathcal{PT}$ operator


The  $\mathcal{PT}$  operator is the anti-linear operator composed of the linear parity operator ( $\mathcal{P}$ ), which performs spatial reflection, and the anti-linear time-reversal operator ( $\mathcal{T}$ ). These operators act on position and momentum operators in the following form

$$\begin{aligned}\mathcal{P}: \quad \hat{x} &\rightarrow -\hat{x}, & \hat{p} &\rightarrow -\hat{p}, \\ \mathcal{T}: \quad \hat{x} &\rightarrow \hat{x}, & \hat{p} &\rightarrow -\hat{p}, \quad i \rightarrow -i\end{aligned}\tag{2.1}$$

Some Hamiltonians may not be symmetric under  $\mathcal{P}$  or  $\mathcal{T}$  separately, but Hamiltonians that remain invariant under the influence of the  $\mathcal{PT}$  operator are labelled as  $\mathcal{PT}$ -symmetric. A Hamiltonian  $\hat{H}$  will possess unbroken  $\mathcal{PT}$  symmetry if the following conditions are satisfied

$$[\tilde{H}, \mathcal{PT}] = 0 \quad \text{and} \quad \mathcal{PT} |\psi_n\rangle = |\psi_n\rangle, \tag{2.2}$$

where  $|\psi_n\rangle$  are the eigenstates of  $\tilde{H}$ . The second condition in 2.2 means that the eigenstates  $|\psi_n\rangle$  are simultaneously eigenstates of the  $\mathcal{PT}$  operator. Unfortunately, the anti-linearity of the  $\mathcal{PT}$  operator is responsible for the possibility that only the first identity in 2.2 could hold, but not the second[6]. If both conditions in 2.2 hold then the  $\mathcal{PT}$ -symmetry of  $\tilde{H}$  is unbroken. This means that the spectrum of  $\tilde{H}$  is fully real and bounded below –This is also known as exact  $\mathcal{PT}$ -symmetry–.



**Figure 2.1:** Illustration of the conditions in equations 2.2.  $\mathcal{PT}$ -symmetry and the reality of eigenvalues. This figure is quite subtle. Unbroken or exact  $\mathcal{PT}$ -symmetry requires that both conditions in 2.2 are true. Since the  $\mathcal{PT}$  operator is antilinear it is possible that eigenstates of a Hamiltonian are not also eigenstates of the  $\mathcal{PT}$  operator. This figure is strongly inspired by figure 2 in reference [6]

Most model  $\mathcal{PT}$ -symmetric Hamiltonians  $\tilde{H}$  in the literature admit tunable parameters such that there will be regions in the parameter space where the eigenstates of  $\tilde{H}$  are not  $\mathcal{PT}$ -symmetric[7]. This means that a system described by  $\tilde{H}$  admits two phases, one for broken and another for unbroken  $\mathcal{PT}$ -symmetry. In the unbroken phase eigenstates of  $\tilde{H}$  become degenerate and this implies that they lose completeness[7].

The effectiveness of  $\mathcal{PT}$  symmetry as a tool to investigate the spectra of some non-Hermitian Hamiltonians has been shown in various works, such as Dorey et al[8], Bender and Boettcher[4], Brody[9], Bender and Mannheim [10], Bender et al[11], Mostafazadeh[12][13] among others. To ensure that the eigenstates  $\{|\psi_n\rangle\}$  of a  $\mathcal{PT}$ -symmetric Hamiltonian  $\tilde{H}$  are orthogonal, it is necessary to specify an inner product. A possible choice for the inner product associated with  $\tilde{H}$  might be[11]

$$\langle\psi_n|\psi_m\rangle \equiv \int_c dx [\psi_n(x)]^{\mathcal{PT}} \psi_m(x) = \int_c dx [\psi_n(-x)]^* \psi_m(x) \quad (2.3)$$

where  $c$  is a complex contour that starts and ends within specific boundary conditions (see section 8.4). Unfortunately this obvious choice for the  $\mathcal{PT}$  inner product is not correct because it does not guarantee a positive definite norm.

## 2.2 The $\mathcal{CPT}$ inner product

To be able to describe precisely the nature of  $\mathcal{PT}$ -symmetric quantum mechanics, we must delve briefly into the inner-product under which our theory satisfies the postulates of conventional quantum mechanics. It is important to note that  $\mathcal{PT}$ -symmetric quantum mechanics is a kind of ‘bootstrap’ theory[5], since infinitely many inner-products exist for a given vector space, we can construct an inner product whose associated norm is positive definite by design. This inner-product is in general dependent on the characteristics of the Hamiltonian in question and it guarantees that the underlying dynamics of any  $\mathcal{PT}$ -symmetric Hamiltonian satisfies unitarity[3]. Firstly, it is necessary to solve for the eigenstates of the Hamiltonian before knowing the Hilbert space and consequentially the associated inner product. To guarantee a positive norm for our theory, we will construct a new linear operator  $\mathcal{C}$  that commutes with both  $\tilde{H}$  and  $\mathcal{PT}$ . We use the symbol  $\mathcal{C}$  to represent this symmetry because its properties are similar to those of the charge conjugation operator in particle physics[5].

### 2.2.1 The $\mathcal{C}$ operator

When the  $\mathcal{PT}$ -symmetry of  $\hat{H}$  is unbroken, the eigenfunctions  $\psi_n(x)$  of  $\hat{H}$  are simultaneously eigenstates of the  $\mathcal{PT}$  operator[14]

$$\mathcal{PT}\psi_n(x) = \lambda_n\psi_n(x), \quad (2.4)$$

Where  $\lambda_n$  is a pure phase. Without loss of generality, for each  $n$  the phase can be absorbed into  $\psi_n(x)$  and this makes the eigenvalue of the  $\mathcal{PT}$  operator unity[14]:

$$\mathcal{PT}\psi_n(x) = \psi_n^*(-x) = \psi_n(x). \quad (2.5)$$

There is strong numerical evidence of the completeness of the eigenfunctions  $\psi_n(x)$ [15, 14][7]. In the coordinate basis, the completeness statement reads:

$$\sum_n (-1)^n \psi_n(x) \psi_n(y) = \delta(x - y), \quad x, y \in \mathbb{R} \quad (2.6)$$

the unconventional  $(-1)^n$  factor in 2.6 can be explained if we define the  $\mathcal{PT}$  inner product as

$$\begin{aligned} (f, g) &= \int dx [\mathcal{PT}f(x)] g(x) \\ &= \int dx f^*(-x) g(x). \end{aligned} \quad (2.7)$$

where the integral above follows a path in the complex plane. In contrast with a Hermitian Hamiltonian –which always commutes with a parity operator–, the definition in 2.7 describes how eigenstate norms alternate in sign depending on the value  $n$ . This means that the metric associated with the  $\mathcal{PT}$  inner product is indefinite[14, 16]. In quantum theory, the norm of states is interpreted as a probability and this means that the indefinite metric described above presents a serious problem for the validity of  $\mathcal{PT}$ -symmetric quantum theory. The solution to this problem lies in finding an interpretation for the negative valued norms[11].

The literature claims that for any theory with unbroken  $\mathcal{PT}$ -symmetry there exists a symmetry of the Hamiltonian that describes the negative and positive norm states. To describe this symmetry of  $\hat{H}$  it is necessary to construct a linear operator denoted by  $\mathcal{C}$ [3, 15, 14]. When represented in position space  $\mathcal{C}$  is a sum over the energy eigenstates of  $\hat{H}$ :

$$\mathcal{C} = \sum_n \psi_n(x) \psi_n(y) \quad (2.8)$$

From this definition, relation  $(\psi_m(y), \psi_n(y)) = \int dy (\mathcal{PT}\psi_m(y)) \psi_n(y)$  and equation 2.5 we can verify that the eigenvalues of  $\mathcal{C}$  are  $\pm 1$

$$\begin{aligned} \mathcal{C}\psi_n(x) &= \int dy \mathcal{C} \psi_n(y) \\ &= \sum_m \psi_m(x) \int dy \psi_m(y) \psi_n(y) \\ &= (-1)^n \psi_n(x) \end{aligned} \quad (2.9)$$

The  $\mathcal{PT}$  norm signatures can therefore be interpreted as the “charge” of the states, while  $\mathcal{C}$  is the operator used to measure this charge[14].

The  $\mathcal{C}$  operator commutes with the  $\mathcal{PT}$  operator, but it does not commute with the parity operator  $\mathcal{P}$ . Notice that  $\mathcal{C}$  and  $\mathcal{P}$  operators are square roots of  $\delta(x - y)$  the unity operator[15]

$$\mathcal{P}^2 = \mathcal{C}^2 = \mathbb{1}, \quad (2.10)$$

where  $\mathcal{P} \neq \mathcal{C}$ , since  $\mathcal{P}$  is real and  $\mathcal{C}$  is complex valued[3][14].

Using the newly constructed  $\mathcal{C}$  operator, we can redesign the  $\mathcal{PT}$  inner product to suit the conventional probabilistic interpretation of the vector norms in quantum mechanics

$$(f, g) = \int dx [\mathcal{CPT}f(x)] g(x). \quad (2.11)$$

We must point out that all experimental evidence to date shows that the nature follows the Dirac inner-product[17]. However, the definitions show much promise as they might allow us to study non-Hermitian dynamics in experimental set-ups.

### 2.3 $2 \times 2$ $\mathcal{PT}$ -symmetric Hamiltonian

An example of the results of unbroken  $\mathcal{PT}$ -symmetric theory that is often found in the literature describes a simple finite non-Hermitian but  $\mathcal{PT}$ -symmetric Hamiltonian

$$\tilde{H} = \begin{pmatrix} re^{i\theta} & s \\ s & re^{-i\theta} \end{pmatrix} \quad (2.12)$$

where the parameters  $r, s$  and  $\theta$  are real. In this set-up the  $\mathcal{T}$  operator performs complex conjugation and the parity operator is written as

$$\mathcal{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (2.13)$$

The Hamiltonian in 2.12 allows two parametric regions. When  $s^2 < r^2 \sin^2 \theta$  the eigenvalues of  $\tilde{H}$  form a complex conjugate pair. This is the broken  $\mathcal{PT}$ -symmetry region. Otherwise, when  $s^2 \geq r^2 \sin^2 \theta$ , then the eigenvalues are real. This is the region of unbroken  $\mathcal{PT}$ -symmetry region. In the unbroken symmetry region the  $\mathcal{PT}$  normalised eigenstates of both  $\tilde{H}$  and  $\mathcal{PT}$  are

$$|E_+\rangle = \frac{1}{\sqrt{2 \cos \alpha}} \begin{pmatrix} e^{i\alpha/2} \\ e^{-i\alpha/2} \end{pmatrix} \quad \text{and} \quad |E_-\rangle = \frac{i}{\sqrt{2 \cos \alpha}} \begin{pmatrix} e^{-i\alpha/2} \\ -e^{i\alpha/2} \end{pmatrix}, \quad (2.14)$$

we use the relation  $\sin \alpha = \frac{r}{s} \sin \theta$ . The corresponding  $\mathcal{C}$  operator for this Hamiltonian is

$$\mathcal{C} = \frac{1}{\cos \alpha} \begin{pmatrix} i \sin \alpha & 1 \\ 1 & -i \sin \alpha \end{pmatrix}. \quad (2.15)$$

This  $\mathcal{C}$  operator commutes with  $\tilde{H}$  and satisfies  $\mathcal{C}^2 = \mathbb{1}$ . The eigenvalues of  $\mathcal{C}$  are precisely the signs of the  $\mathcal{PT}$  norms of the corresponding eigenstates[14]. Using this  $\mathcal{C}$  operator we can guarantee that the  $\mathcal{CPT}$  inner product in the two-dimensional Hilbert space spanned by  $|E_{\pm}\rangle$  is positive definite.


# Broken $\mathcal{PT}$ -symmetry

In contrast with unbroken symmetry, Hamiltonians with broken  $\mathcal{PT}$ -symmetry do not satisfy the spectral theorem and this gives rise to very unconventional quantum mechanical behaviour. Hamiltonians with broken  $\mathcal{PT}$ -symmetry exhibit purely non-Hermitian spectral degeneracies known as exceptional points (EP)[18]. EPs are points in an (at least) two-dimensional parameter space, in which two or more eigenstates become identical[19][20]. That is, in the presence of EPs, the coalescence of eigenstates causes the collapse of the subspace dimension to one, because of this the eigenstate no longer forms a complete basis[21]. The interest in these degeneracies has increased greatly since they give rise to dramatic effects in a great variety of physical problems: in mechanics, electromagnetism, atomic and molecular physics, quantum phase transitions, quantum chaos and more[22]. An illustrative example of EPs can be seen when we consider the well known family of  $\mathcal{PT}$ -symmetric Hamiltonians

$$\hat{H} = \hat{p}^2 + \hat{x}^2(ix)^\epsilon \quad (\epsilon \text{ real}). \quad (3.1)$$


These Hamiltonians have unbroken  $\mathcal{PT}$  symmetry when  $\epsilon \geq 0$  and broken  $\mathcal{PT}$ -symmetry when  $\epsilon < 0$ . The energy spectrum of 3.1 has been proved to be real and bounded below in the region of unbroken  $\mathcal{PT}$  symmetry [23]. In figure 3.1 both these regions and the phase transition the spectrum are visible.





**Figure 3.1:** This figure is my version of Figure 1. in reference [5], it exemplifies the behaviour of the eigenvalue spectra of the Hamiltonians in 3.1 as a function of the real parameter  $\epsilon$ . All the eigenvalues visible in this figure are real valued. The vertical dotted green line corresponds to the value  $\epsilon = 0$ , this is the point where  $\mathcal{PT}$ -symmetry breaks and it corresponds to the Hamiltonian for the classic one dimensional harmonic oscillator. When  $\epsilon \geq 0$  the spectra are real, positive and discrete. Energy levels increase with increasing  $\epsilon$ . In the region corresponding to  $-1 \leq \epsilon < 0$  there is a finite number of real positive eigenvalues and an infinite number of complex-conjugate pairs of eigenvalues (not depicted). The number of real eigenvalues decreases as  $\epsilon$  decreases from 0 to  $-1$ , for  $\epsilon$  that are more negative than the value  $\epsilon = -0.57793$  the only remaining real eigenvalue corresponds to the ground-state energy. At the value  $\epsilon = -1$  the spectrum is null as all eigenvalues diverge towards positive infinity.

In chapter 6 we will investigate the unbroken symmetry region of figure 3.1, specifically the case for  $\epsilon = 2$ . However since in this section we are concerned with EPs, we must zoom into the parametric region  $\epsilon < 0$ . The broken  $\mathcal{PT}$ -symmetry region is visible in more detail in figure 3.2.



**Figure 3.2:** This figure is my version of Figure 3. in reference [24], Similarly to figure 3.1, this is the eigenvalue spectra of the Hamiltonians in 3.1 as a function of the real parameter  $\epsilon$  in the case where  $\epsilon < 0$ . This figure displays a discretised parameter space for  $\epsilon$  as 100 values in the horizontal axis, and the energy range  $-2 \leq E \leq 12$  as the vertical axis. This figure depicts fully real eigenvalues as black dots. Visible here are three epsilon values at which the real eigenvalues become degenerate (coalesce) and then form complex-conjugate pairs. The real parts of these pairs of eigenvalues are represented as blue dots. These energies initially decrease as  $\epsilon$  decreases but blow up suddenly as  $\epsilon$  approaches  $-1$  (The blow up of the real-parts was not captured by my numerics). The imaginary parts of the eigenvalue pairs are represented as red dots. These remain finite and appear to decay to zero at  $\epsilon = -1$ .

In figure 3.2 there are three EPs of order 1 (EP1) and a single infinite order EP at  $\epsilon = -1$ . My numerics did not have enough resolution to represent the infinite order EP.

When physicists test the character and behaviour of states and eigenvalues of EPs is common practice to encircle the EP on a closed contour in parameter space. For examples see chapter 5. There are cases of EPs where eigenvalues and the wave functions are interchanged, For example, if the EP corresponds to a cubic root branch point. For a closed contour around this point a cyclic permutation of all three eigenvalues and eigenfunctions is then expected[20].

Time evolution at EPs differs from generic Hamiltonians in the sense that only one state evolves with the usual oscillations (possibly with overall damping or amplification), while other states remain inextricably coupled to this single eigenstate[18].

# Time evolution

In Hermitian quantum mechanics, time-evolution of states is done by using the unitary operator  $\hat{U} = e^{-i\hat{H}t/\hbar}$ . Because the  $\hat{U}$  operator is unitary this means that as the state  $\psi$  evolves in time, its norm remains constant. This is useful in quantum mechanics because the norm of a state is interpreted as a probability, and therefore unitarity guarantees that a state's probability does not grow or decay with time. As explained in section 2.2, when the  $\mathcal{PT}$ -symmetry of  $\hat{H}$  is unbroken it is possible to have positive definite  $\mathcal{CPT}$  norm states for the non-Hermitian Hamiltonian. This means that time evolution under the unbroken  $\mathcal{PT}$ -symmetric framework is also unitary[2][15][13]. The unitary time evolution of a state  $\vec{\psi}$  from time  $t = 0$  to a time  $t$  is written as

$$\vec{\psi}(t) = \hat{U}\vec{\psi}(0). \quad (4.1)$$

An interesting feature of  $\mathcal{PT}$ -symmetric Hamiltonians has been suggested in [25]. According to this paper it is possible to obtain arbitrarily fast quantum evolutions using non-Hermitian  $\mathcal{PT}$ -symmetric Hamiltonian operators. This is because for such Hamiltonians the path from can be made short. A brief description of this work is described below.

## 4.1 Faster than Hermitian time evolution?

The mechanism described in [25] is similar to that in general relativity in which the distance between two space-time points can be made small if they are connected by a wormhole. If true, The results in [25] will have drastic consequences in quantum computation, because it removes the bound on the time-optimal unitary NOT operations[26] that is obtained in quantum mechanics[27]. However, Bender et al warn us about the possibility of the existence of quantum protection mechanisms that place a lower bound on the time required to switch Hilbert spaces. which would limit the applicability of a Hilbert-space “wormhole” to improve quantum algorithms[25].

Günther and Samsonov have interpreted the  $\mathcal{PT}$ -symmetric set-up for [25] as a quantum system consisting of a non-Hermitian  $\mathcal{PT}$ -symmetric component and a Hermitian component simultaneously. In [28] Günther and Samsonov formulate a general recipe for the construction of partially  $\mathcal{PT}$ -symmetric quantum systems which are not 1:1 equivalent to purely Hermitian systems. Using a strong structural analogy with the reference frames for inertial observers in special relativity, they associate  $\mathcal{PT}$ -symmetric models in different representations with corresponding measurement frames. They claim that the resultant operators which are Dirac Hermitian are connected with non-Dirac-Hermitian operators in another frame. The probabilistic content of the models is frame-independent. According to Günther and Samsonov their geometric analysis of the equivalence mapping

between mutually  $\mathcal{PT}$ -symmetric and Hermitian operators demonstrates the agreement of the vanishing passage-time solution proposed in [25] and the temporal lower bound for Hermitian evolution proposed by Anandan and Aharonov in [29].

Despite the potential importance of  $\mathcal{PT}$ -symmetric theory, the experimental observation in a genuine quantum system is still missing[20]. Therefore the purpose of our research is determine whether it is possible to observe the dynamical behaviour of a  $\mathcal{PT}$ -symmetric quantum in an experiment.

# Experiments

Much research has been carried out on properties of physical systems described by  $\mathcal{PT}$ -symmetry. Such Hamiltonians can be used to model the behaviour of closed quantum systems, but they can also be replicated in open systems which exhibit balanced gain and loss. This has been implemented in laboratory experiments for a wide range of systems[30], here are some brief examples.

## 5.1 Observation of a Topological Transition in the Bulk of a Non-Hermitian System

The work of Zeuner et al in [31] describes topological transitions in a non-Hermitian system. Topological transport is a chiral transport phenomenon of particles due to Hall conductance that can be linked to an integer-valued topological quantity. This kind of phenomenon has been observed in two-dimensional electron gases, silicon platforms, and even in ultra-cold gases[31].

In this paper Zeuner et al present an experiment where they realise the idealised one-dimensional non-Hermitian dimer model. The model used in this investigation permits the loss or absorption of transported particles at every second site of a one-dimensional optical lattice. It is this phenomenological loss of particles that makes the system non-Hermitian. The experimental methodology is based on bulk measurements of topological transitions in fabricated waveguide lattices with alternating distances in fused silica glass. To introduce loss, every second waveguide is sinusoidally oscillated transversely to the plane of the waveguides[31]. The results of this experiment leads Zeuner et al to argue that the features of the idealized tight binding model in reference [32] carries over to experiments. Because loss is an inherent feature of many systems that may exhibit topological effects (such as coupled quantum dots[32], and driven-dissipative exciton-polariton condensates [33]). This intrinsic non-Hermiticity may act as a surprising probe of their potentially rich topological phases[31].

## 5.2 Light Stops at Exceptional Points

In 2018 Goldzak et al published [34] where they relate stopped light pulses and the phenomenon of an exceptional point (EP). In their work they claim to conceptually show that the group velocity of light vanishes if a waveguide is designed exactly at the EP. Goldzak et al explain that this pulse-stop effect is possible in a  $\mathcal{PT}$ -symmetric optical waveguides where two optical modes coalesce.  $\mathcal{PT}$ -symmetry is characterized by accurately balanced loss and gain in time.

In their proposed experiment Goldzak et al “freeze” and release Gaussian light pulses by adiabatically increasing gain and loss parameters. Using a pumped optical cavity as the non-linear medium, Goldzak et al explain that parametric interactions operating at ultra fast rates can be engineered using synchronously pumped optical parametric oscillators [34]. After the light pulse is “released” it should be possible to observe that the carried coherent information is preserved. They explain that it is the non-resonant mechanism of this phenomenon that is of an important technological advantage, since apparently this result can be tuned for optical pulses in a wide range of frequencies and bandwidths. In addition they propose that it should be possible to engineer this effect in any  $\mathcal{PT}$ -symmetric system of two waveguide channels. which in theory should not be limited only to light but could be extended to acoustic waves or other fields in physics satisfying  $\mathcal{PT}$  symmetry [34]. If replicable this result would open new possibilities for designing slow light devices, which would exploit generic properties of EPs offering interesting operational capabilities of a new generation of photonic devices [34].

### 5.3 Topological protection in non-Hermitian Haldane honeycomb lattices

The work by Reséndiz-Vázquez et al focuses in the possibility of observing topologically protected edge states in two-dimensional non-Hermitian Haldane lattices [35]. The Haldane model represents a unique system where the quantum Hall effect is contained as an intrinsic lattice-band-structure property, rather than an external effect due to the presence of a strong magnetic field [36].

The claim in this paper is that edge states can be observed in the so-called broken  $\mathcal{PT}$ -symmetric phase. The experiment is done using finite rectangular Haldane ribbon comprising 1860 sites. They present the results for a lattice several kinds of terminations. Reséndiz-Vázquez et al find that such topologically protected edge states emerge irrespective of the lattice boundaries, namely, zigzag, bearded, or armchair. which implies that, in general, the edge geometry of the sample does not play a role in the observation of topological edge protection as long as the system is described by a ribbon. This contrasts with previous findings. The results reported by Reséndiz-Vázquez et al help elucidate the role of  $\mathcal{PT}$  symmetry in two-dimensional topological phenomena and demonstrate that topological protection can exist in the archetypal Haldane model even in the presence of gain and loss [35].

There is an ongoing quest devoted to investigating the advantages of extending quantum theory via  $\mathcal{PT}$ -symmetry. On occasion this has revealed new phenomena apparently exclusive to non-Hermitian systems, to read more on this please refer to [37] and [38].

# Project statement

We are interested in understanding the dynamics of quantum mechanical super fluids in non-Hermitian traps. Our ultimate goal in this project is the experimental observation of the dynamics of a Bose-Einstein condensate in a three-dimensional Hermitian potential as a quench transforms the potential into a  $\mathcal{PT}$ -symmetric trap in the shape of an inverted quartic in one dimension, with a potential extension into a non-Hermitian  $\mathcal{PT}$ -symmetric trap in three-dimensions. In section 3 we described the well known parametric family of  $\mathcal{PT}$ -symmetric Hamiltonians in 3.1. As explained previously these Hamiltonians have unbroken  $\mathcal{PT}$  symmetry when  $\epsilon \geq 0$  and broken  $\mathcal{PT}$ -symmetry when  $\epsilon < 0$ . The particular Hamiltonian that we use in my project occurs when we let  $\epsilon = 2$  in 3.1. Our Hamiltonian is

$$\hat{H} = \hat{p}^2 - \hat{x}^4; \tag{6.1}$$

this Hamiltonian is interesting because even though it possesses an upside-down potential, it nevertheless appears to have a fully real and bounded below energy spectrum[39] as can be seen in figure 3.1. Our present aim is to simulate the dynamics of a single trapped state after we implement a quench that would instantaneously change the trapping potential from a harmonic oscillator trap to an inverted quartic potential exactly like the one in the Hamiltonian in 6.1.

## 6.1 Quantum simulation: $\mathcal{PT}$ -symmetric time evolution

Here I include some of the progress I made in my work on the dynamics under the  $\mathcal{PT}$ -symmetric framework.

For  $\tilde{H}$  whose eigenvectors span the Hilbert space  $\mathcal{H}$ [25].

# Conclusion



# Appendix

## 8.1 Bi-orthogonality and completeness

In the case of a  $\mathcal{PT}$ -symmetric Hamiltonian  $\tilde{H}$ , it is possible for two (or more) complex eigenvalues to cross. At the crossing point the eigenvalues are degenerate and this is accompanied by the coalescence of the eigenstates of  $\tilde{H}$ . If eigenstates of  $\tilde{H}$  become degenerate this implies that they lose completeness[7]. This special situation is associated with a branch point in the complex energy plane and is commonly termed an “exceptional point”[19]. Moreover, sufficiently close to the branch point the eigenfunctions of the coalescing eigenvalues are nearly self-orthogonal.

We need completeness and closure relations in order to develop, for example, perturbation theory and scattering theories for non-Hermitian Hamiltonians and in order to be able to solve the Schrödinger equation by numerical methods[19].

Suppose that a diagonalizable non-Hermitian Hamiltonian  $\tilde{H}$  has a discrete spectrum and it commutes with the  $\mathcal{PT}$  operator. This means that  $\tilde{H}$  has unbroken  $\mathcal{PT}$ -symmetry. A Hilbert space  $\mathcal{H}$  spanned by the eigenvectors of  $\tilde{H}$  exists, but because  $\tilde{H}$  is non-Hermitian –with respect to the standard inner product of  $\mathcal{H}$ – the time evolution generated by  $\tilde{H}$  will not be unitary[12]. Nevertheless, as explained in section 2.2.1. For the assumed  $\tilde{H}$  above it is possible to construct a different complete positive-definite inner product by constructing and using the antisymmetric  $\mathcal{C}$  operator that corresponds to  $\tilde{H}$ .

The assumed diagonalizability condition of  $\tilde{H}$  may be viewed as a physical requirement without which an energy eigenbasis would not exist. To our knowledge all known non-Hermitian Hamiltonians that are used in physical applications are diagonalizable and therefore admit a complete *bi-orthonormal* set of eigenvectors. This set of bi-orthonormal eigenvectors is  $\{(|\psi_n, a\rangle, |\phi_n, a\rangle)\}$  and will satisfy the following defining relations[40]

$$\tilde{H} |\psi_n, a\rangle = E_n |\psi_n, a\rangle, \quad \tilde{H}^\dagger |\phi_n, a\rangle = E_n^* |\phi_n, a\rangle, \quad (8.1)$$

$$\langle \phi_m, b | \psi_n, a \rangle = \delta_{mn} \delta_{ab}, \quad (8.2)$$

$$\sum_n \sum_{a=1}^{d_n} |\psi_n, a\rangle \langle \phi_n, a| = \hat{\mathbb{I}} \quad (8.3)$$

$$\tilde{H} = \sum_n \sum_{a=1}^{d_n} E_n |\psi_n, a\rangle \langle \phi_n, a|, \quad \tilde{H}^\dagger = \sum_n \sum_{a=1}^{d_n} E_n^* |\phi_n, a\rangle \langle \psi_n, a| \quad (8.4)$$

where  $n$  and  $a$  are the spectral degeneracy levels,  $d_n$  is the degree of degeneracy of  $E_n$  and  $\hat{\mathbb{I}}$  is the identity operator[40].

## 8.2 Equivalent Hamiltonians with distinct symmetries

In parallel to the work of Bender is the research of Mostafazadeh who introduced the notion of pseudo-Hermiticity in [12]. A Hamiltonian is said to be pseudo-Hermitian with respect to a positive-definite, Hermitian operator  $\eta$  if it satisfies

$$\tilde{H}^\dagger = \eta^{-1} \tilde{H} \eta. \quad (8.5)$$

In the case of Hamiltonians with  $\mathcal{PT}$ -symmetry, the role of  $\eta$  is played by the combined action of  $\mathcal{P}$  and  $\mathcal{C}$ . A convenient way to write the  $\mathcal{C}$  operator was proposed in [41]

$$\mathcal{C} = e^Q \mathcal{P} = \eta^{-1} \mathcal{P}, \quad (8.6)$$

here  $Q$  is an antisymmetric Hermitian operator. Mostafazadeh [13] has shown that the square root of the positive-definite Hermitian operator  $\eta = e^{-Q}$  can be used to transform any non-Hermitian Hamiltonian with unbroken  $\mathcal{PT}$ -symmetry into a spectrally equivalent Hermitian Hamiltonian by means of a unitary “similarity transformation” [42][43]. The transformation is as follows, the invertible operator  $\rho = \sqrt{\eta}$  acts on the non-Hermitian  $\mathcal{PT}$ -symmetric Hamiltonian  $\tilde{H}$  and returns an equivalent Hermitian Hamiltonian  $\hat{H}$

$$\hat{H} = \rho^{-1} \tilde{H} \rho = e^{-Q/2} \tilde{H} e^{Q/2}. \quad (8.7)$$

To verify that the similar Hamiltonian  $\hat{H}$  is Hermitian we take the Hermitian conjugate of  $\hat{H}$

$$\begin{aligned} \hat{H}^\dagger &= (e^{-Q/2} \tilde{H} e^{Q/2})^\dagger, \\ &= e^{Q/2} \tilde{H}^\dagger e^{-Q/2}, \end{aligned} \quad (8.8)$$

If we “swap” Hermiticity:  $\tilde{H}^\dagger$  in 8.8 for  $\mathcal{PT}$ -symmetry, and we use equation 8.6

$$\begin{aligned} \hat{H}^\dagger &= e^{Q/2} \mathcal{P} \tilde{H} \mathcal{P}^{-1} e^{-Q/2}, \\ &= e^{-Q/2} \mathcal{C} \tilde{H} \mathcal{C}^{-1} e^{Q/2}, \end{aligned} \quad (8.9)$$

Finally we recall that  $\mathcal{C}$  and  $\tilde{H}$  commute

$$\hat{H}^\dagger = e^{-Q/2} \tilde{H} e^{Q/2} = \hat{H}. \quad (8.10)$$

Mostafazadeh [12], conjectures that because  $\mathcal{CPT}$ -symmetry satisfies the postulates of quantum mechanics –whilst only “swapping” the Hermiticity of a Hamiltonian by  $\mathcal{CPT}$ -symmetry– then this must mean that non Hermitian  $\mathcal{CPT}$ -symmetric theories are equivalent to certain non local Hermitian field theories. This kind of transformation is explored in references [12],[44],[40],[43],[45]. In general this transformation is extremely complicated and so it is quite impractical to perform. Hence, while the mapping from  $\mathcal{PT}$ -symmetric to Hermitian Hamiltonians is of great theoretical interest, it does not have much practical value[5]. Another important fact is that this kind of transformation is a similarity and not a unitary transformation. Thus, while the eigenvalues of both Hamiltonians are identical, the relationships between vectors are changed; pairs of vectors that are orthogonal are mapped into pairs of vectors that are not orthogonal[5][25].


### 8.3 Equivalent $\mathcal{PT}$ -symmetric and Hermitian quantum properties

In the case of unbroken  $\mathcal{PT}$ -symmetry of a Hamiltonian  $\tilde{H}$  with eigenfunctions  $\psi_n$  then the  $\mathcal{PT}$  operator eigenfunctions are:  $\chi_n(x) = e^{-i\alpha/2}\psi_n(x)$

Properties	Hermitian	$\mathcal{PT}$ -symmetry (unbroken)
Eigenvalues and Eigenstates	$\hat{H}  \psi_n\rangle = E_n  \psi_n\rangle$	$\tilde{H}  \psi_n\rangle = E_n  \psi_n\rangle,$ $\langle\phi_n  \tilde{H}^\dagger = \langle\phi_n  E_n^*$
Inner product (ignoring degenerate spectra)	$\langle\psi \phi\rangle = ( \psi\rangle)^\dagger \cdot  \phi\rangle$	$(\psi \phi) = (\mathcal{CPT}  \psi\rangle)^T \cdot  \phi\rangle$
Orthonormality of states	$\langle\psi_n \psi_m\rangle = \delta_{mn}$	$(\psi_n \psi_m) = \delta_{mn}$
Completeness of states	$\sum_{n=0}^{\infty} [\psi(x)]^* \psi(y) = \delta(x - y)$	$\sum_{n=0}^{\infty} \psi_n(x) [\mathcal{CPT} \psi_n(y)] = \delta(x - y)$
Hamiltonian, Greens and zeta functions	$H(x, y) = \sum_{n=0}^{\infty} E_n [\psi_n(x)]^* \psi_n(y),$ $G(x, y) = \sum_{n=0}^{\infty} \frac{1}{E_n} [\psi_n(x)]^* \psi_n(y),$ $\int dx G(x, x) = \sum_{n=0}^{\infty} \frac{1}{E_n}$	$H(x, y) = \sum_{n=0}^{\infty} (-1)^n E_n \chi_n(x) \chi_n(y),$ $G(x, y) = \sum_{n=0}^{\infty} \frac{(-1)^n}{E_n} \chi_n(x) \chi_n(y),$
Time evolution	$\hat{U} = e^{-i\hat{H}t/\hbar}$	$\hat{U} = e^{-i\tilde{H}t/\hbar}$

### 8.4 Stokes wedges and boundary conditions

The boundary conditions for the eigenvalue problem corresponding to the Hamiltonians in equation 3.1 depend on the deformation parameter  $\epsilon$ [5]. Since the eigenvalue problem can be conceptualised as a harmonic oscillator that has been deformed into the complex x-plane[46]. The integration contour we choose to use should be located within two angular sectors in the complex x-plane. These sectors are known as *Stokes wedges*[5]. In general these wedges are centred about the origin in the complex x-plane and we can define their orientation in terms of  $\text{Arg}(x) = \theta$ . The solutions to the differential equation are analytic functions of  $x$ [47], therefore the eigenvalues are independent of the choice of contour (see figure 8.1) in the complex x-plane[11][48]. However, it is crucial that the end points of the contour are located within the stokes wedges, and that solutions to the differential equation must be equal to zero at both ends of the contour in the stokes wedges[6].



**Figure 8.1:** Example of Stokes wedges in the complex plane. The configuration of the wedges varies depending on the problem at hand. Infinitely many possible integration contours are allowed to be used in order to solve the Schrödinger equation for the Hamiltonians in equation 3.1. But it is required that the endpoints of the integration contour must remain within the stokes wedges. The integration contour must be continuous but it need not be smooth and may even cross itself[11].

Stokes wedges must be non-adjacent in order to describe the boundary conditions of a valid coordinate-space eigenvalue problem. If the wedges are adjacent they would not represent a continuous region within the same Riemann sheet. Determining the locations of Stokes wedges does not require that we solve the differential equation exactly but it can be done using asymptotic approximations[11].

# Bibliography

- [1] P. A. M. Dirac. *THE PRINCIPLES OF QUANTUM MECHANICS*. International series of monographs on physics (Oxford, England). Clarendon Pr., Oxf., 4th ed. edition (1958). [3](#)
- [2] K. Jones-Smith. *Non-Hermitian Quantum Mechanics*. PhD thesis, Case Western Reserve University (2010). [3](#), [11](#)
- [3] C. M. Bender, D. C. Brody, and H. F. Jones. *Must a Hamiltonian be Hermitian?* American Journal of Physics **71**, 1095 (2003). DOI: [10.1119/1.1574043](#). [3](#), [4](#), [5](#), [6](#), [7](#)
- [4] C. M. Bender and S. Boettcher. *Real Spectra in Non-Hermitian Hamiltonians Having  $PT$  Symmetry*. Phys. Rev. Lett. **80**, 5243 (1998). DOI: [10.1103/PhysRevLett.80.5243](#). [4](#), [5](#)
- [5] C. M. Bender. *Making sense of non-Hermitian Hamiltonians*. Reports on Progress in Physics **70**, 947 (2007). DOI: [10.1088/0034-4885/70/6/r03](#). [4](#), [5](#), [9](#), [18](#), [19](#)
- [6] C. F. de Morisson Faria and A. Fring. *Non-Hermitian Hamiltonians with real eigenvalues coupled to electric fields: From the time-independent to the time-dependent quantum mechanical formulation*. Laser Physics **17**, 424 (2007). DOI: [10.1134/S1054660X07040196](#). [4](#), [5](#), [19](#)
- [7] D. C. Brody. *Biorthogonal quantum mechanics*. Journal of Physics A: Mathematical and Theoretical **47**, 035305 (2013). DOI: [10.1088/1751-8113/47/3/035305](#). [5](#), [6](#), [17](#)
- [8] P. Dorey, C. Dunning, and R. Tateo. *Spectral equivalences, Bethe ansatz equations, and reality properties in  $PT$ -symmetric quantum mechanics*. Journal of Physics A: Mathematical and General **34**, 5679 (2001). DOI: [10.1088/0305-4470/34/28/305](#). [5](#)
- [9] D. C. Brody. *Consistency of  $PT$ -symmetric quantum mechanics*. Journal of Physics A: Mathematical and Theoretical **49**, 10LT03 (2016). DOI: [10.1088/1751-8113/49/10/10LT03](#). [5](#)
- [10] C. M. Bender and P. D. Mannheim.  *$PT$ -symmetry and necessary and sufficient conditions for the reality of energy eigenvalues*. Physics Letters A **374**, 1616 (2010). DOI: [10.1016/j.physleta.2010.02.032](#). [5](#)
- [11] C. M. Bender, S. Boettcher, and P. N. Meisinger.  *$PT$ -symmetric quantum mechanics*. Journal of Mathematical Physics **40**, 2201 (1999). DOI: [10.1063/1.532860](#). [5](#), [6](#), [19](#), [20](#)

- [12] A. Mostafazadeh. *Pseudo-Hermiticity versus  $PT$  symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian*. Journal of Mathematical Physics **43**, 205 (2002). DOI: [10.1063/1.1418246](https://doi.org/10.1063/1.1418246). 5, 17, 18
- [13] A. Mostafazadeh. *Exact  $PT$ -symmetry is equivalent to Hermiticity*. Journal of Physics A: Mathematical and General **36**, 7081 (2003). DOI: [10.1088/0305-4470/36/25/312](https://doi.org/10.1088/0305-4470/36/25/312). 5, 11, 18
- [14] C. M. Bender, J. Brod, A. Refig, and M. E. Reuter. *The  $C$  operator in  $PT$ -symmetric quantum theories*. Journal of Physics A: Mathematical and General **37**, 10139 (2004). DOI: [10.1088/0305-4470/37/43/009](https://doi.org/10.1088/0305-4470/37/43/009). 6, 7
- [15] C. M. Bender, D. C. Brody, and H. F. Jones. *Complex Extension of Quantum Mechanics*. Phys. Rev. Lett. **89**, 270401 (2002). DOI: [10.1103/PhysRevLett.89.270401](https://doi.org/10.1103/PhysRevLett.89.270401). 6, 7, 11
- [16] A. Mostafazadeh. *A Critique of  $PT$ -Symmetric Quantum Mechanics*, (2003). DOI: [10.48550/ARXIV.QUANT-PH/0310164](https://doi.org/10.48550/ARXIV.QUANT-PH/0310164). 6
- [17] A. Kumar, K. W. Murch, and Y. N. Joglekar. *Maximal quantum entanglement at exceptional points via unitary and thermal dynamics*. Phys. Rev. A **105**, 012422 (2022). DOI: [10.1103/PhysRevA.105.012422](https://doi.org/10.1103/PhysRevA.105.012422). 7
- [18] A. Bossart and R. Fleury. *Non-Hermitian time evolution: From static to parametric instability*. Phys. Rev. A **104**, 042225 (2021). DOI: [10.1103/PhysRevA.104.042225](https://doi.org/10.1103/PhysRevA.104.042225). 8, 10
- [19] N. Moiseyev. *Non-Hermitian Quantum Mechanics*. Cambridge University Press (2011). DOI: [10.1017/CBO9780511976186](https://doi.org/10.1017/CBO9780511976186). 8, 17
- [20] H. Cartarius. *Quantum systems with balanced gain and loss, signatures of branch points, and dissociation effects*. PhD thesis (2014). 8, 10, 12
- [21] F. Bagarello, F. Gargano, M. Lattuca, R. Passante, L. Rizzuto, and S. Spagnolo. *Exceptional Points in a Non-Hermitian Extension of the Jaynes-Cummings Hamiltonian*. In F. Bagarello, R. Passante, and C. Trapani (editors), *Non-Hermitian Hamiltonians in Quantum Physics*, pages 83–95. Springer International Publishing (2016). 8
- [22] W. D. Heiss. *The physics of exceptional points*. Journal of Physics A: Mathematical and Theoretical **45**, 444016 (2012). DOI: [10.1088/1751-8113/45/44/444016](https://doi.org/10.1088/1751-8113/45/44/444016). 8
- [23] P. Dorey, C. Dunning, and R. Tateo. *A Reality Proof in  $PT$ -Symmetric Quantum Mechanics*. Czechoslovak Journal of Physics **54**, 35 (2004). DOI: [10.1023/b:cjop.0000014365.19507.b6](https://doi.org/10.1023/b:cjop.0000014365.19507.b6). 8
- [24] C. M. Bender, N. Hassanpour, D. W. Hook, S. P. Klevansky, C. Sünderhauf, and Z. Wen. *Behavior of eigenvalues in a region of broken  $PT$ -symmetry*. Physical Review A **95** (2017). DOI: [10.1103/physreva.95.052113](https://doi.org/10.1103/physreva.95.052113). 9
- [25] C. M. Bender, D. C. Brody, H. F. Jones, and B. K. Meister. *Faster than Hermitian Quantum Mechanics*. Phys. Rev. Lett. **98**, 040403 (2007). DOI: [10.1103/PhysRevLett.98.040403](https://doi.org/10.1103/PhysRevLett.98.040403). 11, 12, 15, 18

- [26] M. A. Nielsen, M. R. Dowling, M. Gu, and A. C. Doherty. *Optimal control, geometry, and quantum computing*. Phys. Rev. A **73**, 062323 (2006). DOI: [10.1103/PhysRevA.73.062323](https://doi.org/10.1103/PhysRevA.73.062323). 11
- [27] A. Mostafazadeh. *Quantum Brachistochrone Problem and the Geometry of the State Space in Pseudo-Hermitian Quantum Mechanics*. Physical review letters **99**, 130502 (2007). DOI: [10.1103/PhysRevLett.99.130502](https://doi.org/10.1103/PhysRevLett.99.130502). 11
- [28] U. Günther and B. F. Samsonov. *The  $PT$ -symmetric brachistochrone problem, Lorentz boosts and non-unitary operator equivalence classes*. Physical Review A **78** (2008). DOI: [10.1103/physreva.78.042115](https://doi.org/10.1103/physreva.78.042115). 11
- [29] J. Anandan and Y. Aharonov. *Geometry of quantum evolution*. Phys. Rev. Lett. **65**, 1697 (1990). DOI: [10.1103/PhysRevLett.65.1697](https://doi.org/10.1103/PhysRevLett.65.1697). 12
- [30] C. M. Bender, D. C. Brody, L. P. Hughston, and B. K. Meister. *Geometric Aspects of Space-Time Reflection Symmetry in Quantum Mechanics*. In F. Bagarello, R. Pasante, and C. Trapani (editors), *Non-Hermitian Hamiltonians in Quantum Physics*, pages 185–199. Springer International Publishing (2016). 13
- [31] J. M. Zeuner, M. C. Rechtsman, Y. Plotnik, Y. Lumer, S. Nolte, M. S. Rudner, M. Segev, and A. Szameit. *Observation of a Topological Transition in the Bulk of a Non-Hermitian System*. Phys. Rev. Lett. **115**, 040402 (2015). DOI: [10.1103/PhysRevLett.115.040402](https://doi.org/10.1103/PhysRevLett.115.040402). 13
- [32] M. S. Rudner and L. S. Levitov. *Topological Transition in a Non-Hermitian Quantum Walk*. Phys. Rev. Lett. **102**, 065703 (2009). DOI: [10.1103/PhysRevLett.102.065703](https://doi.org/10.1103/PhysRevLett.102.065703). 13
- [33] T. Jacqmin, I. Carusotto, I. Sagnes, M. Abbarchi, D. D. Solnyshkov, G. Malpuech, E. Galopin, A. Lemaître, J. Bloch, and A. Amo. *Direct Observation of Dirac Cones and a Flatband in a Honeycomb Lattice for Polaritons*. Phys. Rev. Lett. **112**, 116402 (2014). DOI: [10.1103/PhysRevLett.112.116402](https://doi.org/10.1103/PhysRevLett.112.116402). 13
- [34] T. Goldzak, A. A. Mailybaev, and N. Moiseyev. *Light Stops at Exceptional Points*. Phys. Rev. Lett. **120**, 013901 (2018). DOI: [10.1103/PhysRevLett.120.013901](https://doi.org/10.1103/PhysRevLett.120.013901). 13, 14
- [35] P. Reséndiz-Vázquez, K. Tschernig, A. Perez-Leija, K. Busch, and R. de J León-Montiel. *Topological protection in non-Hermitian Haldane honeycomb lattices*. Phys. Rev. Research **2**, 013387 (2020). DOI: [10.1103/PhysRevResearch.2.013387](https://doi.org/10.1103/PhysRevResearch.2.013387). 14
- [36] C.-Z. Chang, J. Zhang, X. Feng, J. Shen, Z. Zhang, M. Guo, K. Li, Y. Ou, P. Wei, L.-L. Wang, Z.-Q. Ji, Y. Feng, S. Ji, X. Chen, J. Jia, X. Dai, Z. Fang, S.-C. Zhang, K. He, Y. Wang, L. Lu, X.-C. Ma, and Q.-K. Xue. *Experimental Observation of the Quantum Anomalous Hall Effect in a Magnetic Topological Insulator*. Science **340**, 167 (2013). DOI: [10.1126/science.1234414](https://doi.org/10.1126/science.1234414). 14
- [37] V. M. M. Alvarez, J. E. B. Vargas, and L. E. F. F. Torres. *Non-Hermitian robust edge states in one dimension: Anomalous localization and eigenspace condensation at exceptional points*. Phys. Rev. B **97**, 121401 (2018). DOI: [10.1103/PhysRevB.97.121401](https://doi.org/10.1103/PhysRevB.97.121401). 14



- [38] T. Yoshida, K. Kudo, and Y. Hatsugai. *Non-Hermitian fractional quantum Hall states*. Scientific Reports **9** (2019). DOI: [10.1038/s41598-019-53253-8](https://doi.org/10.1038/s41598-019-53253-8). 14
- [39] C. M. Bender. *Upside-down potentials*. Journal of Physics: Conference Series **343**, 012014 (2012). DOI: [10.1088/1742-6596/343/1/012014](https://doi.org/10.1088/1742-6596/343/1/012014). 15
- [40] A. Mostafazadeh. *Pseudo-Hermiticity versus PT-symmetry III: Equivalence of pseudo-Hermiticity and the presence of antilinear symmetries*. Journal of Mathematical Physics **43**, 3944 (2002). DOI: [10.1063/1.1489072](https://doi.org/10.1063/1.1489072). 17, 18
- [41] C. M. Bender and B. Tan. *Calculation of the hidden symmetry operator for a PT-symmetric square well*. Journal of Physics A: Mathematical and General **39**, 1945 (2006). DOI: [10.1088/0305-4470/39/8/011](https://doi.org/10.1088/0305-4470/39/8/011). 18
- [42] C. M. Bender, J.-H. Chen, and K. A. Milton. *PT-symmetric versus Hermitian formulations of quantum mechanics*. Journal of Physics A: Mathematical and General **39**, 1657 (2006). DOI: [10.1088/0305-4470/39/7/010](https://doi.org/10.1088/0305-4470/39/7/010). 18
- [43] H. F. Jones. *On pseudo-Hermitian Hamiltonians and their Hermitian counterparts*. Journal of Physics A: Mathematical and General **38**, 1741 (2005). DOI: [10.1088/0305-4470/38/8/010](https://doi.org/10.1088/0305-4470/38/8/010). 18
- [44] H. F. Jones and J. Mateo. *Equivalent Hermitian Hamiltonian for the non-Hermitian  $-x^4$  potential*. Phys. Rev. D **73**, 085002 (2006). DOI: [10.1103/PhysRevD.73.085002](https://doi.org/10.1103/PhysRevD.73.085002). 18
- [45] H. F. Jones and J. Mateo. *Pseudo-Hermitian Hamiltonians: Tale of two potentials*. Czechoslovak Journal of Physics **55**, 1117 (2005). DOI: [10.1007/s10582-005-0116-9](https://doi.org/10.1007/s10582-005-0116-9). 18
- [46] C. M. Bender and A. Turbiner. *Analytic continuation of eigenvalue problems*. Physics Letters A **173**, 442 (1993). DOI: [https://doi.org/10.1016/0375-9601\(93\)90153-Q](https://doi.org/10.1016/0375-9601(93)90153-Q). 19
- [47] C. M. Bender and S. A. Orszag. *Advanced Mathematical Methods for Scientists and Engineers I*. Springer New York, NY (1999). DOI: [10.1007/978-1-4757-3069-2](https://doi.org/10.1007/978-1-4757-3069-2). 19
- [48] M. Sorrell. *Complex WKB analysis of a  $\gamma$ -symmetric eigenvalue problem*. Journal of Physics A: Mathematical and Theoretical **40**, 10319 (2007). DOI: [10.1088/1751-8113/40/33/023](https://doi.org/10.1088/1751-8113/40/33/023). 19