

\mathcal{PT} -symmetric quantum mechanics

Ana Fabela Hinojosa ¹

Supervisors:

Dr. Jesper Levinsen

Prof. Meera Parish

SCHOOL OF PHYSICS & ASTRONOMY



MONASH University

November 2022

¹acfab1@student.monash.edu.au

Contents

1	Introduction	1
1.1	\mathcal{PT} -symmetry	1
1.1.1	The \mathcal{PT} operator	2
1.2	A new inner product	2
1.2.1	The \mathcal{CPT} inner product	2
1.2.2	The \mathcal{C} operator	3
1.3	Time evolution	4
1.3.1	2×2 \mathcal{PT} -symmetric Hamiltonian	4
2	Equivalent Hamiltonians with distinct symmetries	6
3	Project statement	7
4	Conclusion	8
5	Appendix	9

Abstract

IN LESS THAN 100 WORDS

Introduction

In “The principles of quantum mechanics”, Paul Dirac advises us that “it is important to remember that science is concerned only with *observable* things and that we can observe an object only by letting it interact with some outside influence”[1]. This means that in order to study valid physical systems, these systems must satisfy the fundamental postulates of quantum mechanical theory. Nearly all these postulates are based in physical properties. For example, one postulate establishes that time evolution of a quantum system must be *unitary* (i.e. probability conserving). Another requires that the energy spectrum of the system is bounded below so a lowest energy state can be measured. In quantum mechanics, the Hamiltonian operator (\hat{H}) encapsulates the total energy of a system. As explained already, a system’s energy spectrum is required to be bounded and real in order to be measurable. According to the fundamental postulates of the standard theory, if the operator \hat{H} satisfies the mathematical property known as Hermiticity –these operators are also known as self-adjoint in mathematics–, then \hat{H} will be an adequate physical observable. Interestingly, the Hermiticity postulate stands out from others in the conventional theory of quantum mechanics because it is mathematical rather than physical in its character[2]. An operator \hat{O} that is Hermitian has the property that its effect on the vectors of the Hilbert space in which \hat{O} is defined is independent of the order in which \hat{O} acts on said vectors[3]

$$\hat{O} |\psi\rangle = \langle\psi| \hat{O}^\dagger. \quad (1.1)$$

Despite of the correctness of Hermiticity, some believe that perhaps we have ended up with an overly restrictive quantum theory. The aim of this review is to summarize the present developments of a quantum theory that uses an alternative postulate to Hermiticity. This alternative postulate is referred to as space–time reflection symmetry (\mathcal{PT} symmetry)[4].

1.1 \mathcal{PT} -symmetry

In the late nineties, Carl Bender et al presented a \mathcal{PT} -symmetric theory of quantum mechanics. Their aim was to explain a conjecture on the reality and positiveness of the spectrum of a non-Hermitian Hamiltonian proposed by Bessis[5]. \mathcal{PT} -symmetric theory can be viewed as an analytical continuation of the conventional theory from real into the complex phase space[6].

An important question that must be answered, is whether a \mathcal{PT} -symmetric Hamiltonian defines a valid physical theory of quantum mechanics. By a physical theory we mean that the following conditions must be satisfied; The energy spectra of a system described by \hat{H} must be real and bounded below. Another condition is related to the probabilistic interpretation of the norm of a state, a norm must be always positive to be

a valid probability. Finally time evolution under the theory must be unitary. This means that as a state vector evolves in time the state's probability does not leak away[4][2].

1.1.1 The \mathcal{PT} operator

The \mathcal{PT} operator is the anti-linear operator composed of the linear parity operator (\mathcal{P}), which performs spatial reflection, and the anti-linear time-reversal operator (\mathcal{T}). These operators act on position and momentum operators in the following form

$$\begin{aligned}\mathcal{P} : \quad \hat{x} &\rightarrow -\hat{x}, & \hat{p} &\rightarrow -\hat{p}, \\ \mathcal{T} : \quad \hat{x} &\rightarrow \hat{x}, & \hat{p} &\rightarrow -\hat{p}, \quad i \rightarrow -i.\end{aligned}\tag{1.2}$$

Some Hamiltonians may not be symmetric under \mathcal{P} or \mathcal{T} separately, but Hamiltonians that remain invariant under the influence of the \mathcal{PT} operator are labelled as \mathcal{PT} -symmetric. A Hamiltonian \hat{H} will possess unbroken \mathcal{PT} symmetry if it's eigenstates are simultaneously eigenstates of the \mathcal{PT} operator, or in other words, if \hat{H} and the \mathcal{PT} operator commute. If the \mathcal{PT} operator and \hat{H} do not commute we say that the Hamiltonian's \mathcal{PT} -symmetry is broken[2][7][4]. If the symmetry is unbroken, then the eigenspectrum of \hat{H} is fully real and bounded below—This is also known as exact \mathcal{PT} -symmetry—. The effectiveness of \mathcal{PT} symmetry as a tool to investigate the spectra of some non-Hermitian Hamiltonians has been proved rigorously in various works, such as Dorey et al[8], Bender and Boettcher[5], Brody[9], Bender and Mannheim [10], Bender et al[?], Mostafazadeh[11][12] amongst others.

1.2 A new inner product

To ensure that the eigenstates $\{|\psi_n\rangle\}$ of a \mathcal{PT} -symmetric Hamiltonian \tilde{H} are orthogonal, it is necessary to specify an inner product. A possible choice for the inner product associated with \tilde{H} might be[6]

$$\langle\psi_n|\psi_m\rangle \equiv \int_c dx [\psi_n(x)]^{\mathcal{PT}} \psi_m(x) = \int_c dx [\psi_n(-x)]^* \psi_m(x)\tag{1.3}$$

where c is a complex contour that starts and ends within specific boundary conditions (see section 5). Unfortunately this obvious choice for the \mathcal{PT} inner product is not correct because it does not guarantee a positive definite norm.

1.2.1 The \mathcal{CPT} inner product

To be able to describe precisely the nature of \mathcal{PT} -symmetric quantum mechanics, we must delve briefly into the inner-product under which our theory satisfies the postulates of conventional quantum mechanics. It is important to note that \mathcal{PT} -symmetric quantum mechanics is a kind of ‘bootstrap’ theory[2], since infinitely many inner-products exist for a given vector space, we can construct an inner product whose associated norm is positive definite by design. This inner-product is in general dependent on the characteristics of the Hamiltonian in question and it guarantees that the underlying dynamics of any \mathcal{PT} -symmetric Hamiltonian satisfies unitarity[4]. Firstly, it is necessary to solve for the eigenstates of the Hamiltonian before knowing the Hilbert space and consequentially the associated inner product. To guarantee a positive norm for our theory, we will construct

a new linear operator \mathcal{C} that commutes with both \hat{H} and \mathcal{PT} . We use the symbol \mathcal{C} to represent this symmetry because its properties are similar to those of the charge conjugation operator in particle physics[2].

1.2.2 The \mathcal{C} operator

When the \mathcal{PT} -symmetry of \hat{H} is unbroken, then \hat{H} and \mathcal{PT} commute. This statement is equivalent to saying that the eigenfunctions $\phi_n(x)$ of \hat{H} are simultaneously eigenstates of the \mathcal{PT} operator[13]

$$\mathcal{PT}\phi_n(x) = \lambda_n\phi_n(x), \quad (1.4)$$

Where λ_n is a pure phase. Without loss of generality, for each n the phase can be absorbed into $\phi_n(x)$ and this makes the eigenvalue of the \mathcal{PT} operator unity[13]:

$$\mathcal{PT}\phi_n(x) = \phi_n^*(-x) = \phi_n(x). \quad (1.5)$$

There is strong numerical evidence of the completeness of the eigenfunctions $\phi_n(x)$ [7][13][14]. In the coordinate basis, the completeness statement reads:

$$\sum_n (-1)^n \phi_n(x) \phi_n(y) = \delta(x - y), \quad x, y \in \mathbb{R} \quad (1.6)$$

the unconventional $(-1)^n$ factor in ?? can be explained if we define the \mathcal{PT} inner product as

$$\begin{aligned} (f, g) &= \int dx [\mathcal{PT}f(x)] g(x) \\ &= \int dx f^*(-x) g(x). \end{aligned} \quad (1.7)$$

where the integral above follows a path in the complex plane. Under this definition the eigenstate norms alternate in sign depending on the value n . This means that the metric associated with the \mathcal{PT} inner product is indefinite[13][15]. In quantum theory, the norm of states is interpreted as a probability and this means that the indefinite metric described above presents a serious problem for the validity of \mathcal{PT} -symmetric quantum theory. The solution to this problem lies in finding an interpretation for the negative valued norms[?].

Most model \mathcal{PT} -symmetric Hamiltonians \tilde{H} in the literature admit tuneable parameters such that there will be regions in the parameter space where the eigenstates of \tilde{H} are not \mathcal{PT} -symmetric[14]. This means that a system described by \tilde{H} admits two phases, one for broken and another for unbroken \mathcal{PT} -symmetry. In the unbroken phase eigenstates of \tilde{H} become degenerate and this implies that they lose completeness[14]. This is the one reason why in this review we are focused in the unbroken phase. The literature claims that for any theory with unbroken \mathcal{PT} -symmetry there exists a symmetry of the Hamiltonian that describes the negative and positive norm states. To describe this symmetry of \hat{H} it is necessary to construct a linear operator denoted by \mathcal{C} [4][7][13]. When represented in position space \mathcal{C} is a sum over the energy eigenstates of \hat{H} :

$$\mathcal{C} = \sum_n \phi_n(x) \phi_n(y) \quad (1.8)$$

From this definition, and the relation $(\phi_m(y), \phi_n(y)) = \int dy \phi_m(y) \phi_n(y)$ we can verify that the eigenvalues of \mathcal{C} are ± 1

$$\begin{aligned} \mathcal{C}\phi_n(x) &= \int dy \mathcal{C} \phi_n(y) \\ &= \sum_m \phi_m(x) \int dy \phi_m(y) \phi_n(y) \\ &= (-1)^n \phi_n(x) \end{aligned} \tag{1.9}$$

The \mathcal{PT} norms signatures can therefore be interpreted as the “charge” of the states, while \mathcal{C} is the operator used to measure this charge[13].

The \mathcal{C} operator commutes with the \mathcal{PT} operator, but it does not commute with the parity operator \mathcal{P} . Notice that \mathcal{C} and \mathcal{P} operators are square roots of $\delta(x - y)$ the unity operator[7]

$$\mathcal{P}^2 = \mathcal{C}^2 = \mathbb{1}, \tag{1.10}$$

where $\mathcal{P} \neq \mathcal{C}$, since \mathcal{P} is real and \mathcal{C} is complex valued[4][13].

Using the newly constructed \mathcal{C} operator, we can redesign the \mathcal{PT} inner product to suit the conventional probabilistic interpretation of the vector norms in quantum mechanics

$$(f, g) = \int dx [\mathcal{CPT}f(x)] g(x). \tag{1.11}$$

1.3 Time evolution

The aim of this section is to illustrate the dynamics of non-Hermitian \mathcal{PT} -symmetric Hamiltonians. In Hermitian quantum mechanics, time-evolution of states is done by using the unitary operator $\hat{U} = e^{-i\hat{H}t/\hbar}$. Because the \hat{U} operator is unitary this means that as the state $\vec{\psi}$ evolves in time, its norm remains constant. This is useful in quantum mechanics because the norm of a state is interpreted as a probability, and therefore unitarity guarantees that a state’s probability does not grow or decay with time. As explained in section 1.2.1, when the \mathcal{PT} -symmetry of \hat{H} is unbroken we have positive definite \mathcal{CPT} norm states. This means that time evolution under the unbroken \mathcal{PT} -symmetric framework is also unitary[3][7][12]. The unitary time evolution of a state $\vec{\psi}$ from time $0 \rightarrow t$ is written as

$$\vec{\psi}(t) = \hat{U}\vec{\psi}(0). \tag{1.12}$$

1.3.1 2×2 \mathcal{PT} -symmetric Hamiltonian

A simple example of Non-Hermitian time evolution that is often found in the literature describes a two-level \mathcal{PT} -symmetric Hamiltonian \tilde{H} whose eigenvectors span the Hilbert space \mathcal{H} [16]. Without loss of generality, since we are dealing with a two-level system, it is useful to use the Bloch sphere to visualise the geometry of this system. Let the eigenvectors of \tilde{H} be

$$|\psi_i\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |\psi_f\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{1.13}$$

There are many possible paths for the time evolution from $|\psi_i\rangle$ to $|\psi_f\rangle$ as is visible in figure 1.1.




Figure 1.1

Equivalent Hamiltonians with distinct symmetries

In parallel to the work of Bender is the research of Mostafazadeh who introduced the notion of pseudo-Hermiticity in [11]. A Hamiltonian is said to be pseudo-Hermitian with respect to a positive-definite, Hermitian operator η if it satisfies

$$\tilde{H}^\dagger = \eta^{-1} \tilde{H} \eta. \quad (2.1)$$

In the case of Hamiltonians with \mathcal{PT} -symmetry, the role of η is played by the combined action of \mathcal{P} and \mathcal{C} . A convenient way to write the \mathcal{C} operator was proposed in [17]

$$\mathcal{C} = e^Q \mathcal{P} = \eta^{-1} \mathcal{P}, \quad (2.2)$$

here Q is an antisymmetric Hermitian operator. Mostafazadeh [12] has shown that the square root of the positive-definite Hermitian operator $\eta = e^{-Q}$ can be used to transform any non-Hermitian Hamiltonian with unbroken \mathcal{PT} -symmetry into a spectrally equivalent Hermitian Hamiltonian by means of a unitary “similarity transformation” [18][19]. The transformation is as follows, the invertible operator $\rho = \sqrt{\eta}$ acts on the non-Hermitian \mathcal{PT} -symmetric Hamiltonian \tilde{H} and returns an equivalent Hermitian Hamiltonian \hat{H}

$$\hat{H} = \rho^{-1} \tilde{H} \rho = e^{-Q/2} \tilde{H} e^{Q/2}. \quad (2.3)$$

To verify that the similar Hamiltonian \hat{H} is Hermitian we take the Hermitian conjugate of \hat{H}

$$\begin{aligned} \hat{H}^\dagger &= (e^{-Q/2} \tilde{H} e^{Q/2})^\dagger, \\ &= e^{Q/2} \tilde{H}^\dagger e^{-Q/2}, \end{aligned} \quad (2.4)$$

If we “swap” Hermiticity: \tilde{H}^\dagger in 2.4 for \mathcal{PT} -symmetry, and we use equation ??

$$\begin{aligned} \hat{H}^\dagger &= e^{Q/2} \mathcal{P} \tilde{H} \mathcal{P}^{-1} e^{-Q/2}, \\ &= e^{-Q/2} \mathcal{C} \tilde{H} \mathcal{C}^{-1} e^{Q/2}, \end{aligned} \quad (2.5)$$

Finally we recall that \mathcal{C} and \tilde{H} commute

$$\hat{H}^\dagger = e^{-Q/2} \tilde{H} e^{Q/2} = \hat{H}. \quad (2.6)$$

Mostafazadeh [11], conjectures that because \mathcal{CPT} -symmetry satisfies the postulates of quantum mechanics –whilst only “swapping” the Hermiticity of a Hamiltonian by \mathcal{CPT} -symmetry– then this must mean that non Hermitian \mathcal{CPT} -symmetric theories are equivalent to certain non local Hermitian field theories. It is natural to notice that this equivalence feature between both theories could provide an advantage to simplify quantum mechanical calculations, this is explored in references [11],[20],[21],[19],[22].

Project statement

Conclusion

Appendix

One-to-one equivalence:

PT-symmetric and Hermitian quantum theories

parametric family of PTsNHH

Stokes wedges and boundary conditions

Bi-orthogonality and completeness

Bibliography

- [1] P. A. M. Dirac. *THE PRINCIPLES OF QUANTUM MECHANICS*. International series of monographs on physics (Oxford, England). Clarendon Pr., Oxf., 4th ed. edition (1958). [1](#)
- [2] C. M. Bender. *Making sense of non-Hermitian Hamiltonians*. Reports on Progress in Physics **70**, 947 (2007). DOI: [10.1088/0034-4885/70/6/r03](#). [1](#), [2](#), [3](#)
- [3] K. Jones-Smith. *Non-Hermitian Quantum Mechanics*. PhD thesis, Case Western Reserve University (2010). [1](#), [4](#)
- [4] C. M. Bender, D. C. Brody, and H. F. Jones. *Must a Hamiltonian be Hermitian?* American Journal of Physics **71**, 1095 (2003). DOI: [10.1119/1.1574043](#). [1](#), [2](#), [3](#), [4](#)
- [5] C. M. Bender and S. Boettcher. *Real Spectra in Non-Hermitian Hamiltonians Having PT -Symmetry*. Phys. Rev. Lett. **80**, 5243 (1998). DOI: [10.1103/PhysRevLett.80.5243](#). [1](#), [2](#)
- [6] C. M. Bender, S. Boettcher, and P. N. Meisinger. *PT -symmetric quantum mechanics*. Journal of Mathematical Physics **40**, 2201 (1999). DOI: [10.1063/1.532860](#). [1](#), [2](#)
- [7] C. M. Bender, D. C. Brody, and H. F. Jones. *Complex Extension of Quantum Mechanics*. Phys. Rev. Lett. **89**, 270401 (2002). DOI: [10.1103/PhysRevLett.89.270401](#). [2](#), [3](#), [4](#)
- [8] P. Dorey, C. Dunning, and R. Tateo. *Spectral equivalences, Bethe ansatz equations, and reality properties in PT -symmetric quantum mechanics*. Journal of Physics A: Mathematical and General **34**, 5679 (2001). DOI: [10.1088/0305-4470/34/28/305](#). [2](#)
- [9] D. C. Brody. *Consistency of PT -symmetric quantum mechanics*. Journal of Physics A: Mathematical and Theoretical **49**, 10LT03 (2016). DOI: [10.1088/1751-8113/49/10/10LT03](#). [2](#)
- [10] C. M. Bender and P. D. Mannheim. *PT -symmetry and necessary and sufficient conditions for the reality of energy eigenvalues*. Physics Letters A **374**, 1616 (2010). DOI: [10.1016/j.physleta.2010.02.032](#). [2](#)
- [11] A. Mostafazadeh. *Pseudo-Hermiticity versus PT symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian*. Journal of Mathematical Physics **43**, 205 (2002). DOI: [10.1063/1.1418246](#). [2](#), [6](#)
- [12] A. Mostafazadeh. *Exact PT -symmetry is equivalent to Hermiticity*. Journal of Physics A: Mathematical and General **36**, 7081 (2003). DOI: [10.1088/0305-4470/36/25/312](#). [2](#), [4](#), [6](#)

- [13] C. M. Bender, J. Brod, A. Refig, and M. E. Reuter. *The C operator in PT -symmetric quantum theories*. Journal of Physics A: Mathematical and General **37**, 10139 (2004). DOI: [10.1088/0305-4470/37/43/009](https://doi.org/10.1088/0305-4470/37/43/009). 3, 4
- [14] D. C. Brody. *Biorthogonal quantum mechanics*. Journal of Physics A: Mathematical and Theoretical **47**, 035305 (2013). DOI: [10.1088/1751-8113/47/3/035305](https://doi.org/10.1088/1751-8113/47/3/035305). 3
- [15] A. Mostafazadeh. *A Critique of PT -Symmetric Quantum Mechanics*, (2003). DOI: [10.48550/ARXIV.QUANT-PH/0310164](https://doi.org/10.48550/ARXIV.QUANT-PH/0310164). 3
- [16] C. M. Bender, D. C. Brody, H. F. Jones, and B. K. Meister. *Faster than Hermitian Quantum Mechanics*. Phys. Rev. Lett. **98**, 040403 (2007). DOI: [10.1103/PhysRevLett.98.040403](https://doi.org/10.1103/PhysRevLett.98.040403). 4
- [17] C. M. Bender and B. Tan. *Calculation of the hidden symmetry operator for a PT -symmetric square well*. Journal of Physics A: Mathematical and General **39**, 1945 (2006). DOI: [10.1088/0305-4470/39/8/011](https://doi.org/10.1088/0305-4470/39/8/011). 6
- [18] C. M. Bender, J.-H. Chen, and K. A. Milton. *PT -symmetric versus Hermitian formulations of quantum mechanics*. Journal of Physics A: Mathematical and General **39**, 1657 (2006). DOI: [10.1088/0305-4470/39/7/010](https://doi.org/10.1088/0305-4470/39/7/010). 6
- [19] H. F. Jones. *On pseudo-Hermitian Hamiltonians and their Hermitian counterparts*. Journal of Physics A: Mathematical and General **38**, 1741 (2005). DOI: [10.1088/0305-4470/38/8/010](https://doi.org/10.1088/0305-4470/38/8/010). 6
- [20] H. F. Jones and J. Mateo. *Equivalent Hermitian Hamiltonian for the non-Hermitian $-x^4$ potential*. Phys. Rev. D **73**, 085002 (2006). DOI: [10.1103/PhysRevD.73.085002](https://doi.org/10.1103/PhysRevD.73.085002). 6
- [21] A. Mostafazadeh. *Pseudo-Hermiticity versus PT -symmetry III: Equivalence of pseudo-Hermiticity and the presence of antilinear symmetries*. Journal of Mathematical Physics **43**, 3944 (2002). DOI: [10.1063/1.1489072](https://doi.org/10.1063/1.1489072). 6
- [22] H. F. Jones and J. Mateo. *Pseudo-hermitian hamiltonians: Tale of two potentials*. Czechoslovak Journal of Physics **55**, 1117 (2005). DOI: [10.1007/s10582-005-0116-9](https://doi.org/10.1007/s10582-005-0116-9). 6