

Faster than Hermitian time-evolution

by Carl M Bender

Ana Fabela Hinojosa



MONASH University

Supervisors:
Jesper Levinsen
Meera Parish

Outline

Introduction

Time evolution

Brachistochrone problem

Conclusion

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Hilbert spaces

A Hilbert space is a vector space that can be infinite dimensional.

Vector spaces are equipped with an inner product.

Inner product:

Define a distance function.




Fig.3: Hilbert space stuff

Hamiltonians as Observables




Fig.1: The set of all possible Hamiltonians.

1. Observables are **self-adjoint** operators $\{\hat{O}, \hat{H}, \dots\}$
2. Real energy spectrum with defined lowest energy
3. A state ψ of the quantum system is a unit vector of \hat{H}
4. Expectation values of observables are given by the inner-product.
5. Unitarity

Hamiltonians as Observables




Fig.1: The set of all possible Hamiltonians.

1. Observables are **self-adjoint** operators $\{\hat{O}, \hat{H}, \dots\}$
2. Real energy spectrum with defined lowest energy
3. A state ψ of the quantum system is a unit vector of \hat{H}
4. Expectation values of observables are given by the inner-product.
5. Unitarity

Hamiltonians as Observables




Fig.1: The set of all possible Hamiltonians.

1. ~~Observables are self-adjoint operators~~ $\{\hat{O}, \hat{H}, \dots\}$
2. Real energy spectrum with defined lowest energy
3. A state ψ of the quantum system is a unit vector of \hat{H}
4. Expectation values of observables are given by the inner-product.
5. Unitarity

CPT - symmetry in short

Hilbert space $\rightarrow \hat{H} = \hat{H}^{CPT}$

$\mathcal{C} \rightarrow$ charge conjugation,

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ complex conjugation and time reversal.

Inner product is determined dynamically in terms of the Hamiltonian.

CPT - symmetry in short

Hilbert space $\rightarrow \hat{H} = \hat{H}^{CPT}$

$\mathcal{C} \rightarrow$ charge conjugation,

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ complex conjugation and time reversal.

Inner product is determined dynamically in terms of the Hamiltonian.

If the eigenfunctions of a \mathcal{PT} -symmetric Hamiltonian are **not** also eigenfunctions of the \mathcal{PT} operator we say the Hamiltonian possesses **broken \mathcal{PT} -symmetry**.

CPT - symmetry in short

Hilbert space $\rightarrow \hat{H} = \hat{H}^{CPT}$

$\mathcal{C} \rightarrow$ charge conjugation,

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ complex conjugation and time reversal.

Inner product is determined dynamically in terms of the Hamiltonian.

If the eigenfunctions of a \mathcal{PT} -symmetric Hamiltonian are **not** also eigenfunctions of the \mathcal{PT} operator we say the Hamiltonian possesses **broken \mathcal{PT} -symmetry**.

Interesting physical phenomena occurs in the broken symmetry region

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Time evolution

$$\vec{\psi}_i \rightarrow \vec{\psi}_f$$

Time evolution

$$\vec{\psi}_i \rightarrow \vec{\psi}_f \quad \vec{\psi}_f = \hat{U} \vec{\psi}_i$$

Time evolution

$$\begin{array}{ccc} \vec{\psi}_i & \rightarrow & \vec{\psi}_f \\ & \hookrightarrow & \end{array} \quad \vec{\psi}_f = \hat{U} \vec{\psi}_i$$

\downarrow

1. Hermitian quantum mechanics:

$$\hat{U} = e^{-i\hat{H}t/\hbar},$$

where $t > 0$.

Time evolution

$$\vec{\psi}_i \rightarrow \vec{\psi}_f \quad \hookrightarrow \quad \vec{\psi}_f = \hat{U} \vec{\psi}_i$$

\downarrow

1. Hermitian quantum mechanics:

$$\hat{U} = e^{-i\hat{H}t/\hbar},$$

where $t > 0$.

2. CPT quantum mechanics: ?

Contents

Introduction

Time evolution


Brachistochrone problem

Conclusion


The Classical Brachistochrone problem




The Classical Brachistochrone problem




The Classical Brachistochrone problem




The Classical Brachistochrone problem



The Classical Brachistochrone problem




The Classical Brachistochrone problem



βράχιστος χρόνος
brákhistos khrónos:
“shortest time”

Fig.2: A particle travels from left to right in time t .
Can we make this trip nearly instantaneous?

The Classical Brachistochrone problem



βράχιστος χρόνος
brákhistos khrónos:
“shortest time”

How fast can we evolve a state?

Fig.2: A particle travels from left to right in time t ,
Can we make this trip nearly instantaneous?

A simple quantum brachistochrone problem

This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.





Fig.3: Bloch sphere with initial and final states

A simple quantum brachistochrone problem




This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

We want the **fastest** time evolution possible, without violating the time-energy uncertainty principle.

Fig.3: Bloch sphere with initial and final states

A simple quantum brachistochrone problem



This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

We want the **fastest** time evolution possible, without violating the time-energy uncertainty principle.

The chosen Hamiltonian must satisfy:

Energy constraint:

$$\omega = E_{\max} - E_{\min}$$

Fig.3: Bloch sphere with initial and final states

A simple quantum brachistochrone problem




Fig.3: Bloch sphere with initial and final states

This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

We want the **fastest** time evolution possible, without violating the time-energy uncertainty principle.

The chosen Hamiltonian must satisfy:

Energy constraint:

$$\omega = E_{\max} - E_{\min}$$

Will a complex non-Hermitian Hamiltonian give time-optimal evolution?

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \{r, s, \theta, u\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = (s - u)^2 + 4r^2,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i\frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i\frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

CPT-symmetric case

$$\tilde{H} = \begin{pmatrix} re^{i\theta} & s \\ s & re^{-i\theta} \end{pmatrix}, \quad \{r, s, \theta\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = 4s^2 - 4r^2 \sin^2 \theta > 0,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\text{where } \sin(\alpha) = \frac{r}{s} \sin(\theta).$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s - u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

CPT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar} - \alpha) \\ -i \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

CPT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega|b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

CPT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

CPT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar} - \alpha) \\ -i \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos \frac{\omega t}{2\hbar} - \alpha}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

Since ω is fixed, $\omega^2 = 4s^2 - 4r^2 \sin^2(\theta)$

$$\rightarrow \omega^2 = 4s^2 \cos^2(\alpha)$$

we require $s, r \gg 1$.

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

CPT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

Since ω is fixed, $\omega^2 = 4s^2 - 4r^2 \sin^2(\theta)$

$$\rightarrow \omega^2 = 4s^2 \cos^2(\alpha)$$

we require $s, r \gg 1$.

Then $t \rightarrow 0$ when $\alpha \rightarrow -\frac{\pi}{2}$.

The geometry of the space

\mathcal{CPT} inner products are defined depending on the Hamiltonian used.

The geometry of the space

\mathcal{CPT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow Hermitian inner product

The geometry of the space

\mathcal{CPT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow Hermitian inner product
2. not orthogonal \rightarrow \mathcal{CPT} inner product

The geometry of the space

\mathcal{CPT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow **Hermitian** inner product
2. not orthogonal \rightarrow **CPT** inner product
vector separation: $\delta = \pi - 2|\alpha|$

The geometry of the space

\mathcal{CPT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow Hermitian inner product
2. not orthogonal \rightarrow \mathcal{CPT} inner product
vector separation: $\delta = \pi - 2|\alpha|$

We can choose α to create a “wormhole” effect in Hilbert space.

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Conclusion

1. This paper demonstrates the theoretical differences in both \mathcal{PT} -symmetric and conventional Hermitian quantum mechanics.
2. The “wormhole” effect could be of importance in design and implementation of fast quantum computation and communication algorithms.
3. There could be quantum protection mechanisms limiting the applicability of Hilbert-space “wormholes”.

References



C. M. Bender, D. C. Brody, H. F. Jones, and B. K. Meister.

Faster than hermitian quantum mechanics.

Phys. Rev. Lett. **98**, 040403 (2007).

DOI: [10.1103/PhysRevLett.98.040403](https://doi.org/10.1103/PhysRevLett.98.040403).