

Faster than Hermitian Quantum Mechanics

by Carl M Bender et al

Ana Fabela Hinojosa



MONASH University

Supervisors:
Jesper Levinsen
Meera Parish

Outline

Introduction

Time evolution

Brachistochrone problem

Conclusion

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Hilbert spaces

A Hilbert space is a vector space that can be infinite dimensional.

Vector spaces are equipped with an inner product.

Inner product:

Define a distance function.



Fig.1: Hilbert space stuff

Hamiltonians as Observables




Fig.2: The set of all possible Hamiltonians.

1. Observables are **self-adjoint** operators.
2. \hat{H} has a real energy spectrum bounded below.
3. Unitarity.

Hamiltonians as Observables




Fig.2: The set of all possible Hamiltonians.

1. Observables are **self-adjoint** operators.
2. \hat{H} has a real energy spectrum bounded below.
3. Unitarity.

PT - symmetry in short

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ time reversal (complex conjugation).

$$\hat{H} = \mathcal{PT} \hat{H} \mathcal{PT} = \hat{H}^{\mathcal{PT}}$$

PT - symmetry in short

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ time reversal (complex conjugation).

$$\hat{H} = \mathcal{PT} \hat{H} \mathcal{PT} = \hat{H}^{\mathcal{PT}}$$

- ▶ The \mathcal{PT} inner product is determined dynamically in terms of the Hamiltonian.

PT - symmetry in short

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ time reversal (complex conjugation).

$$\hat{H} = \mathcal{PT} \hat{H} \mathcal{PT} = \hat{H}^{\mathcal{PT}}$$

- ▶ The \mathcal{PT} inner product is determined dynamically in terms of the Hamiltonian.
- ▶ \mathcal{PT} -symmetry can be **broken** or **unbroken**.

PT - symmetry in short

$\mathcal{P} \rightarrow$ spatial inversion,

$\mathcal{T} \rightarrow$ time reversal (complex conjugation).

$$\hat{H} = \mathcal{PT} \hat{H} \mathcal{PT} = \hat{H}^{\mathcal{PT}}$$

- ▶ The \mathcal{PT} inner product is determined dynamically in terms of the Hamiltonian.
- ▶ \mathcal{PT} -symmetry can be **broken** or **unbroken**.
- ▶ Interesting physical phenomena occurs in the broken symmetry region.

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Time evolution

$$\vec{\psi}_i \rightarrow \vec{\psi}_f$$

Time evolution

$$\begin{array}{l} \vec{\psi}_i \rightarrow \vec{\psi}_f \\ \hookrightarrow \vec{\psi}_f = \hat{U} \vec{\psi}_i \end{array}$$

Time evolution

$$\begin{array}{ccc} \vec{\psi}_i & \rightarrow & \vec{\psi}_f \\ & \hookrightarrow & \\ & \vec{\psi}_f & = \hat{U} \vec{\psi}_i \end{array}$$

\downarrow

► Hermitian quantum mechanics:
 $\hat{U} = e^{-i\hat{H}t/\hbar}$

Time evolution

$$\begin{aligned}\vec{\psi}_i &\rightarrow \vec{\psi}_f \\ &\hookrightarrow \vec{\psi}_f = \hat{U} \vec{\psi}_i \\ &\quad \downarrow\end{aligned}$$

- ▶ Hermitian quantum mechanics:

$$\hat{U} = e^{-i\hat{H}t/\hbar}$$

- ▶ PT quantum mechanics:

$$\hat{U} = e^{-i\tilde{H}t/\hbar}$$

Time evolution

$$\begin{aligned}\vec{\psi}_i &\rightarrow \vec{\psi}_f \\ &\hookrightarrow \vec{\psi}_f = \hat{U} \vec{\psi}_i\end{aligned}$$

\downarrow

► Hermitian quantum mechanics:
 $\hat{U} = e^{-i\hat{H}t/\hbar}$

► PT quantum mechanics:
 $\hat{U} = e^{-i\tilde{H}t/\hbar}$
Unitary?

Contents

Introduction

Time evolution


Brachistochrone problem

Conclusion


The Classical Brachistochrone problem




The Classical Brachistochrone problem




The Classical Brachistochrone problem




The Classical Brachistochrone problem




The Classical Brachistochrone problem



The Classical Brachistochrone problem



The Classical Brachistochrone problem



βράχιστος χρόνος
brákhistos khrónos:
“shortest time”

How fast can we evolve a state?

Fig.3: A particle travels from left to right in time t ,
Can we make this trip nearly instantaneous?

A simple quantum brachistochrone problem

This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.




Fig.4: Bloch sphere with initial and final states

A simple quantum brachistochrone problem





Fig.4: Bloch sphere with initial and final states

This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

What is the **fastest** time evolution possible, that does not violate the time-energy uncertainty principle.

A simple quantum brachistochrone problem



This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

What is the **fastest** time evolution possible, that does not violate the time-energy uncertainty principle.

The Hamiltonian must satisfy an **energy constraint** based on:

$$\omega \equiv E_{\max} - E_{\min} > 0$$

Fig.4: Bloch sphere with initial and final states

A simple quantum brachistochrone problem




Fig.4: Bloch sphere with initial and final states

This space is spanned by $\vec{\psi}_i$ and $\vec{\psi}_f$.

What is the **fastest** time evolution possible, that does not violate the time-energy uncertainty principle.

The Hamiltonian must satisfy an **energy constraint** based on:

$$\omega \equiv E_{\max} - E_{\min} > 0$$

Will a complex non-Hermitian Hamiltonian give time-optimal evolution?

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \{r, s, \theta, u\} \in \mathbb{R},$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \{r, s, \theta, u\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = (s - u)^2 + 4r^2,$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \quad \{r, s, \theta, u\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = (s - u)^2 + 4r^2,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

PT-symmetric case

$$\tilde{H} = \begin{pmatrix} re^{i\theta} & s \\ s & re^{-i\theta} \end{pmatrix}, \quad \{r, s, \theta\} \in \mathbb{R},$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \quad \{r, s, \theta, u\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = (s - u)^2 + 4r^2,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

PT-symmetric case

$$\tilde{H} = \begin{pmatrix} re^{i\theta} & s \\ s & re^{-i\theta} \end{pmatrix}, \quad \{r, s, \theta\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = 4s^2 - 4r^2 \sin^2 \theta > 0,$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hermitian case

$$\hat{H} = \begin{pmatrix} s & re^{-i\theta} \\ re^{i\theta} & u \end{pmatrix}, \{r, s, \theta, u\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = (s - u)^2 + 4r^2,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

PT-symmetric case

$$\tilde{H} = \begin{pmatrix} re^{i\theta} & s \\ s & re^{-i\theta} \end{pmatrix}, \{r, s, \theta\} \in \mathbb{R},$$

Energy constraint

$$\omega^2 = 4s^2 - 4r^2 \sin^2 \theta > 0,$$

Time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar} - \alpha) \\ -i \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

where $\sin(\alpha) = \frac{r}{s} \sin(\theta)$.

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s - u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

PT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s - u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar} - \alpha) \\ -i \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar}) - i \frac{s-u}{\omega} \sin(\frac{\omega t}{2\hbar}) \\ -i \frac{2r}{\omega} e^{i\theta} \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s - u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

PT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos(\frac{\omega t}{2\hbar} - \alpha) \\ -i \sin(\frac{\omega t}{2\hbar}) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos \frac{\omega t}{2\hbar} - \alpha}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

PT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega|b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

PT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

Since ω is fixed, $\omega^2 = 4s^2 - 4r^2 \sin^2(\theta)$
 $\rightarrow \omega^2 = 4s^2 \cos^2(\alpha)$

The Maths

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_r = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Hermitian time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = e^{\frac{-i(s+u)t}{2\hbar}} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar}\right) - i \frac{s-u}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right) \\ -i \frac{2r}{\omega} e^{i\theta} \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |b| = \frac{2r}{\omega} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{2\hbar}{\omega} \arcsin\left(\frac{\omega |b|}{2r}\right),$$

For all $r > 0$ subject to $\omega^2 = (s-u)^2 + 4r^2$
for a fixed ω .

$$\therefore \tau = \frac{2\hbar}{\omega} \arcsin(|b|) = \frac{\pi\hbar}{\omega}$$

is the minimum *passage time*.

PT time evolution

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{e^{\frac{-itr \cos \theta}{\hbar}}}{\cos \alpha} \begin{pmatrix} \cos\left(\frac{\omega t}{2\hbar} - \alpha\right) \\ -i \sin\left(\frac{\omega t}{2\hbar}\right) \end{pmatrix}.$$

$$\Rightarrow |a| = \frac{\cos\left(\frac{\omega t}{2\hbar} - \alpha\right)}{\cos \alpha} \sin\left(\frac{\omega t}{2\hbar}\right),$$

$$\Rightarrow t = \frac{\hbar}{\omega} (2\alpha + \pi),$$

Since ω is fixed, $\omega^2 = 4s^2 - 4r^2 \sin^2(\theta)$
 $\rightarrow \omega^2 = 4s^2 \cos^2(\alpha)$

Then $t \rightarrow 0$ when $\alpha \rightarrow -\frac{\pi}{2}$.
but we also require $s, r \gg 1$.

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow Hermitian inner product

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow **Hermitian** inner product
vector separation: $\delta = 2 \arccos |\langle \psi_f | \psi_i \rangle| = \pi$

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow **Hermitian** inner product
vector separation: $\delta = 2 \arccos |\langle \psi_f | \psi_i \rangle| = \pi$
2. not orthogonal \rightarrow **PT** inner product

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow **Hermitian** inner product
vector separation: $\delta = 2 \arccos |\langle \psi_f | \psi_i \rangle| = \pi$
2. not orthogonal \rightarrow **PT** inner product
vector separation: $\delta = \pi - 2|\alpha|$

The geometry of the space

\mathcal{PT} inner products are defined depending on the Hamiltonian used.

$$\vec{\psi}_i = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \vec{\psi}_f = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

1. orthogonal \rightarrow **Hermitian** inner product
vector separation: $\delta = 2 \arccos |\langle \psi_f | \psi_i \rangle| = \pi$
2. not orthogonal \rightarrow **PT** inner product
vector separation: $\delta = \pi - 2|\alpha|$

We can choose α to create a “wormhole” effect in Hilbert space.

Contents

Introduction

Time evolution

Brachistochrone problem

Conclusion

Conclusion

1. This paper demonstrates the theoretical differences in both \mathcal{PT} -symmetric and conventional Hermitian quantum mechanics.
2. The “wormhole” effect could be of importance in design and implementation of fast quantum computation and communication algorithms.
3. There could be quantum protection mechanisms limiting the applicability of Hilbert-space “wormholes”.

References



C. M. Bender, D. C. Brody, H. F. Jones, and B. K. Meister.

Faster than hermitian quantum mechanics.

Phys. Rev. Lett. **98**, 040403 (2007).

DOI: [10.1103/PhysRevLett.98.040403](https://doi.org/10.1103/PhysRevLett.98.040403).

Light Stops at Exceptional Points

Tamar Goldzak,¹ Alexei A. Mailybaev,^{2,*} and Nimrod Moiseyev^{1,†}

¹*Schulich Faculty of Chemistry and Faculty of Physics, Technion—Israel Institute of Technology, Haifa 32000, Israel*

²*Instituto Nacional de Matemática Pura e Aplicada—IMPA, 22460-320 Rio de Janeiro, Brazil*



(Received 29 September 2017; published 3 January 2018)

Almost twenty years ago, light was slowed down to less than 10^{-7} of its vacuum speed in a cloud of ultracold atoms of sodium. Upon a sudden turn-off of the coupling laser, a slow light pulse can be imprinted on cold atoms such that it can be read out and converted into a photon again. In this process, the light is stopped by absorbing it and storing its shape within the atomic ensemble. Alternatively, the light can be stopped at the band edge in photonic-crystal waveguides, where the group speed vanishes. Here, we extend the phenomenon of stopped light to the new field of parity-time (PT) symmetric systems. We show that zero group speed in PT symmetric optical waveguides can be achieved if the system is prepared at an exceptional point, where two optical modes coalesce. This effect can be tuned for optical pulses in a wide range of frequencies and bandwidths, as we demonstrate in a system of coupled waveguides with gain and loss.

DOI: [10.1103/PhysRevLett.120.013901](https://doi.org/10.1103/PhysRevLett.120.013901)

Broken PT-symmetry phenomena




Fig 5. The probability density of the wave functions corresponding to the first and second excited states of the harmonic oscillator. States' probability densities are symmetric about zero. As the exceptional point parametric value is approached from the left, the probability densities begin to overlap in space. We can see the densities of the wave functions behave as mirror images of each other after we pass the exceptional point in the negative direction.

Constructing the \mathcal{PT} inner product

The \mathcal{PT} inner product is **not positive definite**.

To patch up this problem, we use the \mathcal{C} operator:

$\mathcal{C} \rightarrow$ charge conjugation

$$\mathcal{C}^2 = 1, \quad [\mathcal{C}, \hat{H}] = 0, \quad [\mathcal{C}, \mathcal{PT}] = 0.$$

We construct \mathcal{C} from the \hat{H} eigenstates ψ_n .

$$\mathcal{C} = \sum_n \psi_n(x) \psi_n(y)$$

Inner product:

$$(\mathcal{CPT}\psi_n(x)) \cdot \psi_m(y) = \mathcal{C}\psi_n^*(-x)) \cdot \psi_m(y)$$