



ERASMUS MUNDUS MASTER
IN ASTROPHYSICS AND SPACE SCIENCE

MASTER'S THESIS

RECONSTRUCTING PHOTOMETRIC LIGHTCURVES
OF ACTIVE GALACTIC NUCLEI USING NEURAL
STOCHASTIC FLOWS

SUPERVISORS

Prof. Anđelka Kovačević
Astrophysics Department, University of Belgrade

Prof. Médéric Boquien
Laboratoire Lagrange, Université Côte d'Azur

Prof. Dragana Ilić
Astrophysics Department, University of Belgrade

AUTHOR

Ana Cecilia Luis Ramirez

ACADEMIC YEAR 2025/2026



Cofunded by
the European Union

This Master thesis is submitted in partial fulfillment of the requirements for the degree of Master in Astrophysics and Space Science as part of a multiple degree awarded in the framework of the Erasmus Mundus Joint Master in Astrophysics and Space Science — MASS jointly delivered by a Consortium of four Universities: Tor Vergata University of Rome, University of Belgrade, University of Bremen, and Université Cote d'Azur, regulated by the MASS Consortium Agreement and funded by the EU under the call ERASMUS-EDU-2021-PEX-EMJM-MOB.

УНИВЕРЗИТЕТ У БЕОГРАДУ
МАТЕМАТИЧКИ ФАКУЛТЕТ



Ana Cecilia Luis Ramirez

Реконструкција фотометријских светлосних
кривих активних галактичких језгара
применом стохастичких Neural Flow модела

Master Thesis

Београд, 2026.

Contents

1	Background	6
1.1	Unified Model of AGN	8
1.1.1	Accretion Disk	10
1.1.2	Corona	11
1.1.3	Broad-line Region	11
1.1.4	Dusty Torus	12
1.1.5	Relativistic Jets	13
1.1.6	Narrow-line Region	14
2	AGN Variability Physics	15
2.1	Physical Origin and Timescales	15
2.1.1	Reverberation Mapping	16
2.2	Variability Characterization	18
3	Simulation Framework	20
3.1	The Vera C. Rubin Observatory LSST	20
3.2	Damped Random Walk	20
3.3	Multiband Light Curve Simulation	21
3.3.1	DRW Parameters and MacLeod Scaling Relations	21
3.3.2	EzTao Implementation	22
3.3.3	Noise Model	22
3.3.4	Sampling Strategy	22
3.4	Simulation Regime	22
4	Models	23
	Bibliography	24

List of Figures

1.1	AGN spectral energy distribution (SED) from radio frequencies to hard X-ray . .	7
1.2	Typical optical spectra of different AGN classes	8
1.3	NGC 1068 spectropolarimetry revealing the hidden broad line region (BLR) . . .	9
1.4	Schematic view of the the innermost regions of (non-jetted) AGN	10
1.5	Eddington ratio distribution of broad-line AGN at $1.0 < z < 2.2$	11
1.6	Schematic of the BLR formation via a failed disk wind	12
1.7	Radio contours of a giant radio galaxy extending to Mpc scales	13
1.8	SDSS galaxies in the classical BPT diagram.	14
2.1	Schematic of AGN reverberation mapping across different physical scales	16
2.2	Continuum and reverberation lags as a function of energy and Fourier frequency	17
2.3	X-ray light curves and PSDs for four AGN spanning four decades in black hole mass	18
3.1	Connection between exponential ACFs and structure functions for different stochastic models	21

List of Tables

2.1	AGN region size scales and variability timescales	16
-----	---	----

1 Background

Compared to ordinary galaxies, some have extraordinarily bright central regions, which can sometimes even outshine the host-galaxy emission by many orders of magnitude; these galaxies are called *active galaxies* and their nuclei are referred to as *active galactic nuclei* (AGN). The physical processes necessary to create such high luminosities ($L = 10^{43-48} \text{ erg s}^{-1}$; Lusso et al. 2012) within a small volume were a subject of debate for years, but now are widely accepted to be connected to accretion onto the galaxies' central black holes (BHs). These accreting objects have masses $M_{BH} \gtrsim 10^6 M_{\odot}$ (B. M. Peterson and Horne 2004) and an Eddington ratio exceeding $\lambda_{Edd}^1 \approx 10^{-5}$. This threshold, chosen somewhat arbitrarily, excludes galactic nuclei with no significant accretion activity, such as that of the Milky Way, while still covering the full range of observed AGN luminosities (Netzer 2015).

The “unified model” discussed in Section 1.1, places the BH at the center of a layered structure that places each component of the AGN at a characteristic distance (Figure 1.4). Each of these regions contributes emission into different wavelength regimes, resulting in these objects having detectable emission across the whole electromagnetic spectrum, as illustrated by the spectral energy distribution (SED) in Figure 1.1.

The gravitational energy released as the circumnuclear material falls inwards and loses angular momentum forms an *accretion disk* (§1.1.1) around the black hole, which acts as the primary source of AGN continuum emission. The disk gets heated up via viscous heating (B. M. Peterson 1993), producing a thermal spectrum that has a blackbody temperature peaking in the ultra violet (UV; $\approx 10 - 400 \text{ nm}$) for a “typical” AGN (i.e., with $M_{BH} \approx 10^8 M_{\odot}$; $\lambda_{Edd} \approx 0.1$); this is often referred to as the “big blue bump” (Harrison 2014; Figure 1.1)

The thermal disk photons, also called *seed photons*, are Comptonized by highly energetic electrons that are located at the *corona* (§1.1.2) to form the X-ray power-law spectrum observed in Figure 1.1. These high energy photons can also reflect off the accretion disk and/or the *torus* (§1.1.4), producing the additional “reflection” X-ray component. In a majority of AGN X-ray spectra, there is an additional “soft excess” at energies less than $\sim 2 \text{ keV}$. This emission is above what would be expected from simple accretion disk models and its origin is still a topic of discussion. Two proposed models (which could be simultaneously true) to explain this soft excess are: relativistically blurred reflection from the inner accretion disc, or Comptonization in a “warm corona” located near the surface of the accretion flow (e.g., Ballantyne and Xiang 2022).

¹The ratio L/L_{Edd} , where L is the bolometric luminosity and $L_{Edd} = 1.5 \times 10^{38} M_{BH}/M_{\odot} \text{ erg s}^{-1}$ is the Eddington luminosity for a solar composition gas. This is the maximum isotropic luminosity a body can achieve while maintaining balance between radiation pressure and gravitational force.




Figure 1.1: AGN SED from radio frequencies to hard X-ray, loosely based on the observed SEDs of radio-quiet quasars (e.g., Elvis et al. 1994; Richards et al. 2006). The black solid curve represents the total emission and the colored curves (shifted down for clarity) represent the individual components. The intrinsic shape of the SED in the mm-far infrared (FIR) regime is uncertain; however, it is credited to have a minimal contribution (to an overall galaxy SED) compared to star formation (SF), except in the most intrinsically luminous quasars and powerful jetted AGN. It should be noted that the relative contribution of each of the components can vary dramatically for different AGN types. Adapted from Harrison (2014).

The infrared (IR; $\approx 1 - 1000 \mu\text{m}$) band, is mostly sensitive to the reprocessed emission of the accretion disk by the torus. Although the shape of this IR emission varies from source to source, the peak of this dusty component is usually around $\lambda \approx 20 - 50 \mu\text{m}$ and falls steeply at longer wavelengths (Harrison 2014). In the sub-mm/far-infrared (FIR), or sometimes even in the mid-infrared (MIR), stellar heated emission from cooler dust dominates over AGN heated emission (Lutz 2014).

AGN cover a wide range of radio luminosities, which can differ by several orders of magnitude between radio-quiet (RQ, also called non-jetted) and radio-loud (RL, also called jetted) sources (orange lines in Figure 1.1). The latter type makes up only a minority of the AGN population (Padovani 2011) with objects that present highly collimated, relativistic radio *jets* (§1.1.5) and extended radio structures such as lobes and hotspots, that together emit strong non-thermal radiation and, occasionally, a γ -ray jet. On the other hand, the spectrum of non-jetted objects is dominated by thermal emission in most bands and the origin of their radio emission has not been clearly established. Some possibilities include coronal activity or synchrotron radiation from particles accelerated in shocks (see e.g., Laor and Behar 2008; Ishibashi and Courvoisier 2011).




Figure 1.2: Optical spectra of representative AGN classes, from the (essentially featureless) BL Lac objects (Top left) through the radio galaxies and QSOs (Seward and Charles 2010). Note the presence of both broad and narrow emission lines in quasars, Seyfert 1 galaxies and broad-line radio galaxies (BLRG), while Seyfert 2 galaxies, low-ionisation nuclear emission region (LINER) galaxies and narrow-line radio galaxies (NLRG) only show narrow permitted and forbidden lines. A normal galaxy spectrum (Bottom left) is shown for comparison.

1.1 Unified Model of AGN

Because different wavebands probe different physical components, AGN selected at particular wavelengths often exhibit distinct properties, leading to a rich phenomenological classification (Figure 1.2). Reality, however, is much simpler because now we know that these classes differ from each other only by a small number of parameters, such as: orientation, accretion rate, the presence of jets, and possibly the host galaxy and environment (see Padovani et al. (2017) and references there-in). This suggests that these numerous brands do not necessarily correspond to fundamentally different physical objects, but instead motivate a unifying framework capable of explaining their apparent diversity.

This unification idea arose shortly after the discovery of *quasars* (QSOs)², when the observed variety of AGN prompted questions about the role of the observer’s orientation relative to the AGN axis. Blandford and Rees (1978) proposed that BL Lac objects were radio galaxies viewed down the axis of a relativistic jet. Relativistic beaming (§1.1.5) would cause the non-thermal continuum to appear very bright and the emission lines weak in comparison. The same object, viewed from the side, would have normal emission-line equivalent widths, and the radio structure

²The term quasar historically stands for “quasi-stellar radio source”; throughout this thesis it is used more broadly to refer to very luminous AGN.




Figure 1.3: Optical spectra of the nucleus of NGC 1068 (J. S. Miller, Goodrich, and Mathews 1991). The total spectrum (Top) shows the narrow emission lines expected for Type 2 AGN, while the polarized flux spectrum (Bottom) reveals the broad permitted emission lines characteristic of unobscured Type 1 AGN.

would be dominated by the extended lobes rather than the core (Shields 2004).

Using spectropolarimetry, the measurement of the polarization state of the incoming light from a source as a function of wavelength, Antonucci and J. S. Miller (1985) found that the polarized flux of NGC 1068, a Seyfert 2 galaxy (hereafter Seyfert 2 or Type 2; characterized by narrow emission lines only), resembled that of a Seyfert 1 spectrum (hereafter Seyfert 1 or Type 1), which additionally exhibits broad permitted emission lines (Figure 1.3). This was interpreted as the broad-line region (BLR; §1.1.3) and the central continuum source being blocked from direct view by a dusty torus, while free electrons above the nucleus scatter a fraction of the nuclear light towards the observer, polarizing it in the process. The broad lines had gone unnoticed in direct light because this scattered component was too faint relative to the strong narrow-line emission from the narrow-line region (NLR; §1.1.6), which extends beyond the obscuring torus and is therefore visible regardless of orientation.

Multiple efforts such as those recounted above (Antonucci 1984; Orr and Browne 1982) were fundamental to the development of the “favoured” unified model of AGN (Antonucci 1993; Urry and Padovani 1995) introduced in this section. Whilst it has been successful in connecting Type I and II Seyferts, and Flat Spectrum Radio Quasars (FSRQs) and High-Excitation Galaxies (HEGs), recent work suggests that a picture based primarily on orientation and obscuring material is incomplete (Netzer 2015; Padovani et al. 2017). Nonetheless, the major aspects of the model are sufficient for the purposes of this thesis; in the sections below, the primary components are explained briefly. AGN structure, from the very center to host-galaxy scales, is mainly constituted by: a massive central BH, an X-ray corona, an accretion disk, a broad-line region, a torus, and




Figure 1.4: Overview of the innermost regions around a non-jetted accreting supermassive black hole (SMBH). Typical scales along the polar and equatorial directions (for nearby AGN) are shown. Different colors depict regions with different densities or compositions. Taken from Ramos Almeida and Claudio Ricci 2017.

a narrow-line region (Figure 1.4), with radio-loud AGN also hosting relativistic jets. Additional features such as polar dust and large-scale outflows (also visible in Figure 1.4) are discussed briefly in the context of the relevant components but are not treated as separate subsections here.

1.1.1 Accretion Disk

The standard framework of the optically thick, geometrically thin accretion disk (Shakura and Sunyaev 1973) described above applies to SMBHs with moderate to high Eddington ratios ($10^{-2} \lesssim \lambda_{Edd} \lesssim 1$; C. Ricci 2026), where the accretion flow is considered radiatively efficient. This scheme describes the disk as a composition of Keplerian rings, nested and coupled to each other by viscous forces, likely driven by turbulence and magnetic instabilities, the magneto-rotational instability (MRI; Balbus and Hawley 1998) being the most widely quoted. The outer, slower ring brakes the inner faster one, transporting the angular momentum outwards, allowing material to spiral inward. Friction between the rings produces heat that is dissipated as blackbody emission with a characteristic maximum temperature of $T \sim 10^5$ K for the “typical” AGN described above (Netzer 2013), yielding the multi-temperature spectrum depicted in Figure 1.1. The continuum emission of the accretion disk varies with time depending on the dynamics of the infalling matter (see Chapter 2).

Most QSOs in the local universe are thought to accrete through thin disks, yet contrasting accretion scenarios are possible depending on the Eddington ratio (Figure 1.5). At $\lambda_{Edd} \lesssim 10^{-2}$, the disk becomes optically thin and with an Advection Dominated Accretion Flow (ADAF, Narayan and Yi 1994), radiating weakly and with minimal optical/UV emission. Conversely, at $\lambda_{Edd} \gtrsim 1$, theoretical works have predicted that the SMBHs accrete rapidly through “slim” (or “thick”) disks, showing very different SEDs than their slower counterparts and powerful outflows (e.g., Kubota and Done 2018; Giustini and Proga 2019).




Figure 1.5: Eddington ratio distribution of broad-line AGNs at $1.0 < z < 2.2$. Different X-ray surveys are represented as different color histograms, and the black histogram pictures all the combined distributions. The gray shade illustrates the Eddington limit ($\lambda_{Edd} = 1$) and the red solid line corresponds to a log-normal fit with a peak of $\log L/L_{Edd} = -0.6$ (Suh et al. 2015).

1.1.2 Corona

Numerical simulations of magnetohydrodynamical accretion disks have shown that the dissipation of the magnetic energy built up by the MRI naturally produces a hot, active corona located within a few gravitational radii of the accretion disk (Balbus and Hawley 1998; K. A. Miller and Stone 2000; Fabian et al. 2009). This compact region of ionized gas reaches temperatures of $T \sim 10^9$ K (Kara and García 2025), and its rapid variability (see Chapter 2) is consistent with such a small emitting region. The corona contributes approximately 5% of the AGN bolometric output (C. Ricci 2026), a fraction that decreases with increasing λ_{Edd} , with lower λ_{Edd} systems showing stronger X-ray emission relative to the disk (Padovani et al. 2017).

1.1.3 Broad-line Region

The broad-line region consists of high density gas clouds predominantly in Keplerian motion at luminosity-dependent distances of $0.01 - 1$ pc from the central black hole (Netzer 2015), well within the *dust sublimation radius* (see §1.1.4). The BLR is therefore dust free, emitting broad permitted and semi-forbidden emission lines with velocity widths ranging from 1,000 to 20,000 km s^{-1} (C. Ricci 2026), wideness caused by *Doppler broadening*³. Ionizing photons from the accretion disk and corona illuminate these clouds, which subsequently exhibit ionization stratification with respect to radius, with higher ionization species located closer to the central source (Netzer 2013). The physical origin of the BLR remains an open question, some of the models presented in the literature include radiation pressure driven disk winds (Murray et al. 1995) and failed dust-driven wind from the disk surface (Figure 1.6; Czerny and Hryniewicz 2011). The distance of the

³Doppler broadening refers to the widening of spectral emission lines caused by the motions of gas clouds in the BLR. Clouds moving towards the observer have their emission blueshifted, while those moving away are redshifted; the superposition of all these shifted lines produces a single, broadened emission feature whose width reflects the velocity dispersion of the gas.




Figure 1.6: Schematic of the BLR geometry and its location relative to the accretion disk and dusty torus. The figure also illustrates the failed wind scenario of Czerny and Hryniewicz (2011): dust forms in the atmosphere of the inner accretion disk at radii where $T_{eff} < T_{sub}$, is launched as a wind, but then is suppressed by the radiation from the central source and partially falls back, giving rise to the inflowing and outflowing clouds that make up the BLR.

BLR from the central source scales with the AGN luminosity as $R_{BLR} \propto L^{1/2}$ (Kaspi et al. 2000), a relationship that is fundamental in reverberation mapping studies (§2.1.1).

1.1.4 Dusty Torus

As posed previously, the presence of an assymetric “dusty obscurer” has successfully unified apparently different AGN classes. When observed edge-on (i.e., at high inclination angles), this torus⁴ obscures the innermost regions of the AGN and hides the broad emission lines; conversely, when viewed face-on (at lower inclination angles), there is a clear line of sight to the BLR and accretion disk (Seyfert 1 vs Seyfert 2 in Figure 1.2). This structure is conceptualized as an optically thick structure surrounding the central engine at scales larger than that of the BLR and luminosity dependent dimensions ranging from 0.1 – 10 pc (Netzer 2015). Its inner boundary coincides is the so-called dust sublimation radius (R_{sub}), beyond which the dust can survive the radiation from the central engine. This radius scales with luminosity as $R_{sub} \propto L^{1/2}$, and corresponds to the distance at which the dust temperature reaches the sublimation threshold ($T_{sub} \sim 1400 - 1800$ K for silicate and graphite grains respectively; (Barvainis 1987; Netzer 2015)).

In addition to being responsible for the IR reprocessed emission on the AGN SED (Figure 1.1), the torus acts as a gas and dust reservoir that regulates the material available for accretion onto the black hole. Its geometry, believed to contain a central opening, also collimates the AGN radiation into bi-conical ionization cones along the polar direction (Pogge 1988), which

⁴For convenience, we keep referring to this obscuring matter as the torus despite the fact that it could have a different (and more complex) geometrical distribution (e.g., Nenkova et al. 2008; Stalevski et al. 2012).




Figure 1.7: SDSS optical image (composite from bands g , r and i) of the galaxy group UGC 9555 overlaid with NVSS contours (red) and LOFAR M8SS radio contours (white) revealing the size of the giant radio galaxy (GRG). The white bar indicates a physical scale of 1 Mpc, illustrating how AGN jets can extend well beyond the boundaries of their host galaxies. Adapted from Clarke et al. 2017.

illuminate the NLR (§1.1.6). AGN also drive large scale winds and outflows along these directions, powered by radiation and magnetic pressure from the central engine, which can interact with and regulate the host galaxy interstellar medium (C. Ricci 2026).

1.1.5 Relativistic Jets

Jets are thought to be powered by the extraction of rotational energy from the SMBH via strong electromagnetic fields (Blandford and Znajek 1977), and can extend to Mpc scales from their host galaxy (Figure 1.7; Clarke et al. 2017).

Their launch point coincides with the location of the corona, in the immediate vicinity of the accreting object, and their apparent velocities range from mildly to highly relativistic motion. The broadband emission of jets is characterised by a double-peaked SED: the low-energy peak, spanning radio to optical wavelengths, arises from synchrotron radiation of relativistic electrons spiralling in the jet magnetic field, while the high-energy peak, from X-ray to γ -ray energies, is thought to result from inverse-Compton scattering of soft photons from the BLR and accretion disk by the same relativistic electrons (Netzer 2013).

Their observed properties depend strongly on orientation because relativistic beaming could make an approaching jet appear significantly brighter than a receding one. This effect, known as *Doppler boosting*, arises when the jet plasma has bulk relativistic motion that concentrates the emitted radiation into a narrow cone in the direction of motion, amplifying the observed flux and variability for jets viewed close to face-on and explaining the extreme characteristics observed on blazars (Urry and Padovani 1995). Jets are also thought to play a role in galaxy evolution by injecting energy into the surrounding environment, a process referred to as radio-mode feedback (Fabian 2012).




Figure 1.8: Select galaxy objects from the Sloan Digital Sky Survey (SDSS) catalog shown in the BPT diagram. It plots the intensity of the [OIII] $\lambda 5007$ forbidden transition relative to that of the $H\beta$ line versus the ratio of [NII] $\lambda 6583$ to $H\alpha$ on logarithmic axes. Using this to infer ionization states and abundances of the emitting medium enables us to reconstruct the origin of the dominant radiation in the galactic host, which has led to various approaches for the classification of these objects. Depicted is the theoretical delimiter by Kewley et al. (2001) as well as the empirical line by Kauffmann et al. (2003) to distinguish predominantly starforming galaxies from AGN. Additionally, the Schawinski et al. (2007) boundary sorts AGN into efficiently accreting Seyfert and inefficient LINER subtypes. Courtesy of Fritz A. Agildere.

1.1.6 Narrow-line Region

This region is extended enough to be resolved optically, spanning hundreds or even thousands parsecs along the ionization cones (Figure 1.4). The NLR, just as the BLR, is photoionized by the central engine and exhibits optical emission lines, kinematically broadened with typical widths of $10^2\text{--}3 \text{ km s}^{-1}$ (Bennert 2005), with most lines falling within the $350\text{--}500 \text{ km s}^{-1}$ range (C. Ricci 2026).

Compared to the BLR, the NLR has a lower electron density, allowing forbidden transitions to occur without being collisionally suppressed. The presence of these forbidden lines, together with the high ionization state of the gas, makes the NLR emission diagnostically distinct from that of star-forming regions, a separation commonly visualised using emission line ratio diagrams such as the Baldwin-Phillips-Terlevich (BPT) diagram (Baldwin, Phillips, and Terlevich 1981) (Figure 1.8).

2 AGN Variability Physics

The first quasar discovered, 3C 273 (Edge et al. 1959), had been documented in photographic plates at least as early as 1887 (Slavcheva-Mihova et al. 2013). This emblematic radio-loud object showed variable emission in the optical and radio bands on timescales from days to years (Smith and Hoffleit 1963; Dent 1965), indicating that the size of the emitting region spanned only a few light-days in size. This compactness, coupled with the finding of its distant redshift (Schmidt 1963), was very difficult to explain by physical processes that operate in normal galaxies, and variability thus became one of the key arguments that the enormous luminosities of quasars had to originate from an exceptionally compact and energetic central engine (Salpeter 1964).

Now we know that variability in all wavelengths is an ubiquitous property of AGN, and so, its studies have been essential in understanding the physics of their nuclear regions, which, in general, cannot be resolved by current telescopes. This had been strictly true until the observations of Sgr A* and M87 (The Event Horizon Telescope Collaboration et al. 2019) and, more recently, with the GRAVITY interferometer on the VLT resolving the BLR in 3C 273 (Gravity Collaboration et al. 2018); still, such surveys are limited to massive sources in the very nearby universe. Variability experiments mostly can be categorised by the region being probed and the type of *reverberation signal* (§2.1.1) being measured, as illustrated in Figure 2.1. The primary observational product of such investigations is the *light curve*, a time series recording how the brightness of the AGN, measured within a given wavelength band, varies over a monitoring period. Moreover, the obtained time scales, spectral changes, and the correlations and delays between variations in different continuum or line components provide crucial information on the nature and location of these components and on their interdependencies (Ulrich et al. 1997).

2.1 Physical Origin and Timescales

Early campaigns quickly revealed that AGN vary much more rapidly in X-rays than in the optical (Winkler and White 1975), sometimes on timescales as short as a few minutes (McHardy, Lawrence, et al. 1981), and that variability at different wavelengths is correlated (Ulrich et al. 1997). This stochastic behaviour can generally be explained by the *propagating fluctuations* theory (Lyubarskii 1997). Local fluctuations in the disk’s surface density, presumably driven by the MRI (§ 1.1.1), occur at all radii and propagate inwards at the local viscous timescale¹. As these

¹The viscous timescale, $t_{\text{visc}} \sim r^2/\nu$, where ν is the kinematic viscosity, indicates the time for material to drift radially through the disk and increases with radius.




Figure 2.1: Cross-sectional schematic of an AGN illustrating the four main types of reverberation mapping and their corresponding radial scales from the central black hole. Taken from Cackett et al. 2021.

fluctuations travel, they modulate the local mass accretion rate at progressively smaller radii, coupling the variability of the outer disk to that of the hotter inner regions and corona, where the shorter local viscous timescale drives faster fluctuations in the emitted luminosity (Phil Uttley and Malzac 2023). The X-ray variability produced in the innermost regions then irradiates the outer accretion disk and is reprocessed, driving the optical and UV variability observed at longer wavelengths with a characteristic time delay (Padovani et al. 2017), as discussed below. Other mechanisms, including changes in line-of-sight obscuration, jet activity, or external perturbations like tidal disruption events (TDEs), can also contribute to the overall variability but are not described further here.

2.1.1 Reverberation Mapping

Reverberation mapping (RM) exploits the time delays between variability in different AGN components to infer their physical properties and spatial scales (Blandford and McKee 1982). The underlying principle is that any region illuminated by the central engine responds to its variability with a characteristic lag $\tau = R/c$, where R is the light travel distance to the responding region, providing a direct physical scale (B. Peterson 2006). The characteristic timescales and size scales probed by different types of reverberation are summarised in Table 2.1 and Figure 2.1.

Table 2.1: Order of magnitude size scales and their equivalent light-crossing times for the different regions of an AGN. Adapted from Cackett et al. 2021.

Region	Size scale (R_g)	Light-crossing time	
		$10^7 M_\odot$	$10^9 M_\odot$
X-ray corona	10	500 s	14 hr
UV/optical disk	10^2 – 10^4	0.06–6 days	6–600 days
BLR	10^3 – 10^5	0.6–60 days	60–6000 days
Dusty torus	$> 10^5$	> 60 days	> 6000 days

In the optical/UV, the most established application is BLR reverberation mapping, which measures the lag between continuum and broad emission line variations; since both the continuum and the broad lines are observed to vary in a correlated fashion, the measured lag directly yields the BLR size via $R_{\text{BLR}} = c\tau$ (B. Peterson 2006). As introduced in Section 1.1.3, this size




Figure 2.2: Frequency-resolved X-ray time lags as a function of energy for an AGN ($\sim 10^6 M_\odot$, left) and a black hole X-ray binary ($\sim 10 M_\odot$, right), illustrating the universality of the two lag regimes. At low Fourier frequencies (top, blue), hard X-rays lag soft ones, consistent with inward-propagating accretion fluctuations. At high Fourier frequencies (bottom, red), soft X-rays lag hard ones, consistent with disk reverberation. Taken from Kara and García 2025.

scales with AGN luminosity as $R_{\text{BLR}} \propto L^{1/2}$ (Kaspi et al. 2000), a relationship empirically calibrated through RM campaigns and used to estimate black hole masses via the virial theorem². Continuum reverberation mapping instead measures the wavelength-dependent lags between different optical/UV bands, probing the accretion disk size and temperature profile (Cackett et al. 2021).

Both propagating fluctuation lags and reverberation lags are simultaneously present in the data, but dominate at different Fourier frequencies and have opposite energy dependences (Figure 2.2): at low frequencies ($\nu \lesssim 10^{-5}$ Hz for a $10^6 M_\odot$ black hole, or equivalently timescales $\gtrsim 1$ day), hard X-ray variations lag soft ones as accretion fluctuations propagate inward; at high frequencies ($\nu \gtrsim 10^{-3}$ Hz, timescales of minutes), soft X-rays lag hard ones as the disk responds to the driving corona (Kara and García 2025). Simple cross-correlation averages over all timescales and can obscure this distinction, making Fourier analysis essential for disentangling the two contributions.

Observed continuum reverberation lags may be contaminated by diffuse BLR emission to the broadband continuum, which lies farther out in the accretion flow (e.g. Korista & Goad 2001; Cackett2018 ADD REFERENCE) and makes them appear longer than they really are. Variability-based methods are more effective than SED-based methods at identifying low mass and low luminosity AGN.

² $M_{\text{BH}} = f R_{\text{BLR}} \Delta V^2 / G$, where ΔV is the emission line width and f is a geometric factor (B. Peterson 2006).




Figure 2.3: Left: X-ray light curves of four Seyfert 1 AGN with black hole masses spanning 10^5 – $10^8 M_\odot$, observed with XMM-Newton over 70 ks. Smaller black holes vary more rapidly, with the $10^5 M_\odot$ AGN (NGC 4395, top) showing variability on timescales of tens of seconds, while the $10^8 M_\odot$ AGN (Ark 120, bottom) varies on timescales of hours. Right: The corresponding PSDs, showing the red-noise power-law shape characteristic of AGN X-ray variability. Taken from Kara and García 2025.

2.2 Variability Characterization

Variability has been studied under a variety of methods, starting from simple statistical quantities tracing the amplitude and timescales of the observed variability to more sophisticated approaches in both the time and frequency domains (Paolillo and Papadakis 2025). In general, these statistics attempt to measure the absolute or fractional variance of the light curve over some characteristic timescale Δt , or express the degree of correlation between different timescales or photometric bands (Ulrich et al. 1997; De Cicco et al. 2025).

The distribution of variability power over temporal frequencies (or, equivalently, $1/\text{timescale}$) is quantified by the *power spectral density* (PSD), defined as the squared amplitude of the flux (e.g., P. Uttley et al. 2002). AGN X-ray PSDs typically follow a “red-noise” power law, $P(\nu) \propto \nu^{-\alpha}$ with $\alpha \sim 1$ at low frequencies, steepening to $\alpha \gtrsim 2^3$ above a characteristic break frequency ν_b , consequently displaying larger amplitude variations on longer timescales. This break timescale scales with black hole mass and accretion rate (McHardy, Koerding, et al. 2006), with smaller black holes varying more rapidly than larger ones for the same observing duration (Figure 2.3). The anticorrelation can be interpreted as a consequence of the fact that the most luminous AGN are likely to host larger mass black holes (if all quasars accrete at similar Eddington rates), in which case, their variability should be diluted by the larger emitting region (Paolillo and Papadakis 2025).

The PSD also depends strongly on the observing waveband, with optical PSDs showing steeper slopes and lower normalization than X-ray PSDs, consistent with the propagating fluctuations

³ $\alpha \sim 2$ corresponds to a *Damped random walk* that breaks to flat slope at long timescales.

picture where outer disk regions contribute power at lower frequencies and inner regions at higher ones (Kara and García 2025).

A complementary time-domain characterization is provided by the *structure function* (SF), which quantifies the mean variability amplitude as a function of time lag Δt ; on short timescales the SF rises as a power law, flattening above a characteristic damping timescale τ that encodes the same physical information as the PSD break frequency (Kozłowski 2016).

Deriving the PSD/SF features and their uncertainties from an irregularly sampled light curve is very challenging, specially for stochastic light curves such as those of AGN; thereon, high quality light curves are indispensable to enable robust characterization of the UV/optical variability and interband lags.

Beyond the persistent stochastic variability described above, more extreme variability phenomena also exist, including *changing-look* AGN, which undergo dramatic changes in their broad emission lines and continuum on timescales of months to years, possibly driven by rapid changes in accretion rate or obscuration (LaMassa et al. 2015), as well as quasi-periodic eruptions (QPEs) and tidal disruption events (TDEs), they represent transient phenomena superimposed on the underlying stochastic variability.

3 Simulation Framework

This chapter describes the framework used to simulate realistic AGN multiband light curves in the LSST *ugriz* bands. The simulation is based on the damped random walk (DRW) model, whose parameters are assigned using empirical wavelength-dependent scaling relations, and implemented using the EZTAO Gaussian process library (Yu and Richards 2022).

3.1 The Vera C. Rubin Observatory LSST

3.2 Damped Random Walk

The simplest stochastic model consistent with the observed AGN PSD properties is the *damped random walk* (DRW), also known as a first-order continuous autoregressive process, CAR(1), or Ornstein-Uhlenbeck process (Kelly, Bechtold, and Siemiginowska 2009). It is described by the stochastic differential equation:

$$df(t) = -\frac{f(t)}{\tau_{\text{DRW}}} dt + \sigma_{\text{DRW}} \sqrt{dt} \epsilon(t), \quad (3.1)$$

where τ_{DRW} is the damping (or characteristic) timescale, σ_{DRW} is the asymptotic variability amplitude, and $\epsilon(t)$ is a white noise process. The DRW has an exponential autocorrelation function:

$$\text{ACF}(\Delta t) = \exp\left(-\frac{|\Delta t|}{\tau_{\text{DRW}}}\right), \quad (3.2)$$

corresponding to a PSD of the form $P(\nu) \propto \nu^{-2}$ at high frequencies, flattening to white noise below the break frequency $\nu_b = 1/(2\pi\tau_{\text{DRW}})$. In the structure function representation, the DRW produces a power-law rise with slope $\gamma = 0.5$ on short timescales ($\Delta t \ll \tau_{\text{DRW}}$), flattening to a constant value set by σ_{DRW} above the characteristic timescale (Figure 3.1; Kozłowski 2016). The DRW has been shown to provide a good statistical description of optical AGN light curves (Kelly, Bechtold, and Siemiginowska 2009; Kozłowski et al. 2010; MacLeod et al. 2010), with τ_{DRW} found to correlate with black hole mass and luminosity, consistent with the physical picture developed in Chapter 2.

The DRW is the simplest member of the broader family of continuous autoregressive moving average (CARMA) models (Kelly, Becker, et al. 2014), corresponding to the special case




Figure 3.1: Left: Exponential autocorrelation functions $ACF(\Delta t) = \exp(-(|\Delta t|/\tau)^\beta)$ for different values of β . Right: The corresponding structure functions. The DRW model ($\beta = 1$, red) produces a SF with slope $\gamma = 0.5$ on timescales $\Delta t \ll \tau$, flattening above the characteristic damping timescale. Adapted from Kozłowski 2016.

CARMA(1, 0). More flexible CARMA(p, q) processes can reproduce a wider range of PSD shapes; in particular, the CARMA(2, 1) process, also known as the damped harmonic oscillator (DHO), has been shown to provide superior fits to observed AGN light curves compared to the DRW (YuEzTao2025). The simulation framework described here assumes a DRW process for the underlying variability, though the implementation supports more general CARMA kernels.

3.3 Multiband Light Curve Simulation

3.3.1 DRW Parameters and MacLeod Scaling Relations

The DRW parameters τ_{DRW} and σ_{DRW} depend on the observing wavelength, black hole mass, and luminosity. Following MacLeod2010DRW, the wavelength dependence can be expressed as power-law scaling relations:

$$\tau_{\text{DRW}} \propto \lambda^{0.17}, \quad \sigma_{\text{DRW}} \propto \lambda^{-0.479}, \quad (3.3)$$

where λ is the effective wavelength of the observing filter. These relations imply that redder bands have slightly longer damping timescales and lower variability amplitudes than bluer bands, consistent with the wavelength dependence of AGN variability discussed in Chapter 2.

In the simulation framework, a *driver band* is defined as the reference band from which the DRW parameters of all other bands are derived via Equation 3.3. The driver band parameters ($\tau_{\text{DRW}}^g = 100$ days, $\sigma_{\text{DRW}}^g = 0.112$ mag) are taken as fiducial values representative of typical quasar optical variability (YuEzTao2025), with the g band chosen as the reference. The parameter space explored in this work follows YuEzTao2025:

$$\tau_{\text{DRW}} \in [0.01, 500] \text{ days} \quad (\text{log-uniform}), \quad \sigma_{\text{DRW}} \in [0.02, 0.2] \text{ mag} \quad (\text{log-uniform}). \quad (3.4)$$

3.3.2 EzTao Implementation

3.3.3 Noise Model

Observational noise is simulated by adding Gaussian random values drawn from $\mathcal{N}(0, \sigma_{\text{LSST}}^2)$ to each flux measurement, where $\sigma_{\text{LSST}} = 0.01$ mag is the expected LSST single-visit photometric uncertainty¹.

3.3.4 Sampling Strategy

The simulation supports two modes: a single-band driver light curve (**makeDriver**), which generates a single DRW realisation at the reference band parameters on a uniform time grid with step Δt up to a maximum baseline t_{max} ; and a multiband mode (**makeMultiband**), which generates correlated light curves in all requested bands simultaneously using the wavelength-scaled parameters. A sparse sampling mode (**makeSparse**) is also implemented to mimic realistic LSST observing cadences by subsampling the uniform grid with gaps and irregular spacing.

3.4 Simulation Regime

¹<https://rubinobservatory.org/for-scientists/rubin-101/key-numbers>

4 Models

Bibliography

- Antonucci, Robert (Mar. 1984). “Optical spectropolarimetry of radio galaxies.” In: *The Astrophysical Journal* 278, pp. 499–520. DOI: 10.1086/161816.
- (Jan. 1993). “Unified models for active galactic nuclei and quasars.” In: *Annual Review of Astronomy and Astrophysics* 31, pp. 473–521. DOI: 10.1146/annurev.aa.31.090193.002353.
- Antonucci, Robert and J. S. Miller (Oct. 1985). “Spectropolarimetry and the nature of NGC 1068.” In: *The Astrophysical Journal* 297, pp. 621–632. DOI: 10.1086/163559.
- Balbus, Steven A. and John F. Hawley (Jan. 1998). “Instability, turbulence, and enhanced transport in accretion disks”. In: *Reviews of Modern Physics* 70.1, pp. 1–53. DOI: 10.1103/RevModPhys.70.1.
- Baldwin, J. A., M. M. Phillips, and R. Terlevich (Feb. 1981). “Classification parameters for the emission-line spectra of extragalactic objects.” In: *Publications of the Astronomical Society of the Pacific* 93, pp. 5–19. DOI: 10.1086/130766.
- Ballantyne, David and Xin Xiang (Apr. 2022). “A Model for the X-ray Soft Excess in Active Galactic Nuclei: Combining the Effects of Reflection and a Warm Corona”. In: *AAS/High Energy Astrophysics Division*. Vol. 19. AAS/High Energy Astrophysics Division, 106.39, p. 106.39.
- Barvainis, Richard (Sept. 1987). “Hot Dust and the Near-Infrared Bump in the Continuum Spectra of Quasars and Active Galactic Nuclei”. In: *The Astrophysical Journal* 320, p. 537. DOI: 10.1086/165571.
- Bennert, Nicola (2005). “The narrow-line region of active galaxies”. doctoralthesis. Ruhr-Universität Bochum, Universitätsbibliothek.
- Blandford, R. D. and C. F. McKee (Apr. 1982). “Reverberation mapping of the emission line regions of Seyfert galaxies and quasars.” In: *The Astrophysical Journal* 255, pp. 419–439. DOI: 10.1086/159843.
- Blandford, R. D. and M. J. Rees (Jan. 1978). “Some comments on radiation mechanisms in Lacer-tids.” In: *BL Lac Objects*. Ed. by A. M. Wolfe, pp. 328–341.
- Blandford, R. D. and R. L. Znajek (May 1977). “Electromagnetic extraction of energy from Kerr black holes.” In: *Monthly Notices of the Royal Astronomical Society* 179, pp. 433–456. DOI: 10.1093/mnras/179.3.433.
- Cackett, Edward M. et al. (June 2021). “Reverberation mapping of active galactic nuclei: From X-ray corona to dusty torus”. In: *iScience* 24.6, p. 102557. ISSN: 2589-0042. DOI: 10.1016/j.isci.2021.102557. URL: <http://dx.doi.org/10.1016/j.isci.2021.102557>.

- Clarke, A. O. et al. (May 2017). “LOFAR MSSS: Discovery of a 2.56 Mpc giant radio galaxy associated with a disturbed galaxy group”. In: *Astronomy and Astrophysics* 601, A25, A25. DOI: 10.1051/0004-6361/201630152. arXiv: 1702.01571 [astro-ph.GA].
- Czerny, B. and K. Hryniewicz (Jan. 2011). “The origin of the broad line region in active galactic nuclei”. In: *Astronomy and Astrophysics* 525, L8, p. L8. DOI: 10.1051/0004-6361/201016025. arXiv: 1010.6201 [astro-ph.CO].
- De Cicco, D. et al. (May 2025). “Selection of optically variable active galactic nuclei via a random forest algorithm”. In: *Astronomy and Astrophysics* 697, A204, A204. DOI: 10.1051/0004-6361/202453630. arXiv: 2505.15819 [astro-ph.GA].
- Dent, W. A. (Nov. 1965). “Variation in the Radio Emission of 3C273 and Other Quasi-Stellar Sources.” In: *Astronomical Journal* 70, p. 672. DOI: 10.1086/109792.
- Edge, D. O. et al. (Jan. 1959). “A survey of radio sources at a frequency of 159 Mc/s.” In: *Memoirs of the Royal Astronomical Society* 68, pp. 37–60.
- Elvis, Martin et al. (Nov. 1994). “Atlas of Quasar Energy Distributions”. In: *The Astrophysical Journal Supplement Series* 95, p. 1. DOI: 10.1086/192093.
- Fabian, A. C. (Sept. 2012). “Observational Evidence of Active Galactic Nuclei Feedback”. In: *Annual Review of Astronomy and Astrophysics* 50, pp. 455–489. DOI: 10.1146/annurev-astro-081811-125521. arXiv: 1204.4114 [astro-ph.CO].
- Fabian, A. C. et al. (2009). “Broad line emission from iron K- and L-shell transitions in the active galaxy 1H 0707-495”. In: *Nature* 459.7246, pp. 540–542. DOI: 10.1038/nature08007. URL: <https://doi.org/10.1038/nature08007>.
- Giustini, Margherita and Daniel Proga (2019). “A global view of the inner accretion and ejection flow around super massive black holes: Radiation-driven accretion disk winds in a physical context”. In: *Astronomy and Astrophysics* 630, A94. ISSN: 1432-0746. URL: <http://dx.doi.org/10.1051/0004-6361/201833810>.
- Gravity Collaboration et al. (Nov. 2018). “Spatially resolved rotation of the broad-line region of a quasar at sub-parsec scale”. In: *Nature* 563.7733, pp. 657–660. DOI: 10.1038/s41586-018-0731-9. arXiv: 1811.11195 [astro-ph.GA].
- Harrison, Christopher (2014). “Observational constraints on the influence of active galactic nuclei on the evolution of galaxies”. PhD Thesis. Durham University, UK.
- Ishibashi, W. and T. J.-L. Courvoisier (Jan. 2011). “Synchrotron radio emission in radio-quiet AGNs”. In: *Astronomy and Astrophysics* 525, A118, A118. DOI: 10.1051/0004-6361/201014987. arXiv: 1010.5591 [astro-ph.HE].
- Kara, Erin and Javier García (2025). *Supermassive Black Holes in X-rays: From Standard Accretion to Extreme Transients*. arXiv: 2503.22791 [astro-ph.HE]. URL: <https://arxiv.org/abs/2503.22791>.
- Kaspi, Shai et al. (Apr. 2000). “Reverberation Measurements for 17 Quasars and the Size-Mass-Luminosity Relations in Active Galactic Nuclei”. In: *The Astrophysical Journal* 533.2, pp. 631–649. DOI: 10.1086/308704. arXiv: astro-ph/9911476 [astro-ph].
- Kauffmann, Guinevere et al. (Dec. 2003). “The host galaxies of active galactic nuclei”. In: *Monthly Notices of the Royal Astronomical Society* 346.4, pp. 1055–1077. DOI: 10.1111/j.1365-2966.2003.07154.x. arXiv: astro-ph/0304239 [astro-ph].
- Kelly, Brandon C., Jill Bechtold, and Aneta Siemiginowska (May 2009). “ARE THE VARIATIONS IN QUASAR OPTICAL FLUX DRIVEN BY THERMAL FLUCTUATIONS?” In: *The Astrophysical Journal* 698.1, p. 895. DOI: 10.1088/0004-637X/698/1/895. URL: <https://doi.org/10.1088/0004-637X/698/1/895>.

- Kelly, Brandon C., Andrew C. Becker, et al. (2014). “FLEXIBLE AND SCALABLE METHODS FOR QUANTIFYING STOCHASTIC VARIABILITY IN THE ERA OF MASSIVE TIME-DOMAIN ASTRONOMICAL DATA SETS”. In: *The Astrophysical Journal* 788.1. ISSN: 1538-4357. DOI: 10.1088/0004-637x/788/1/33. URL: <http://dx.doi.org/10.1088/0004-637x/788/1/33>.
- Kewley, L. J. et al. (July 2001). “Theoretical Modeling of Starburst Galaxies”. In: *The Astrophysical Journal* 556.1, pp. 121–140. DOI: 10.1086/321545. arXiv: astro-ph/0106324 [astro-ph].
- Kozłowski, Szymon (Aug. 2016). “Revisiting Stochastic Variability of AGNs with Structure Functions”. In: *The Astrophysical Journal* 826.2, 118, p. 118. DOI: 10.3847/0004-637x/826/2/118. arXiv: 1604.05858 [astro-ph.GA].
- Kozłowski, Szymon et al. (Jan. 2010). “Quantifying Quasar Variability as Part of a General Approach to Classifying Continuously Varying Sources”. In: *The Astrophysical Journal* 708.2, pp. 927–945. DOI: 10.1088/0004-637x/708/2/927. arXiv: 0909.1326 [astro-ph.CO].
- Kubota, Aya and Chris Done (Oct. 2018). “A physical model of the broad-band continuum of AGN and its implications for the UV/X relation and optical variability”. In: *Monthly Notices of the Royal Astronomical Society* 480.1, pp. 1247–1262. DOI: 10.1093/mnras/sty1890. arXiv: 1804.00171 [astro-ph.HE].
- LaMassa, Stephanie M. et al. (Feb. 2015). “THE DISCOVERY OF THE FIRST “CHANGING LOOK” QUASAR: NEW INSIGHTS INTO THE PHYSICS AND PHENOMENOLOGY OF ACTIVE GALACTIC NUCLEI”. In: *The Astrophysical Journal* 800.2, p. 144. DOI: 10.1088/0004-637x/800/2/144. URL: <https://doi.org/10.1088/0004-637x/800/2/144>.
- Laor, Ari and Ehud Behar (Oct. 2008). “On the origin of radio emission in radio-quiet quasars”. In: *Monthly Notices of the Royal Astronomical Society* 390.2, pp. 847–862. ISSN: 1365-2966. DOI: 10.1111/j.1365-2966.2008.13806.x. URL: <http://dx.doi.org/10.1111/j.1365-2966.2008.13806.x>.
- Lusso, E. et al. (Sept. 2012). “Bolometric luminosities and Eddington ratios of X-ray selected active galactic nuclei in the XMM-COSMOS survey”. In: *Monthly Notices of the Royal Astronomical Society* 425.1, pp. 623–640. DOI: 10.1111/j.1365-2966.2012.21513.x. arXiv: 1206.2642 [astro-ph.CO].
- Lutz, Dieter (Aug. 2014). “Far-Infrared Surveys of Galaxy Evolution”. In: *Annual Review of Astronomy and Astrophysics* 52, pp. 373–414. DOI: 10.1146/annurev-astro-081913-035953. arXiv: 1403.3334 [astro-ph.GA].
- Lyubarskii, Yu. E. (Dec. 1997). “Flicker noise in accretion discs”. In: *Monthly Notices of the Royal Astronomical Society* 292.3, pp. 679–685. DOI: 10.1093/mnras/292.3.679.
- MacLeod, C. L. et al. (Oct. 2010). “Modeling the Time Variability of SDSS Stripe 82 Quasars as a Damped Random Walk”. In: *The Astrophysical Journal* 721.2, pp. 1014–1033. DOI: 10.1088/0004-637x/721/2/1014. arXiv: 1004.0276 [astro-ph.CO].
- McHardy, I. M., E. Koerding, et al. (Dec. 2006). “Active galactic nuclei as scaled-up Galactic black holes”. In: *Nature* 444.7120, pp. 730–732. DOI: 10.1038/nature05389. arXiv: astro-ph/0612273 [astro-ph].
- McHardy, I. M., A. Lawrence, et al. (Dec. 1981). “The Ariel V (3A) catalogue of X-ray sources. II.” In: *Monthly Notices of the Royal Astronomical Society* 197, pp. 893–919. DOI: 10.1093/mnras/197.4.893.
- Miller, J. S., R. W. Goodrich, and William G. Mathews (Sept. 1991). “Multidirectional Views of the Active Nucleus of NGC 1068”. In: *The Astrophysical Journal* 378, p. 47. DOI: 10.1086/170406.

- Miller, Kristen A. and James M. Stone (May 2000). “The Formation and Structure of a Strongly Magnetized Corona above a Weakly Magnetized Accretion Disk”. In: *The Astrophysical Journal* 534.1, p. 398. DOI: 10.1086/308736. URL: <https://doi.org/10.1086/308736>.
- Murray, N. et al. (Oct. 1995). “Accretion Disk Winds from Active Galactic Nuclei”. In: *The Astrophysical Journal* 451, p. 498. DOI: 10.1086/176238.
- Narayan, Ramesh and Insu Yi (June 1994). “Advection-dominated Accretion: A Self-similar Solution”. In: *The Astrophysical Journal Letters* 428, p. L13. DOI: 10.1086/187381. arXiv: [astro-ph/9403052](https://arxiv.org/abs/astro-ph/9403052) [astro-ph].
- Nenkova, Maia et al. (Sept. 2008). “AGN Dusty Tori. I. Handling of Clumpy Media”. In: *The Astrophysical Journal* 685.1, pp. 147–159. DOI: 10.1086/590482. arXiv: 0806.0511 [astro-ph].
- Netzer, Hagai (2013). *The Physics and Evolution of Active Galactic Nuclei*.
- (Aug. 2015). “Revisiting the Unified Model of Active Galactic Nuclei”. In: *Annual Review of Astronomy and Astrophysics* 53, pp. 365–408. DOI: 10.1146/annurev-astro-082214-122302. arXiv: 1505.00811 [astro-ph.GA].
- Orr, M. J. L. and I. W. A. Browne (Sept. 1982). “Relativistic beaming and quasar statistics.” In: *Monthly Notices of the Royal Astronomical Society* 200, pp. 1067–1080. DOI: 10.1093/mnras/200.4.1067.
- Padovani, P. (Mar. 2011). “The microjansky and nanojansky radio sky: source population and multiwavelength properties”. In: *Monthly Notices of the Royal Astronomical Society* 411.3, pp. 1547–1561. DOI: 10.1111/j.1365-2966.2010.17789.x. arXiv: 1009.6116 [astro-ph.CO].
- Padovani, P. et al. (Aug. 2017). “Active galactic nuclei: what’s in a name?” In: *Astronomy & Astrophysics Review* 25.1, 2, p. 2. DOI: 10.1007/s00159-017-0102-9. arXiv: 1707.07134 [astro-ph.GA].
- Paolillo, Maurizio and Iossif Papadakis (Aug. 2025). “Continuum optical-UV and X-ray variability of AGN: current results and future challenges”. In: *La Rivista del Nuovo Cimento* 48.8, pp. 537–621. ISSN: 1826-9850. DOI: 10.1007/s40766-025-00072-5. URL: <http://dx.doi.org/10.1007/s40766-025-00072-5>.
- Peterson, B. (Jan. 2006). “The Broad-Line Region in Active Galactic Nuclei”. In: *Lecture Notes in Physics* 693. DOI: 10.1007/3-540-34621-X_3.
- Peterson, B. M. (Mar. 1993). “Reverberation Mapping of Active Galactic Nuclei”. In: *Publications of the Astronomical Society of the Pacific* 105, p. 247. DOI: 10.1086/133140.
- Peterson, B. M. and K. Horne (Mar. 2004). “Echo mapping of active galactic nuclei”. In: *Astronomische Nachrichten* 325.3, pp. 248–251. DOI: 10.1002/asna.200310207. arXiv: [astro-ph/0407538](https://arxiv.org/abs/astro-ph/0407538) [astro-ph].
- Pogge, Richard W. (May 1988). “An Extended Ionizing Radiation Cone from the Nucleus of the Seyfert 2 Galaxy NGC 1068”. In: *The Astrophysical Journal* 328, p. 519. DOI: 10.1086/166309.
- Ramos Almeida, Cristina and Claudio Ricci (Oct. 2017). “Nuclear obscuration in active galactic nuclei”. In: *Nature Astronomy* 1, pp. 679–689. DOI: 10.1038/s41550-017-0232-z. arXiv: 1709.00019 [astro-ph.GA].
- Ricci, C. (2026). *Unification models of Active Galactic Nuclei*. arXiv: 2606.12511 [astro-ph.GA]. URL: <https://arxiv.org/abs/2606.12511>.
- Richards, Gordon T. et al. (Oct. 2006). “Spectral Energy Distributions and Multiwavelength Selection of Type 1 Quasars”. In: *The Astrophysical Journal Supplement Series* 166.2, pp. 470–497. DOI: 10.1086/506525. arXiv: [astro-ph/0601558](https://arxiv.org/abs/astro-ph/0601558) [astro-ph].
- Salpeter, E. E. (Aug. 1964). “Accretion of Interstellar Matter by Massive Objects.” In: *The Astrophysical Journal* 140, pp. 796–800. DOI: 10.1086/147973.

- Schawinski, Kevin et al. (Dec. 2007). “Observational evidence for AGN feedback in early-type galaxies”. In: *Monthly Notices of the Royal Astronomical Society* 382.4, pp. 1415–1431. DOI: 10.1111/j.1365-2966.2007.12487.x. arXiv: 0709.3015 [astro-ph].
- Schmidt, M. (Mar. 1963). “3C 273 : A Star-Like Object with Large Red-Shift”. In: *Nature* 197.4872, p. 1040. DOI: 10.1038/1971040a0.
- Seward, Frederick D. and Philip A. Charles (2010). *Exploring the X-Ray Universe*. 2nd. Chap. Active Galactic Nuclei, pp. 277–305.
- Shakura, N. I. and R. A. Sunyaev (Jan. 1973). “Black holes in binary systems. Observational appearance.” In: *Astronomy and Astrophysics* 24, pp. 337–355.
- Shields, Gregory A. (2004). *A Brief History of Active Galactic Nuclei*. <https://ned.ipac.caltech.edu/level5/Sept04/Shields/frames.html>. Accessed: 2026-05-28.
- Slavcheva-Mihova, L. et al. (2013). *3C 273 - half a century later*. arXiv: 1401.5987 [astro-ph.HE]. URL: <https://arxiv.org/abs/1401.5987>.
- Smith, Harlan J. and Dorrit Hoffleit (May 1963). “Light Variations in the Superluminous Radio Galaxy 3C273”. In: *Nature* 198.4881, pp. 650–651. DOI: 10.1038/198650a0.
- Stalevski, Marko et al. (Mar. 2012). “3D radiative transfer modelling of the dusty tori around active galactic nuclei as a clumpy two-phase medium”. In: *Monthly Notices of the Royal Astronomical Society* 420.4, pp. 2756–2772. ISSN: 0035-8711. DOI: 10.1111/j.1365-2966.2011.19775.x. eprint: <https://academic.oup.com/mnras/article-pdf/420/4/2756/2926101/mnras0420-2756.pdf>. URL: <https://doi.org/10.1111/j.1365-2966.2011.19775.x>.
- Suh, Hyewon et al. (Dec. 2015). “Eddington Ratio Distribution of X-ray-selected Broad-line AGNs at $1.0 < z < 2.2$ ”. In: *The Astrophysical Journal* 815.2, p. 129. DOI: 10.1088/0004-637X/815/2/129. URL: <https://doi.org/10.1088/0004-637X/815/2/129>.
- The Event Horizon Telescope Collaboration et al. (Apr. 2019). “First M87 Event Horizon Telescope Results. IV. Imaging the Central Supermassive Black Hole”. In: *The Astrophysical Journal Letters* 875.1, p. L4. DOI: 10.3847/2041-8213/ab0e85. URL: <https://doi.org/10.3847/2041-8213/ab0e85>.
- Ulrich, Marie-Helene et al. (1997). “Variability of active galactic nuclei”. In: *Ann. Rev. Astron. Astrophys.* 35, pp. 445–502. DOI: 10.1146/annurev.astro.35.1.445.
- Urry, C. Megan and P. Padovani (Sept. 1995). “Unified Schemes for Radio-Loud Active Galactic Nuclei”. In: *Publications of the Astronomical Society of the Pacific* 107, p. 803. DOI: 10.1086/133630. arXiv: astro-ph/9506063 [astro-ph].
- Uttley, P. et al. (May 2002). “Measuring the broad-band power spectra of active galactic nuclei with RXTE”. In: *Monthly Notices of the Royal Astronomical Society* 332.1, pp. 231–250. DOI: 10.1046/j.1365-8711.2002.05298.x. arXiv: astro-ph/0201134 [astro-ph].
- Uttley, Phil and Julien Malzac (2023). *Large and complex X-ray time lags from black hole accretion disks with compact inner coronae*. arXiv: 2312.08302 [astro-ph.HE]. URL: <https://arxiv.org/abs/2312.08302>.
- Winkler Jr., P. F. and A. E. White (Aug. 1975). “A sudden increase in the X-ray flux from Centaurus A.” In: *The Astrophysical Journal Letters* 199, pp. L139–L142. DOI: 10.1086/181868.
- Yu, Weixiang and Gordon T. Richards (Jan. 2022). *EzTao: Easier CARMA Modeling*. Astrophysics Source Code Library, record ascl:2201.001. ascl: 2201.001.