
BIVARIATE ANALYSIS OF Prov AND LOG_FD806 FOR ANA.MODEL2 RELATION BETWEEN LOG_FD806 AND Prov

The MEANS Procedure

	Analysis Variable : LOG_FD806													
Geography	N Obs	N	N Miss	Minimum	Lower Quartile	Median	Mean	Upper Quartile	Maximum	Quartile Range	Coeff of Variation	Lower 95% CL for Mean	Upper 95% CL for Mean	Skewness
Alberta	958697	958697	0	2.47	4.90	5.60	5.60	6.44	7.32	1.54	17.11	5.60	5.60	-0.29
Atlantic provinces	629768	629768	0	3.07	4.97	5.74	5.62	6.35	7.35	1.38	16.90	5.61	5.62	-0.41
British Columbia	969907	969907	0	3.22	4.97	5.55	5.49	6.10	7.36	1.14	16.17	5.49	5.49	-0.29
Manitoba	299857	299857	0	2.71	4.76	5.45	5.39	6.07	7.34	1.32	17.20	5.39	5.40	-0.23
Ontario	2882375	2882375	0	2.64	4.62	5.27	5.30	5.95	7.29	1.33	17.62	5.30	5.30	0.02
Quebec	2104687	2104687	0	3.02	4.52	5.29	5.24	5.96	7.25	1.44	17.79	5.24	5.24	-0.20
Saskatchewan	271874	271874	0	3.08	5.01	5.66	5.64	6.33	7.26	1.31	15.62	5.64	5.64	-0.29
Territorial capitals	11711	11711	0	3.57	4.99	5.76	5.71	6.45	7.36	1.47	16.60	5.70	5.73	-0.28



One-way ANOVA Assumptions

In order to run a one-way ANOVA the following assumptions must be met:

1. The response of interest is continuous and normally distributed for each treatment group:

Normality test: PROC UNIVARIATE NORMAL and QQPlot for each group.

- 2.Treatment groups are independent of one another. Experimental units only receive one treatment, and they do not overlap.
- 3. There are no major outliers.
- 4.A check for unequal variances will help determine which version of a one-way ANOVA is most appropriate

(Levene's test, Null hypothesis: variances are equal between groups):

- A .If variances are equal, then the assumptions of a standard one-way ANOVA are met.
- B. If variances are unequal, then a Welch's one-way ANOVA is appropriate.

Normal Distribution?

Null hypothesis: sample has a normal distribution

CLT:

a.If it looks normal and each group have more than 30 observations

b.lf moderately skewed, each group must have more than 100 observations

*rule of thumb: If skewness is between -1 and -0.5 or between 0.5 and 1, the distribution is moderately skewed.

*if the sample size is over 2000, the Kolmgorov test should be used. If the sample size is less than 2000, the Shapiro test is better.

The UNIVARIATE Procedure Variable: LOG_FD806

Freq: WeightD

Geography=Alberta

Moments					
N	958697	Sum Weights	958697		
Mean	5.59782167	Sum Observations	5366614.84		
Std Deviation	0.95784837	Variance	0.91747349		
Skewness	-0.2933758	Kurtosis	-0.4036799		
Uncorrected SS	30920931	Corrected SS	879578.166		
Coeff Variation	17.1110911	Std Error Mean	0.00097826		

Basic Statistical Measures					
Location		Variability			
Mean	5.597822	Std Deviation	0.95785		
Median	5.595083	Variance	0.91747		
Mode	6.439478	Range	4.85031		
		Interquartile Range	1.54433		

Freq: WeightD

Geography=Alberta

Tests for Location: Mu0=0					
Test	St	atistic	p Val	lue	
Student's t	t 5722.199		Pr > t	<.0001	
Sign	М	479348.5	Pr >= M	<.0001	
Signed Rank	S	2.298E11	Pr >= S	<.0001	

Tests for Normality					
Test	Statistic p Value			ue	
Kolmogorov-Smirnov	D	0.065737	Pr > D	<0.0100	
Cramer-von Mises	W-Sq	548.8601	Pr > W-Sq	<0.0050	
Anderson-Darling	A-Sq	3884.935	Pr > A-Sq	<0.0050	

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.32095			
99%	7.30953			
95%	7.08369			
90%	6.84050			
75% Q3	6.43948			
50% Median	5.59508			
25% Q1	4.89515			
10%	4.35773			
5%	4.01241			
1%	3.39786			
0% Min	2.47064			

Freq: WeightD

Geography=Alberta

Extreme Observations						
Lowest			Highest			
Value	Freq	Obs	Value	Freq	Obs	
2.47064	3940	86	7.18200	4355	49	
3.08374	3250	43	7.27414	2361	20	
3.39786	5372	151	7.30361	5912	110	
3.43108	3237	35	7.30953	8411	40	
3.52812	3248	32	7.32095	5649	80	

The UNIVARIATE Procedure Variable: LOG_FD806

Freq: WeightD

Geography=Atlantic provinces

Moments					
N	629768	Sum Weights 629			
Mean	5.61588116	Sum Observations	3536702.25		
Std Deviation	0.94909276	Variance	0.90077707		
Skewness	-0.4111892	Kurtosis	-0.3415584		
Uncorrected SS	20428979.2	Corrected SS	567279.674		
Coeff Variation	16.9001575	Std Error Mean	0.00119596		

Freq: WeightD

Geography=Atlantic provinces

	Basic Statistical Measures					
Location		Variability				
Mean	5.615881	Std Deviation	0.94909			
Median	5.738828	Variance	0.90078			
Mode	4.333099	Range	4.28184			
		Interquartile Range	1.37771			

Tests for Location: Mu0=0					
Test	St	atistic	p Val	lue	
Student's t	t	4695.691	Pr > t	<.0001	
Sign	М	314884	Pr >= M	<.0001	
Signed Rank	S	9.915E10	Pr >= S	<.0001	

Tests for Normality					
Test	Statistic p Value				
Kolmogorov-Smirnov	D	0.052851	Pr > D	<0.0100	
Cramer-von Mises	W-Sq	365.0699	Pr > W-Sq	<0.0050	
Anderson-Darling	A-Sq	2531.387	Pr > A-Sq	<0.0050	

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.35361			
99%	7.32214			
95%	7.03277			
90%	6.73863			

Freq: WeightD

Geography=Atlantic provinces

Quantiles (Definition 5)					
Level Quantil					
75% Q3	6.35141				
50% Median	5.73883				
25% Q1	4.97369				
10%	4.32665				
5%	3.89182				
1%	3.25810				
0% Min	3.07177				

Extreme Observations							
Le	owest		Hi	ghest			
Value Freq Obs Value Freq					Obs		
3.07177	840	797	7.34003	776	400		
3.07177	347	659	7.34357	669	308		
3.15274	132	361	7.35014	1360	487		
3.21165	209	863	7.35109	759	881		
3.21727	186	213	7.35361	491	375		

Freq: WeightD

Geography=British Columbia

Moments						
N	969907	969907 Sum Weights 969				
Mean	5.49258536	Sum Observations	5327296.99			
Std Deviation	0.88839624	Variance	0.78924789			
Skewness	-0.2858383	Kurtosis	-0.3704013			
Uncorrected SS	30026129.7	Corrected SS	765496.26			
Coeff Variation	16.174464	Std Error Mean	0.00090207			

Basic Statistical Measures						
Location Variability						
Mean	5.492585	Std Deviation	0.88840			
Median	5.553541	Variance	0.78925			
Mode	6.246301	Range	4.14116			
		Interquartile Range	1.13596			

Tests for Location: Mu0=0						
Test	Statistic p Value					
Student's t	t	6088.848	Pr > t	<.0001		
Sign	M 484953.5		Pr >= M	<.0001		
Signed Rank	s	2.352E11	Pr >= S	<.0001		

Freq: WeightD

Geography=British Columbia

Tests for Normality							
Test	Test Statistic p Value						
Kolmogorov-Smirnov	D	0.051237	Pr > D	<0.0100			
Cramer-von Mises	W-Sq	491.1901	Pr > W-Sq	<0.0050			
Anderson-Darling	A-Sq	3618.11	Pr > A-Sq	<0.0050			

Quantiles (Definition 5)					
Level	Quantile				
100% Max	7.35843				
99%	7.20510				
95%	7.04319				
90%	6.60932				
75% Q3	6.10216				
50% Median	5.55354				
25% Q1	4.96620				
10%	4.14804				
5%	3.87867				
1%	3.36246				
0% Min	3.21727				

Freq: WeightD

Geography=British Columbia

Extreme Observations							
L	owest		н	ighest			
Value	Freq	Obs	Value	Freq	Obs		
3.21727	1553	1060	7.10462	9539	1053		
3.33220	6907	1089	7.19090	3791	946		
3.36246	3183	932	7.20510	5467	1076		
3.44042	4303	1064	7.21127	1683	1094		
3.63627	3779	1007	7.35843	4907	987		

The UNIVARIATE Procedure Variable: LOG_FD806

Freq: WeightD

Geography=Manitoba

Moments							
N	299857	299857 Sum Weights 299					
Mean	5.39213173	Sum Observations	1616868.45				
Std Deviation	0.92721768	Variance 0.85973					
Skewness	-0.2306575	Kurtosis	-0.4166667				
Uncorrected SS	8976163.64	Corrected SS	257795.988				
Coeff Variation	17.1957535	Std Error Mean	0.00169326				

Freq: WeightD

Geography=Manitoba

	Basic Statistical Measures						
Loc	Location Variability						
Mean	5.392132	Std Deviation	0.92722				
Median	5.451725	Variance	0.85973				
Mode	4.481872	Range	4.62243				
		Interquartile Range	1.31947				

Tests for Location: Mu0=0							
Test	Statistic p Value						
Student's t	t	3184.461	Pr > t	<.0001			
Sign	M 149928.5		Pr >= M	<.0001			
Signed Rank	s	2.248E10	Pr >= S	<.0001			

Tests for Normality							
Test Statistic p Value							
Kolmogorov-Smirnov	D	0.037604	Pr > D	<0.0100			
Cramer-von Mises	W-Sq	95.38586	Pr > W-Sq	<0.0050			
Anderson-Darling	A-Sq	635.7575	Pr > A-Sq	<0.0050			

Quantiles (Definition 5)			
Level	Quantile		
100% Max	7.33580		
99%	7.16667		
95%	6.82778		
90%	6.63637		

Freq: WeightD

Geography=Manitoba

Quantiles (Definition 5)				
Level	Quantile			
75% Q3	6.07496			
50% Median	5.45172			
25% Q1	4.75548			
10%	4.16200			
5%	3.81110			
1%	3.25810			
0% Min	2.71337			

Extreme Observations					
L	owest		Highest		
Value	Freq	Obs	Value	Freq	Obs
2.71337	1520	1314	7.08978	1941	1131
3.10727	381	1270	7.13096	1147	1256
3.25810	2580	1228	7.16667	1520	1148
3.26805	1551	1243	7.27891	505	1267
3.38608	846	1252	7.33580	1566	1130

Freq: WeightD

Geography=Ontario

Moments					
N	2882375	Sum Weights	2882375		
Mean	5.30040293	Sum Observations	15277748.9		
Std Deviation	0.93412537	Variance	0.87259021		
Skewness	0.02419881	Kurtosis	-0.5652113		
Uncorrected SS	83493356.2	Corrected SS	2515131.33		
Coeff Variation	17.6236672	Std Error Mean	0.00055021		

Basic Statistical Measures					
Loc	Location Variability				
Mean	5.300403	Std Deviation	0.93413		
Median	5.267394	Variance	0.87259		
Mode	4.678792	Range	4.64785		
		Interquartile Range	1.32530		

Tests for Location: Mu0=0					
Test	Statistic p Value				
Student's t	t 9633.386		Pr > t	<.0001	
Sign	М	1441188	Pr >= M	<.0001	
Signed Rank	s	2.077E12	Pr >= S	<.0001	

Freq: WeightD

Geography=Ontario

Tests for Normality					
Test	Statistic p Value				
Kolmogorov-Smirnov	D	0.039775	Pr > D	<0.0100	
Cramer-von Mises	W-Sq	847.0641	Pr > W-Sq	<0.0050	
Anderson-Darling	A-Sq	7154.464	Pr > A-Sq	<0.0050	

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.29261			
99%	7.24642			
95%	6.89946			
90%	6.58360			
75% Q3	5.94694			
50% Median	5.26739			
25% Q1	4.62163			
10%	4.01530			
5%	3.90258			
1%	3.30689			
0% Min	2.64476			

Freq: WeightD

Geography=Ontario

Extreme Observations					
L	.owest		Highest		
Value	Freq	Obs	Value	Freq	Obs
2.64476	15313	1596	7.17072	18757	1363
3.30689	15881	1579	7.23335	5785	1355
3.37143	1772	1464	7.24642	6213	1341
3.58129	14896	1552	7.28909	12702	1432
3.58574	6267	1572	7.29261	12196	1415

The UNIVARIATE Procedure Variable: LOG_FD806

Freq: WeightD

Geography=Quebec

Moments						
N	2104687	Sum Weights	2104687			
Mean	5.23877013	Sum Observations	11025971.4			
Std Deviation	0.93216345	Variance	0.86892871			
Skewness	-0.1950948	Kurtosis	-0.6476333			
Uncorrected SS	59591351.7	Corrected SS	1828822.08			
Coeff Variation	17.7935552	Std Error Mean	0.00064254			

Freq: WeightD

Geography=Quebec

	Basic Statistical Measures					
Loc	Location Variability					
Mean	5.238770	Std Deviation	0.93216			
Median	5.292802	Variance	0.86893			
Mode	5.754000	Range	4.22920			
		Interquartile Range	1.44218			

Tests for Location: Mu0=0					
Test	Statistic p Value				
Student's t	t 8153.255		Pr > t	<.0001	
Sign	M 1052344		Pr >= M	<.0001	
Signed Rank	S	1.107E12	Pr >= S	<.0001	

Tests for Normality						
Test	Statistic p Value					
Kolmogorov-Smirnov	D	0.041222	Pr > D	<0.0100		
Cramer-von Mises	W-Sq	875.923	Pr > W-Sq	<0.0050		
Anderson-Darling	A-Sq	6349.62	Pr > A-Sq	<0.0050		

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.25157			
99%	7.09923			
95%	6.65562			
90%	6.42652			

Freq: WeightD

Geography=Quebec

Quantiles (Definition 5)				
Level Quanti				
75% Q3	5.96343			
50% Median	5.29280			
25% Q1	4.52124			
10%	3.96119			
5%	3.53777			
1%	3.27790			
0% Min	3.02237			

Extreme Observations					
Lowest			F	lighest	
Value	Freq	Obs	Value	Freq	Obs
3.02237	4062	1878	6.99027	10294	1908
3.03495	8060	1899	6.99866	5455	1907
3.05777	7665	1906	7.09923	7410	1893
3.27790	16121	1864	7.17749	11846	1787
3.30689	7416	1635	7.25157	2957	1686

Freq: WeightD

Geography=Saskatchewan

Moments					
N	271874	271874 Sum Weights 2			
Mean	5.63878298	Sum Observations	1533038.48		
Std Deviation	0.88073409	Variance	0.77569254		
Skewness	-0.2887928	Kurtosis	-0.5745512		
Uncorrected SS	Uncorrected SS 8855361.17 Corrected SS		210889.857		
Coeff Variation	15.619223	Std Error Mean	0.00168912		

Basic Statistical Measures					
Location Variability					
Mean	5.638783	Std Deviation	0.88073		
Median	5.658018	Variance	0.77569		
Mode	3.814631	Range	4.17511		
		Interquartile Range	1.31152		

Tests for Location: Mu0=0						
Test	Statistic p Value					
Student's t	t 3338.293		Pr > t	<.0001		
Sign	М	135937	Pr >= M	<.0001		
Signed Rank	s	1.848E10	Pr >= S	<.0001		

Freq: WeightD

Geography=Saskatchewan

Tests for Normality					
Test	Statistic p Value				
Kolmogorov-Smirnov	D	0.055935	Pr > D	<0.0100	
Cramer-von Mises	W-Sq	175.6252	Pr > W-Sq	<0.0050	
Anderson-Darling	A-Sq	1242.856	Pr > A-Sq	<0.0050	

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.25885			
99%	7.18062			
95%	6.97271			
90%	6.76542			
75% Q3	6.32522			
50% Median	5.65802			
25% Q1	5.01370			
10%	4.50347			
5%	4.15685			
1%	3.58269			
0% Min	3.08374			

Freq: WeightD

Geography=Saskatchewan

Extreme Observations						
Lowest			Highest			
Value	Freq	Obs	Value	Freq	Obs	
3.08374	1372	2065	7.14419	1518	2141	
3.34850	523	2010	7.18054	888	1925	
3.58269	838	2152	7.18062	1691	2027	
3.70549	629	2082	7.20541	1060	1976	
3.72014	497	1953	7.25885	1377	1981	

The UNIVARIATE Procedure Variable: LOG_FD806

Freq: WeightD

Geography=Territorial capitals

Moments						
N	11711	11711 Sum Weights				
Mean	5.71392282	Sum Observations	66915.7502			
Std Deviation	0.94859878	Variance	0.89983964			
Skewness	-0.2810241	Kurtosis	-0.9449775			
Uncorrected SS	392888.554	Corrected SS	10537.1222			
Coeff Variation	16.6015329	Std Error Mean	0.00876568			

Freq: WeightD

Geography=Territorial capitals

	Basic Statistical Measures					
Location Variability						
Mean	5.713923	Std Deviation	0.94860			
Median	5.759879	Variance	0.89984			
Mode	4.574092	Range	3.78783			
		Interquartile Range	1.46558			

Tests for Location: Mu0=0						
Test	Statistic p Value					
Student's t	t	651.8517	Pr > t	<.0001		
Sign	M 5855.5		Pr >= M	<.0001		
Signed Rank	s	34289808	Pr >= S	<.0001		

Tests for Normality							
Test Statistic p Value							
Kolmogorov-Smirnov	D	0.082104	Pr > D	<0.0100			
Cramer-von Mises	W-Sq	16.42393	Pr > W-Sq	<0.0050			
Anderson-Darling	A-Sq	115.1312	Pr > A-Sq	<0.0050			

Quantiles (Definition 5)				
Level	Quantile			
100% Max	7.36074			
99%	7.30971			
95%	7.08256			
90%	6.93674			

Freq: WeightD

Geography=Territorial capitals

Quantiles (Definition 5)					
Level	Quantile				
75% Q3	6.45063				
50% Median	5.75988				
25% Q1	4.98504				
10%	4.25135				
5%	4.13772				
1%	3.78328				
0% Min	3.57291				

Extreme Observations							
L	owest		Highest				
Value Freq Obs			Value	Freq	Obs		
3.57291	18	2172	7.27783	46	2311		
3.78328	115	2270	7.28773	45	2168		
3.85757	157	2298	7.30971	58	2171		
3.91562	58	2310	7.32818	59	2180		
3.94119	33	2299	7.36074	43	2163		

Null hypothesis: equal variances

a.If variances are equal, then a pooled t-test is appropriate

b.lf variances are unequal, then a Satterthwaite (also known as Welch's) test is appropriate

The GLM Procedure

	Class Level Information						
Class Levels Values							
Prov	8	Alberta Atlantic provinces British Columbia Manitoba Ontario Quebec Saskatchewan Territorial capitals					

Number of Observations Read	2327
Number of Observations Used	2327
Sum of Frequencies Read	8128876
Sum of Frequencies Used	8128876

The GLM Procedure

Dependent Variable: LOG_FD806

Frequency: WeightD

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	7	172580.537	24654.362	28485.7	<.0001
Error	8.13E6	7035530.477	0.865		
Corrected Total	8.13E6	7208111.014			

R-Square Coeff Var		Root MSE	LOG_FD806 Mean
0.023943	17.28520	0.930322	5.382190

Source	DF	Type I SS	Mean Square	F Value	Pr > F
Prov	7	172580.5368	24654.3624	28485.7	<.0001

Source	DF	Type III SS	Mean Square	F Value	Pr > F
Prov	7	172580.5368	24654.3624	28485.7	<.0001

The GLM Procedure

Levene's Test for Homogeneity of LOG_FD806 Variance ANOVA of Absolute Deviations from Group Means							
Source	F Value	Pr > F					
Prov	7	4166.6	595.2	2107.67	<.0001		
Error	8.13E6	2295703	0.2824				

Welch's ANOVA for LOG_FD806					
Source	DF	F Value	Pr > F		
Prov	7.0000	28314.2	<.0001		
Error	220971				

The GLM Procedure


		LOG_FD806		
Level of Prov	N	Mean	Std Dev	
Alberta	958697	5.59782167	0.95784837	
Atlantic provinces	629768	5.61588116	0.94909276	
British Columbia	969907	5.49258536	0.88839624	
Manitoba	299857	5.39213173	0.92721768	
Ontario	2882375	5.30040293	0.93412537	
Quebec	2104687	5.23877013	0.93216345	
Saskatchewan	271874	5.63878298	0.88073409	
Territorial capitals	11711	5.71392282	0.94859878	

The GLM Procedure Least Squares Means Adjustment for Multiple Comparisons: Tukey-Kramer

Prov	LOG_FD806 LSMEAN	LSMEAN Number	
Alberta	5.59782167	1	
Atlantic provinces	5.61588116	2	
British Columbia	5.49258536	3	
Manitoba	5.39213173	4	
Ontario	5.30040293	5	
Quebec	5.23877013	6	
Saskatchewan	5.63878298	7	
Territorial capitals	5.71392282	8	

Least Squares Means for effect Prov Pr > t for H0: LSMean(i)=LSMean(j)									
	Dependent Variable: LOG_FD806								
i/j	1	2	3	4	5	6	7	8	
1		<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	
2	<.0001		<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	
3	<.0001	<.0001		<.0001	<.0001	<.0001	<.0001	<.0001	
4	<.0001	<.0001	<.0001		<.0001	<.0001	<.0001	<.0001	
5	<.0001	<.0001	<.0001	<.0001		<.0001	<.0001	<.0001	
6	<.0001	<.0001	<.0001	<.0001	<.0001		<.0001	<.0001	
7	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001		<.0001	
8	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001		

The GLM Procedure Least Squares Means Adjustment for Multiple Comparisons: Tukey-Kramer



The GLM Procedure Least Squares Means Adjustment for Multiple Comparisons: Tukey-Kramer

