



## Practice quiz on Probability Concepts

Practice Quiz • 25 min



### Congratulations! You passed!

TO PASS 75% or higher

Keep Learning

GRADE

100%

## Practice quiz on Probability Concepts

TOTAL POINTS 9

1. If  $x = \text{"It is raining,"}$  what is  $\sim (\sim x)$ ?

1 / 1 point

- ☐ "It is never raining"
- ☐ "It is always raining"
- ☒ "It is raining"
- ☐ "It is not raining"



Correct

The second negation cancels out the first one.

Similarly  $\sim (\sim (\sim x)) = \sim x$

2. If the statement "I am 25 years old" is assigned probability 0, what probability is assigned to the statement "I am not 25 years old"?

1 / 1 point

- ☐ 0
- ☐ Unknown
- ☐ -1
- ☒ 1



Correct

It is always the case that  $p(x) + p(\sim x) = 1$ .

3. If I assign to the statement  $x = \text{"it will rain today"}$  a probability of  $p(x) = 0.35$ , what probability must I assign to the statement "it will not rain today?"

1 / 1 point

- ☐ 0

- ☐ .35
- ☒ .65
- ☐ .5



Correct

$$p(x) + p(\sim x) = 1$$

4. Is the following collection of statements a probability distribution?

1 / 1 point

1. I own a Toyota pickup truck
2. I do not own a Toyota pickup truck
3. I own a non-Toyota pickup truck
4. I do not own a non-Toyota pickup truck

- ☒ No
- ☐ Yes



Correct

The statements are not *exclusive*: 1 and 4 could both be true, 2 and 3 could both be true, 2 and 4 could both be true, and even (1) and (3) could both be true (if I owned more than one pickup truck).

5. I don't know what it means to be "ingenuous." What probability would I assign to the statement, "I am ingenuous OR I am not ingenuous"?

1 / 1 point

- ☐ -1
- ☒ 1
- ☐ 0
- ☐ .5



Correct

It is always the case, regardless of the content of the statement  $x$ , that  $p(x \text{ or } \sim x) = 1$ .

$$\sim x) = 1$$

6. A friend of mine circumscribes a circle inside a square, so that the diameter of the circle and the edge of the square are the same length. He asks me to close my eyes and pick a point at random inside the square. He says the probability that my point will also be inside the circle is  $\frac{\pi}{4}$

1 / 1 point

Is this correct?

☒ Yes

☐ No



Correct

Probabilities can be any real number between 0 and 1. They do not need to be rational numbers – a numerator that is a transcendental number like  $\pi$  is acceptable.

Note that the correct probability does not depend on the length  $r$  of the circle's radius. For a circle with any radius  $r$  to be circumscribed inside a square, the square must have sides each of length  $2r$ . The area of the circle is  $\pi r^2$  and the area of the square is  $(2r)^2 = 4r^2$ . The probability of landing in a circle of area  $\pi r^2$  when it is known that one is in the area of the square is equal to the ratio of the area of the circle to the area of the square in which it is circumscribed, or  $\pi r^2 / 4r^2$ , which equals  $\pi/4$ .

7. The probability of drawing a straight flush (including a Royal Flush) in a five-card poker hand is 0.0000153908

1 / 1 point

What is the probability of **not** drawing a straight flush?

☐ .9999745688

☒ .9999846092

☐ .9996582672

☐ .9967253809



Correct

$$p(\sim x) = 1 - p(x)$$

8. What is the probability that a fair, six-sided die will come up with a prime number? (Recall that prime numbers are positive integers other than 1 that are divisible only by themselves and 1)

1 / 1 point

- ☐  $\frac{1}{6}$
- ☐  $\frac{1}{3}$
- ☐  $\frac{2}{3}$
- ☒  $\frac{1}{2}$

✓ Correct

The faces with 2, 3 and 5 satisfy the condition – which makes 3 relevant outcomes out of the “universe” of 6 outcomes  $= \frac{3}{6} = \frac{1}{2}$

9. The joint probability  $p$  (the die will come up 5, the next card will be a heart) is equal to the joint probability:

1 / 1 point

- ☒  $p$  (the next card will be a heart, the die will come up 5)
- ☐  $p$  (the next card will **not** come up 5, the next card will be a heart)
- ☐  $p$  (the die will **not** come up 5, the next card will **not** be a heart)
- ☐  $p$  (the next card will be a heart, the die will **not** come up 5)

✓ Correct

In joint probabilities, the order does not change the probability:  $p(A, B) = p(B, A)$